Volumetric and viscometric behavior of the binary system: (Hexan-1-ol + p -Xylene)

著者	Das Kamalendra N., Habibullah M., Rahman Ismail M. M., Hasegawa Hiroshi, Uddin M. Ashraf, Saifuddin Khaled
journal or publication title	Journal of Chemical and Engineering Data
volume	55
number	11
page range	5311-5313
year	2010-11-11
URL	http://hdl.handle.net/2297/26260

doi: 10.1021/je100390b

Volumetric and Viscometric Behavior of the Binary System: (Hexan-1-ol + *p*-Xylene)

Kamalendra N. Das^{a,}, M. Habibullah^a, Ismail M. M. Rahman^{a, b,*}, Hiroshi Hasegawa^b, M. Ashraf
Uddin^a, Khaled Saifuddin^a*

^aDepartment of Chemistry, Faculty of Science, University of Chittagong, Chittagong-4331, Bangladesh

^bGraduate School of Natural Science and Technology, Kanazawa University, Kakuma, Kanazawa 920-
1192, Japan

*Author(s) for correspondence.

E-mail: kamalctgu@hotmail.com (K. N. Das); I.M.M.Rahman@gmail.com (I. M. M. Rahman).

Tel/ Fax: +81-76-234-4792

1 **Abstract**

2 Densities and viscosities for the binary mixtures of hexan-1-ol with *p*-xylene have been measured at a
3 number of mole fractions at $T = (303.15, 313.15, \text{ and } 323.15)$ K. The excess molar volumes and
4 viscosity deviations have been calculated from the experimental results and have been fitted to the
5 Redlich-Kister polynomial equation.

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

1 Introduction

2 Densities and viscosities of multi-component solvent mixtures are required for the process design
3 involving heat transfer, mass transfer, fluidity and so forth as well as for the development of our
4 understanding of molecular interactions in such systems^{1,2}. In this work, which is a part of our ongoing
5 research on the accumulation of the binary physical property data of organic liquid mixtures³⁻⁷, we are
6 reporting the density and viscosity data of the binary mixtures of hexan-1-ol with *p*-xylene at $T =$
7 (303.15, 313.15, and 323.15) K. Among the two components of the studied binary mixtures, hexan-1-ol
8 is an aliphatic polar solvent with strong self-associating nature via hydrogen bonding and Van der
9 Waals interactions while *p*-xylene is a non-associating solvent. Therefore, a favorable interaction with
10 hydroxyl group of hexan-1-ol is possible due to the presence of aromatic rings with delocalized π -
11 electrons in *p*-xylene⁸, and the experimental results will allow us to study the interactions. Excess molar
12 volumes and viscosity deviations for the binary systems have been calculated from the experimental
13 results and fitted to the Redlich–Kister polynomial equation⁹ to estimate the binary coefficients and the
14 standard deviations. The experimental and the derived data have been interpreted using some previous
15 observations for the similar binary systems. The results are compared to those of binary systems of
16 hexan-1-ol with *o*- and *m*-xylenes^{6,7}, which were previously reported.

18 Experimental Section

19 Hexan-1-ol and *p*-xylene from Merck-Schuchard, Germany (0.98 mass fraction purity) were used
20 without further treatment. The results of the measurements of their densities and viscosities at $T =$
21 (303.15, 313.15, and 323.15) K were in satisfactory agreement with the published values (Table 1).
22 Hence, the materials were taken to be reasonably pure.

23 All binary mixtures were prepared by mass on an analytical balance (B 204-S, Mettler Toledo,
24 Switzerland) with uncertainty of ± 0.0001 g and operating in a dry box. Mixtures were completely
25 miscible over the whole composition range. Caution was taken to prevent evaporation of the samples
26 after preparation. The mole fraction of each mixture was obtained to an uncertainty of $\pm 1.0 \times 10^{-4}$. A

transparent glass-walled thermostatic water bath was used. The bath temperature was controlled by the Thermo Haake DC10 controller (Thermo Fisher Scientific, MA, USA), and was monitored with a minimum-maximum thermometer (Brannan Thermometers, Cumberland, UK). The uncertainty in the temperature during the measurements was ± 0.05 K. A 10 mL bi-capillary pycnometer and an A-type Ostwald viscometer, previously calibrated with redistilled water, were used to measure the densities and viscosities respectively. The average uncertainties in the measured densities and viscosities were of ± 0.0004 g·cm⁻³ and ± 0.005 mPa·s, respectively. For all mixture compositions and pure solvents, triplicate measurements were performed, and the average of these values was considered in all calculations.

Results and Discussion

Binary compositions of the mixtures of hexan-1-ol with *p*-xylene, the experimental densities and viscosities, excess molar volumes, V_m^E and the viscosity deviations, $\Delta\eta$ at $T = (303.15, 313.15, \text{ and } 323.15)$ K are summarized in Table 2. The V_m^E and $\Delta\eta$ values have been calculated using the following relations:

$$V_m^E = \left[(x_1 M_1 + x_2 M_2) / \rho - \{ (x_1 M_1) / \rho_1 + (x_2 M_2) / \rho_2 \} \right] \quad (1)$$

$$\Delta\eta = \eta - \exp(x_1 \ln \eta_1 + x_2 \ln \eta_2) \quad (2)$$

where x_1 , M_1 , ρ_1 and η_1 represent respectively mole fraction, molar mass, density and viscosity of hexan-1-ol, and x_2 , M_2 , ρ_2 and η_2 are the corresponding quantities of *p*-xylene. ρ and η are the density and viscosity of the binary mixtures. The uncertainties in V_m^E and $\Delta\eta$ are found to be within ± 0.0001 g·cm⁻³ and ± 0.001 mPa·s, respectively.

The V_m^E profiles for the binary mixtures varying with the mole fraction of hexan-1-ol (x_1) at $T = (303.15, 313.15 \text{ and } 323.15)$ K is observed positive over the whole range of composition (Fig. 1). The V_m^E values increase sharply, reach a maximum and then decrease gradually with the increase of hexan-1-ol mole fraction. The maxima are at about 0.3000 mole fraction. The magnitude of V_m^E values

1 increases with the increase of temperature. The $\delta V_m^E/\delta T$ values for all temperatures are positive. The
 2 sign of V_m^E depends upon the resultant of the factors contributing to the expansion of volume and the
 3 factors contributing to the contraction of volume. The expansion of volume as indicated by the positive
 4 V_m^E values can be attributed to the fact that the total positive contributing factors outweigh the factors
 5 responsible for negative contributing effects indicating weak interaction between the two components.
 6 Hexan-1-ol, like all other alcohols, has strongly associated molecules through H-bonding and added to a
 7 non-associating solvent, *p*-xylene to prepare the binary mixtures. The initial sharp increase in V_m^E
 8 values in the hexan-1-ol low concentration region may be due to the maximum or complete
 9 disintegration of associated molecules of hexan-1-ol in the solution containing the non-polar component
 10 *p*-xylene. At hexan-1-ol rich region of the mixtures, maximum concentration of the alcohol remain in
 11 the associated state facilitating the partial or full accommodation of the *p*-xylene into the network of
 12 alcohol. These may be the cause of slow gradual decrease of V_m^E values at hexan-1-ol rich region.
 13 Lower V_m^E values obtained for hexan-1-ol with *p*-xylene than those reported for hexan-1-ol with *o*- and
 14 *m*-xylenes^{6, 7} indicate the probability of weaker interactions between the delocalized π -electrons of the
 15 *p*-xylene ring and the –OH group of hexan-1-ol beside the disruption of H-bonding.

16 The $\Delta\eta$ values for the binary mixtures of hexan-1-ol with *p*-xylene are negative throughout the whole
 17 range of mole fraction. Figure 2 illustrates the trend of $\Delta\eta$ values as a function of the binary
 18 compositions at $T = (303.15, 313.15, \text{ and } 323.15)$ K with a minima at around 0.6000 mole fraction of
 19 alcohol. Molecular interactions between the components of the mixture determine the viscometric
 20 behavior of a solvent mixture. The negative $\Delta\eta$ values observed for the hexan-1-ol and *p*-xylene system
 21 can be described by the dominance of dispersion force compared to the induced dipole–dipole
 22 interactions which correspond to a system containing an associated component¹⁰. It is anticipated that
 23 the associated structure of hexan-1-ol via H-bonding undergoes disruption when added to *p*-xylene
 24 forming monomeric, oligomeric or small scaled multimeric species, which experience much less
 25 resistance to flow, resulting in the negative viscosity deviations. An increase of temperature leads to an
 26 increase in $\Delta\eta$, i.e., $\delta\eta/\delta T$ is positive. Skewed curves with minima in the hexan-1-ol high concentration

region may be due to the so called interstitial accommodation effect. The *p*-xylene molecules occupy the void spaces within the hexan-1-ol network structures which may result more compact and well-shaped flowing species with favorable geometric fitting. Therefore, the frictional resistance is reduced remarkably, and $\Delta\eta$ decreases sharply. The $\Delta\eta$ values for the hexan-1-ol and *p*-xylene binary system have higher magnitudes than those observed for hexan-1-ol with *o*- and *m*-xylenes^{6, 7}.

The composition dependence of V_m^E or $\Delta\eta$ can be represented by a Redlich–Kister⁹ type equation:

$$\delta = x_1 x_2 \sum_{i=0}^n A_i (1 - 2x_1)^i \quad (3)$$

where δ refers to V_m^E or $\Delta\eta$, and x_1 and x_2 are the mole fractions of hexan-1-ol and *p*-xylene. The variables A_i are the equation coefficients, which are obtained by fitting the equation to the experimental values with a least-squares regression method (Table 3). Data processing and analyses are performed with LAB Fit¹¹ and LSM¹² curve-fitting program. Sma4 for windows¹³ are used for graphical representation of data.

Excess molar volumes (V_m^E) for the hexan-1-ol and *p*-xylene system were reported by Rodríguez-Núñez *et al.*¹⁴ at 298.15 K. The V_m^E values observed at 298.15 K were relatively small showing negative-positive trend in the profile. Whereas, the trend of V_m^E profiles are positive over the entire range of binary compositions at $T = (303.15, 313.15 \text{ and } 323.15) \text{ K}$.

There are reports on the volumetric behavior of *p*-xylene with different alkanols¹⁵⁻¹⁹ other than hexan-1-ol. However, no literature data are available with which to compare the present results for hexan-1-ol with *p*-xylene at $T = (303.15, 313.15 \text{ and } 323.15) \text{ K}$.

1 **Acknowledgment**

2 One of the authors, Ismail M. M. Rahman, acknowledges the support from the Japanese Government
3 (Monbukagakusho: MEXT) Scholarship program that facilitated this research.

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

Literature Cited:

- (1) Rowlinson, J. S.; Swinton, F. L., *Liquids and Liquid Mixtures*. Butterworth Scientific: London, 1982.
- (2) Reichardt, C., *Solvents and Solvent Effects in Organic Chemistry*. 3rd ed.; Wiley-VCH: Weinheim, 2003.
- (3) Adhikhari, A. B.; Rahman, I. M. M.; Uddin, M. A.; Hasegawa, H.; Majid, M. A., Volumetric behavior of the binary mixtures of methyl ethyl ketone with *n*-hexane, cyclohexane, and benzene at $T = (303.15, 313.15, \text{ and } 323.15) \text{ K}$. *J. Chem. Eng. Data* **2009**, *54*, 1138-1141.
- (4) Chowdhury, M. A.; Majid, M. A.; Saleh, M. A., Volumetric and viscometric behaviour of binary systems: (1-hexanol + hydrocarbons). *J. Chem. Thermodyn.* **2001**, *33*, 347-360
- (5) Saleh, M. A.; Habibullah, M.; Ahmed, M. S.; Uddin, M. A.; Uddin, S. M. H.; Uddin, M. A.; Khan, F. M., Excess molar volumes and viscosities of some alkanols with cumene. *Phys. Chem. Liq.* **2006**, *44*, 31-43.
- (6) Das, K. N.; Habibullah, M.; Rahman, I. M. M.; Hasegawa, H.; Uddin, M. A.; Saifuddin, K., Thermodynamic properties of the binary mixture of hexan-1-ol with *m*-xylene at $T = (303.15, 313.15, \text{ and } 323.15) \text{ K}$. *J. Chem. Eng. Data* **2009**, *54*, 3300-3302.
- (7) Das, K. N.; Habibullah, M.; Uddin, K. S.; Barua, P., The thermodynamic excess properties of binary mixtures: I. hexanol-1+*o*-xylene. *J. Solution Chem.* **2006**, *35*, 1017-1028.
- (8) Barker, J. A.; Smith, F., Statistical thermodynamics of associated solutions. *J. Chem. Phys.* **1954**, *22*, 375-380.
- (9) Redlich, O.; Kister, A. T., Algebraic representation of thermodynamic properties and the classification of solutions. *Ind. Eng. Chem.* **1948**, *40*, 345-348.
- (10) Hawrylak, B.; Andrezyk, S.; Gabriel, C. E.; Gracie, K.; Palepu, R., Viscosity, surface tension, and refractive index measurements of mixtures of isomeric butanediols with water. *J. Solution Chem.* **1998**, *27*, 827-841.
- (11) *LAB Fit Curve Fitting Software (Nonlinear Regression and Treatment of Data Program)*; v. 7.2.47; online, available from - www.labfit.net (shareware); Wilton and Cleide Pereira da Silva - DF/CCT/UFPB: Brazil, 1999–2010.

- 1 (12) *LSM (Least Squares Method Curve Fitting Program)*; v. 2.2; online, available from -
2 <http://www.prz.edu.pl/~janand/> (freeware); J. A. Mamczur: Poland, 1998–2010.
- 3 (13) *Sma4 for Windows*; v. 1.48 E; online, available from -
4 <http://hp.vector.co.jp/authors/VA002995/soft/sma4win.htm> (shareware); Toshio Suzuki: Japan,
5 1994–1998.
- 6 (14) Rodriguez-Nuñez, E.; Paz-Andrade, M. I.; Bravo, R.; Ortega, J., Thermodynamic properties of (a
7 xylene + an *n*-alkanol) III. Excess molar volumes at 298.15 K for 1-hexanol, 1-heptanol, and 1-
8 octanol. *J. Chem. Thermodyn.* **1985**, *17*, 817-821.
- 9 (15) Yadav, B. L.; Maken, S.; Kalra, K. C.; Singh, K. C., Excess volumes of (an alkanol + an aromatic
10 hydrocarbon) at the temperature 308.15 K. *J. Chem. Thermodyn.* **1993**, *25*, 1345-1350.
- 11 (16) Letcher, T. M.; Koteswari Prasad, A.; Mercer-Chalmers, J.; Radloff, S. E., Excess molar volumes of
12 (a xylene + an alkanol) at 298.15 K. *J. Chem. Thermodyn.* **1990**, *22*, 1175-1179.
- 13 (17) Rodriguez-Nuñez, E.; Paz-Andrade, M. I.; Ortega, J., Thermodynamic properties of (a xylene + an
14 *n*-alkanol) IV. Excess molar volumes at 298.15 K for 1-propanol, 1-butanol, and 1-pentanol. *J.*
15 *Chem. Thermodyn.* **1986**, *18*, 303-308.
- 16 (18) Bhardwaj, U.; Maken, S.; Singh, K. C., Excess volumes of 1-butanol, 2-butanol, 2-methylpropan-1-
17 ol, and 2-methylpropan-2-ol with xylenes at 308.15 K. *J. Chem. Eng. Data* **1996**, *41*, 1043-1045.
- 18 (19) Ouyang, G.; Huang, Z.; Ou, J.; Wu, W.; Kang, B., Excess molar volumes and surface tensions of
19 xylene with 2-propanol or 2-methyl-2-propanol at 298.15 K. *J. Chem. Eng. Data* **2002**, *48*, 195-197.
- 20 (20) Lee, M. J.; Lin, T. K.; Pai, Y. H.; Lin, K. S., Density and viscosity for monoethanolamine + 1-
21 propanol, + 1-hexanol, and + 1-octanol. *J. Chem. Eng. Data* **1997**, *42*, 854-857.
- 22 (21) TRC Thermodynamic Tables–Non-Hydrocarbons. Texas Engineering Experiment Station, Texas A
23 & M University System: 1994.
- 24 (22) Weng, W. L., Viscosities and densities for binary mixtures of anisole with 1-butanol, 1-pentanol, 1-
25 hexanol, 1-heptanol, and 1-octanol. *J. Chem. Eng. Data* **1999**, *44*, 63-66.
- 26 (23) Trenzado, J. L.; Matos, J. S.; Gonzalez, E.; Romano, E.; Caro, M. N., Study on properties derived
27 from densities and viscosities for the ternary systems (methyl pentanoate or methyl heptanoate) plus

octane plus 1-hexanol and their binary subsystems at various temperatures. *J. Chem. Eng. Data* **2003**, *48*, 1004-1014.

(24) Matsuo, S.; Makita, T., Viscosities of six 1-alkanols at temperatures in the range 298–348 K and pressures up to 200 MPa. *Int. J. Thermophys.* **1989**, *10*, 833-843.

(25) Riddick, J. A.; Bunger, W. B.; Sakano, T. K., *Organic Solvents: Physical Properties and Methods of Purification*. 4th ed.; Wiley: New York, 1986.

(26) Serrano, L.; Silva, J. A.; Farelo, F., Densities and viscosities of binary and ternary liquid systems containing xylene. *J. Chem. Eng. Data* **1990**, *35*, 288.

(27) Yang, C.; Xu, W.; Ma, P., Thermodynamic properties of binary mixtures of *p*-xylene with cyclohexane, heptane, octane, and N-methyl-2-pyrrolidone at several temperatures. *J. Chem. Eng. Data* **2004**, *49*, 1794-1801.

(28) Rathnam, M. V.; Mohite, S.; Kumar, M. S., Thermophysical properties of isoamyl acetate or methyl benzoate + hydrocarbon binary mixtures, at (303.15 and 313.15) K. *J. Chem. Eng. Data* **2009**, *54*, 305-309.

(29) Rathnam, M. V.; Mohite, S.; Kumar, M. S. S., Viscosity, density, and refractive index of some (ester plus hydrocarbon) binary mixtures at 303.15 K and 313.15 K. *J. Chem. Eng. Data* **2005**, *50*, 325-329.

(30) George, J.; Sastry, N. V., Densities, excess molar volumes, viscosities, speeds of sound, excess isentropic compressibilities, and relative permittivities for $C_mH_{2m+1}(OCH_2CH_2)_nOH$ ($m = 1$ or 2 or 4 and $n = 1$) + benzene, + toluene, + (*o*-, *m*-, and *p*-) xylenes, + ethylbenzene, and + cyclohexane. *J. Chem. Eng. Data* **2003**, *48*, 977-989.

Table 1. Comparison of Experimental Densities, ρ and Viscosities, η of Pure Liquids with Literature Values at $T = (303.15, 313.15, \text{ and } 323.15) \text{ K}$.

Liquid (s)	T/K	$\rho / \text{g} \cdot \text{cm}^{-3}$	$\eta / \text{mPa} \cdot \text{s}$
------------	-------	--	------------------------------------

		Exp.	Lit.	Ref.	Exp.	Lit.	Ref.
Hexan-1-ol	303.15	0.8119	0.8120	20	3.829	3.84	20
			0.8121	21		3.769	21
			0.8115	22		3.861	22
	313.15	0.8046	0.8046	20, 21	2.879	2.90	20
			0.80428	23		2.89	24
	323.15	0.7975	0.7972	20	2.215	2.23	20
0.7979			21	2.210		24	
<i>p</i> -xylene	303.15	0.8524	0.8523	25	0.573	0.568	25
			0.85239	26		0.5762	26
			0.85235	27			
			0.8521	28			
	313.15	0.8439	0.8435	28	0.512	0.517	28, 29
			0.8440	29			
			0.8436	30			
	323.15	0.8352	0.8349	27	0.461	0.4599	27

Table 2. Compositions, Experimental Densities, ρ , Excess Molar Volumes, V_m^E , Viscosities, η and the Viscosity Deviations, $\Delta\eta$, for the Binary Mixtures of Hexan-1-ol with *p*-Xylene at $T = (303.15, 313.15, \text{ and } 323.15) \text{ K}$.

$T/\text{K} = 303.15$			$T/\text{K} = 313.15$		$T/\text{K} = 323.15$	
Densities and excess molar volumes of hexan-1-ol (1) + <i>p</i> -xylene (2)						
x_1	$\rho / \text{g}\cdot\text{cm}^{-3}$	$V_m^E / \text{cm}^3\cdot\text{mol}^{-1}$	$\rho / \text{g}\cdot\text{cm}^{-3}$	$V_m^E / \text{cm}^3\cdot\text{mol}^{-1}$	$\rho / \text{g}\cdot\text{cm}^{-3}$	$V_m^E / \text{cm}^3\cdot\text{mol}^{-1}$
0.0000	0.8524	0.0000	0.8439	0.0000	0.8352	0.0000
0.0999	0.8478	0.0759	0.8393	0.0961	0.8305	0.1389
0.1999	0.8435	0.1090	0.8351	0.1335	0.8264	0.1873
0.2999	0.8394	0.1140	0.8311	0.1421	0.8226	0.1907
0.4001	0.8354	0.1041	0.8272	0.1355	0.8189	0.1783
0.4997	0.8314	0.0990	0.8233	0.1335	0.8152	0.1702
0.6000	0.8275	0.0756	0.8195	0.1129	0.8116	0.1430
0.7000	0.8236	0.0551	0.8157	0.0951	0.8080	0.1183
0.8000	0.8197	0.0357	0.8120	0.0626	0.8044	0.0942
0.8999	0.8158	0.0179	0.8083	0.0315	0.8009	0.0555
1.0000	0.8119	0.0000	0.8046	0.0000	0.7975	0.0000
Viscosities, and deviations of the viscosity of hexan-1-ol (1) + <i>p</i> -xylene (2)						
x_1	$\eta / \text{mPa}\cdot\text{s}$	$\Delta\eta / \text{mPa}\cdot\text{s}$	$\eta / \text{mPa}\cdot\text{s}$	$\Delta\eta / \text{mPa}\cdot\text{s}$	$\eta / \text{mPa}\cdot\text{s}$	$\Delta\eta / \text{mPa}\cdot\text{s}$
0.0000	0.573	0.000	0.512	0.000	0.461	0.000
0.0999	0.616	−0.077	0.544	−0.064	0.485	−0.054
0.1999	0.693	−0.145	0.605	−0.118	0.533	−0.098
0.2999	0.799	−0.214	0.689	−0.170	0.600	−0.138
0.4001	0.946	−0.279	0.802	−0.220	0.689	−0.175
0.4997	1.148	−0.332	0.956	−0.257	0.807	−0.203
0.6000	1.420	−0.371	1.156	−0.287	0.958	−0.224
0.7000	1.801	−0.365	1.433	−0.282	1.161	−0.222
0.8000	2.323	−0.296	1.806	−0.232	1.433	−0.185
0.8999	3.004	−0.162	2.295	−0.127	1.788	−0.105
1.0000	3.829	0.000	2.879	0.000	2.215	0.000

Table 3. Coefficients, A_i , of Redlich-Kister Equation (Equation 3), Expressing V_m^E ($\text{cm}^3 \cdot \text{mol}^{-1}$) and $\Delta\eta$ ($\text{mPa} \cdot \text{s}$), and Standard Deviation, σ , for the Binary Mixtures of Hexan-1-ol with *p*-Xylene at $T = (303.15, 313.15, \text{ and } 323.15) \text{ K}$.

T/K	A_0	A_1	A_2	A_3	σ
Correlated results for excess volumes of hexan-1-ol (1) + <i>p</i> -xylene (2)					
303.15	0.3746	0.3220	0.2129	0.1423	0.0026
313.15	0.5167	0.2282	0.2833	0.3601	0.0024
323.15	0.6511	0.3702	0.6326	0.3272	0.0037
Correlated results for viscosity deviations of hexan-1-ol (1) + <i>p</i> -xylene (2)					
303.15	-1.3531	1.0008	-0.0358	-0.6278	0.0042
313.15	-1.0506	0.7411	-0.0823	-0.4537	0.0037
323.15	-0.8258	0.5440	-0.1365	-0.2822	0.0025

Figure 1. Excess molar volumes for hexan-1-ol (1) + *p*-xylene (2): ●, 303.15 K; ▲, 313.15 K; ■, 323.15 K

1

2 Figure 2. Viscosity deviations for hexan-1-ol (1) + *p*-xylene (2): ●, 303.15 K; ▲, 313.15 K; ■, 323.15 K

3

4

5