Chemical characteristics of chromian spinel in plutonic rocks: Implications for deep magma processes and discrimination of tectonic setting

著者	Arai Shoji, Okamura Hidenobu, Kadoshima Kazuyuki, Tanaka Chima, Suzuki Kenji, Ishimaru Satoko
著者別表示	荒井 章司, 鈴木 健之, 石丸 聡子
journal or publication title	Island Arc
volume	20
number	1
page range	125-137
year	2011-03-01
URL	http://doi.org/10.24517/00010315

doi: 10.1111/j.1440-1738.2010.00747.x

Chemical characteristics of chromian spinel in plutonic rocks:
implications for deep magma processes and discrimination of
tectonic setting

SHOJI ARAI¹, HIDENOBU OKAMURA¹, KAZUYUKI KADOSHIMA¹, CHIMA TANAKA¹,
KENJI SUZUKI¹ AND SATOKO ISHIMARU¹

¹ *Department of Earth Sciences, Kanazawa University, Kakuma, Kanazawa 920-1192,
Japan (email: ultrasa@kenroku.kanazawa-u.ac.jp)*

* Correspondence

Abstract We summarize chemical characteristics of chromian spinels from ultramafic
to mafic plutonic rocks (lherzolites, harzburgites, dunites, wehrlites, troctolites and
olivine gabbros) with regard to three tectonic settings (mid-ocean ridge, arc and oceanic
hotspot). The chemical range of spinels is distinguishable between the three settings in
terms of Cr# (= Cr/(Cr + Al) atomic ratio) and Ti content. The relationships are almost
parallel with those of chromian spinels in volcanic rocks, but the Ti content is slightly

lower in plutonics than in volcanics at a given tectonic environment. The Cr# of spinels in plutonic rocks is highly diverse; its ranges overlap between the three settings, but extend to higher values (up to 0.8) in arc and oceanic hotspot environments. The Ti content of spinels in plutonics increases, for a given lithology, from the arc to oceanic hotspot settings via mid-ocean ridge on average. This chemical diversity is consistent with that of erupted magmas from the three settings. If we systematically know the chemistry of chromian spinels from a series of plutonic rocks, we can estimate their tectonic environments of formation. The spinel chemistry is especially useful in dunitic rocks, in which chromian spinel is the only discriminating mineral. Applying this, discordant dunites cutting mantle peridotites were possible precipitated from arc-related magmas in the Oman ophiolite, and from an intraplate tholeiite in the Lizard ophiolite, Cornwall.

Key words: ultramafic plutonics, chromian spinel, tectonic setting, Ti content, Cr/(Cr + Al) ratio

Running title: Chromian spinel in plutonic rocks

35

36 INTRODUCTION

37 Chromian spinel is common to ultramafic and related rocks, and is a very good indicator
38 of petrological characteristics of involved magmas (e.g. Irvine 1965, 1967; Dick &
39 Bullen 1984; Roeder 1994; Kamenetsky *et al.* 2001). It has a general formula, $(\text{Mg},$
40 $\text{Fe}^{2+})(\text{Cr}, \text{Al}, \text{Fe}^{3+})_2\text{O}_4$, where Fe^{3+} is only minor in peridotitic rocks. $\text{Cr}/(\text{Cr} + \text{Al})$ atomic
41 ratio (= Cr#) is highly variable and serves as an important petrogenetic indicator for
42 ultramafic and related rocks (Irvine 1967; Dick & Bullen 1984). $\text{Mg}/(\text{Mg} + \text{Fe}^{2+})$ atomic
43 ratio (= Mg#) varies inversely with the Cr# in chromian spinel (e.g. Irvine 1967). Small
44 amounts of Ti are possibly incorporated as Fe_2TiO_4 (= ulvospinel component) in
45 chromian spinel. Arai (1992) summarized the chemistry of chromian spinel in volcanic
46 rocks (or magmas) as a potential indicator of magma chemistry for three main tectonic
47 settings, *i.e.*, the mid-ocean ridge, arc and intraplate. Irvine (1967) and Dick and Bullen
48 (1984) referred to the Cr# and Mg# of chromian spinel, and Arai (1994a, b) discussed
49 the relationship between the Cr# of chromian spinel and Fo of coexisting olivine in
50 peridotites and related rocks. The range of Cr# of chromian spinel alone, however, has
51 considerable overlaps between the different settings (Arai 1994a). The Mg# of chromian

52 spinel is strongly dependent on the equilibrium temperature (Irvine 1965; Jackson 1969),
53 and is changeable at subsolidus stage also depending on its modal amount (Arai 1980).
54 We should notice that the Mg#-Cr# relationship commonly used for descriptions and
55 discussions of chromian spinel in igneous rocks (Irvine 1967) is strongly dependent on
56 their subsolidus cooling histories after the igneous stage.

57 In this article, we review and summarize the chromian spinel chemistry (mainly the
58 Cr# and Ti content) in deep-seated rocks (lherzolites, harzburgites, dunites, wehrlites,
59 troctolites and gabbros) to define its chemical spread for discrimination of their tectonic
60 settings and deep magmatic processes. We examined chromian spinel compositions in
61 the ultramafic and mafic plutonic rocks (mantle peridotites and ultramafic/mafic
62 cumulates) for which derived tectonic settings are well constrained. Each of the three
63 settings, i.e., the mid-ocean ridge, arc (mantle wedge) and oceanic hotspot (intraplate),
64 produces the magmas that are distinguishable from those of the other settings in
65 geochemistry (e.g. Pearce 1975). There have been plenty of articles dealing with
66 ultramafic xenoliths from continental areas captured by intra-plate basalts. The
67 deep-seated rocks from non-arc continental areas are not used in this article, because
68 they may have complicated histories, i.e., multi-setting generation/modification, and are

not appropriate to the purpose of this study. The chromian spinel chemistry has been systematically discussed both for mantle peridotites (e.g. Dick & Bullen 1984; Arai 1994a) and for volcanic rocks (e.g. Arai, 1992; Kamenetsky *et al.* 2001). No works has ever discussed chromian spinel in possibly cumulative plutonic rocks (dunites, wehrlites, troctolites and gabbros) in more systematic way than this article that deals with chromian spinels from plutonic rocks only with well-constrained derivations. The result of this work is potentially useful, because such rocks are quite commonly found from various geologic bodies. We also present two examples of application of our result to two ophiolitic dunites, which are apparently unknown or debated for the tectonic setting of formation. This article is supplementary to Arai (1992), which deals with chromian spinels in volcanic rocks from the three tectonic settings.

DATA ACQUISITION

We collected data of chromian spinel in deep-seated rocks from three main tectonic settings, i.e., mid-ocean ridge, arc and oceanic hotspot. The source of spinel chemical data treated here is unpublished theses of our laboratory in addition to the literature. Fe^{2+} and Fe^{3+} amounts were calculated assuming spinel stoichiometry. Titanium is

86 assumed to form the ulvospinel component.

87

88 MID-OCEAN RIDGES

89 We can obtain deep-seated rocks (lherzolites, harzburgites, dunites, wehrlites, troctolites

90 and gabbros) from the present-day ocean floor by dredging, drilling and submersible

91 diving (e.g. Dick 1989). The oceanic fracture zones (FZ), which are more prominently

92 developed in the slow to ultraslow spreading ridge systems than in fast spreading ones,

93 are the main loci for obtaining abyssal deep-seated rocks. Hess Deep, the East Pacific

94 Rise, is one of the non-FZ localities where deep-seated rocks are exposed on the ocean

95 floor (e.g. Arai & Matsukage 1996; Dick & Natland 1996; Allan & Dick 1996). We can

96 interpret these plutonics as deep-seated magmatic products beneath the mid-ocean ridge.

97 Harzburgites and lherzolites are predominant in the uppermost mantle of fast spreading

98 ridges and of slow spreading ridges, respectively (Niu & Hékinian 1997). Dunites and

99 related rocks (troctolites and olivine gabbros) are commonly found from Hess Deep,

100 and they may represent the Moho transition zone of the fast spreading ridge (e.g. Arai &

101 Matsukage 1996).

102

103 OPHIOLITES

104 Plutonic rocks from ophiolites may represent the deep-seated rocks of some sorts of
105 oceanic lithosphere (e.g. Coleman 1977; Nicolas 1989). Their tectonic setting for
106 genesis has been a problem in controversy since the pioneering paper of Miyashiro
107 (1973), and therefore, the data from ophiolitic plutonic rocks are not considered in this
108 section. Some ophiolites that exhibit both mid-ocean ridge and island-arc characteristics
109 are called “supra-subduction zone” (SSZ) ophiolites (Pearce *et al.* 1984). Many people
110 have favored the back-arc basin as the locus of the SSZ ophiolite formation (e.g. Pearce
111 *et al.* 1984; Moores *et al.* 1984). The polygenetic nature of some ophiolites has been
112 recently recognized as well. For example, some peridotites from the northern Oman
113 ophiolite are of arc origin (e.g. Tamura & Arai 2006; Arai *et al.* 2006), although the
114 main portion of the mantle section was of mid-ocean ridge origin (e.g. Nicolas 1989).
115 Wehrlitic rocks around the Moho transition zone were interpreted as mid-ocean ridge
116 products in the southern Oman ophiolite (Koga *et al.* 2001). For another example, the
117 mantle member of the Coast Range ophiolite, California, is composed of a mixture of
118 SSZ harzburgites and abyssal lherzolites (Choi *et al.* 2008; Jean *et al.* 2010).

119

120 ARCS

121 Deep-seated rocks from the sub-arc mantle (mantle wedge) are more difficult to obtain
122 systematically. Alkali basalts that carry deep-seated rocks as xenoliths most frequently
123 erupt on non-arc regions, i.e., on continental rift zones or oceanic hotspots, and the
124 xenoliths in kimberlites represent the upper mantle beneath cratons (e.g. Nixon 1987).
125 Genesis of calc-alkaline magmas is related with the subduction of slab (e.g. Tatsumi &
126 Eggins 1995), and their deep-seated xenoliths undoubtedly represent the sub-arc
127 deep-seated materials. Calc-alkaline andesites and basaltic andesites from Megata and
128 Oshima-Ôshima volcanoes (Northeast Japan arc), Iraya volcano (Luzon arc), and
129 Avacha and Shiveluch volcanoes (Kamchatka arc) contain peridotite xenoliths that are
130 derived from lithosphere of the mantle wedge (e.g. Takahashi 1978; Ninomiya & Arai
131 1992; Arai *et al.* 2003, 2004; Ishimaru *et al.* 2007; Bryant *et al.* 2007). It is noteworthy
132 that some of ultramafic rocks treated here form composite xenoliths with gabbros and
133 hornblendites in calc-alkaline volcanics (e.g. Ninomiya & Arai 1992; Arai *et al.* 1996,
134 2003, 2004). A variety of peridotites, i.e. lherzolites to highly refractory harzburgites
135 (Cr# of spinel < 0.8), may constitute the mantle wedge (Arai 1994a; Arai *et al.* 2003;
136 Arai & Ishimaru 2008). Dunites and related rocks form a thick cumulus mantle beneath

the Southwest Japan arc (Takahashi 1978). Dunites are relatively abundant from the Oshima-Ôshima volcano (e.g. Yamamoto 1984), and are small in amount in other localities (especially the Megata, Iraya and Avacha volcanoes).

Although not treated here, we can obtain large amounts of ultramafic xenoliths carried by non-arc type alkaline basalts on past or present-day arcs, e.g., the Japan arcs (e.g. Takahashi 1978; Aoki 1987; Abe *et al.* 1998, 1999; Arai *et al.* 1998, 2000, 2007).

These materials from the Japan arcs may represent sub-arc mantle materials because their eruption ages are mostly younger than Miocene (Uto 1990), when the present arc setting had been established for the Japan island arcs (e.g. Otofujii *et al.* 1985).

Ultramafic to mafic xenoliths from the Southwest Japan arc have been affected to various extents (Arai *et al.* 2000) by Cenozoic non-arc type alkali basalt magmas (Nakamura *et al.* 1987).

Peridotites and related rocks are exposed on the ocean floor of fore-arc regions (e.g. Fisher & Engel 1969), and the dredged and drilled peridotites distinctively represent fore-arc mantle materials (e.g. Bloomer 1983; Parkinson & Pearce 1998). Refractory harzburgites are dominant in the fore-arc mantle (Arai 1994a). In this article, we treat only these two sets of deep-seated rocks, i.e., the ultramafic xenoliths in arc-type

154 volcanics and the ultramafic rocks exposed on the present-day fore-arc ocean floor, as
155 “genuine” sub-arc materials with well-defined derivations.
156

157 OCEANIC HOTSPOTS

158 Deep-seated rocks from the oceanic hotspot areas have been almost solely obtained as
159 ultramafic and mafic xenoliths in their volcanic rocks (e.g. Nixon 1987). The xenoliths
160 in alkaline basalts especially represent the lower crust to upper mantle of that tectonic
161 setting because the xenolith-bearing magmas postdate the main stage of hotspot
162 volcanism (e.g. Jackson & Wright 1970). Lherzolites are apparently dominant in amount
163 as the upper mantle material from the oceanic hotspot (Arai 1994a). Some peridotites
164 may be related with the hotspot magmatism (cumulates or restites), and the others are
165 only representative of the sub-oceanic mantle that hosts mantle plumes relevant to the
166 hotspot activity. Arai (1994b) predicted predominance of refractory harzburgites with
167 high Cr# (around 0.7) of chromian spinel as residual peridotites after the hotspot
168 tholeiite genesis. Some basalt magmas contain a large amount of dunite xenoliths,
169 indicating a thick dunite layer at the uppermost mantle as a result of extensive tholeiitic
170 magmatism (Jackson & Wright 1970; Sen & Presnall 1986). Hawaiian and French

Polynesian hotspots are especially important for occurrences of ultramafic xenoliths (Nixon 1987), and our data accumulation owes the literature dealing with them.

CHEMICAL SPECTRA OF CHROMIAN SPINELS IN PLUTONIC ROCKS FROM THE THREE TECTONIC SETTINGS

MID-OCEAN RIDGES

As is well known, the Cr# of chromian spinel in abyssal mantle peridotites ranges from 0.1 to 0.6 (e.g. Dick & Bullen 1984; Arai 1994a; Niu & Hékinian 1997) (Fig. 1a). It changes from around 0.4 to 0.6 in harzburgites to <0.4 in lherzolites, in response to a decrease of degrees of partial melting (e.g. Dick & Bullen, 1984; Arai 1994a). The TiO₂ content is mostly lower than 0.3 wt% in their chromian spinel (Fig. 1a). The chromian spinel interestingly displays the same range of Cr# between plagioclase-bearing and -free varieties of mantle peridotites (Fig. 1a). The TiO₂ content of chromian spinel is, however, systematically higher in the plagioclase-bearing peridotites than in the plagioclase-free ones (Dick 1989). The chromian spinel exhibits relatively wide ranges of Cr#, 0.2-0.6, and TiO₂, nil to 2 wt% (mostly <1 wt%) in abyssal dunites (Fig. 1a).

Other plutonic rocks, especially troctolites and olivine gabbros from Hess Deep, show a narrow range of Cr#, mostly 0.5 to 0.6, and a wide range of TiO₂ content, <3 wt% in chromian spinel. Chromian spinel in abyssal plutonic rocks is characterized by overall low Fe³⁺ contents as that in MORB (Arai 1992). The Mg# is negatively correlated with the Cr# in chromian spinel for all peridotitic rocks including dunites from the mid-ocean ridges (Fig. 3a). Y_{Fe} (= Fe³⁺/(Cr + Al + Fe³⁺) atomic ratio) of chromian spinel is mostly lower than 0.1 in peridotitic rocks, and lower than 0.2 in troctolites and gabbros (Figs. 2a and 4a). Chromian spinel shows lower Mg#s at a Cr# around 0.5 in abyssal gabbros and troctolites. The TiO₂ content is well correlated positively with the Y_{Fe} in chromian spinel from troctolites and gabbros, being 2.5 to 3 wt% at around Y_{Fe} of 0.2 (Fig. 4a).

Rocks of dunite-troctolite-olivine gabbro suite from Hess Deep, East Pacific Rise, were interpreted as a reaction product between the primary MORB and mantle harzburgite (Arai & Matsukage 1996; Dick & Natland 1996). These rocks are expected to be in equilibrium with MORB in terms of mineral chemistry (e.g. Kelemen *et al.* 1995; Arai 2005). Plagioclase in the harzburgites/lherzolites is calcic, and is a melt-impregnation product (e.g. Dick 1989). The formation of plagioclase-bearing peridotites is the very initiation of melt/peridotite reaction. The slightly but clearly

higher TiO₂ content of chromian spinel in plagioclase-bearing peridotites (Fig. 1a) is consistent with this interpretation (e.g. Dick 1989).

ARCS

The Cr# of chromian spinel exhibits a wide range, from than <0.2 to 0.9, for mantle peridotites (lherzolite to harzburgite) and dunites (Fig. 1b). This is consistent with the wide range of spinel Cr# in sub-arc mantle restites estimated from arc magmas (Arai 1994b). The TiO₂ content of chromian spinel is, however, slightly higher in dunites than in mantle peridotites (Fig. 1b). Some of dunite, wehrlite and clinopyroxenite treated here possibly have initially formed composite xenoliths with younger gabbros or hornblendites, and have been chemically influenced by evolved magmas that formed the latter younger rocks (e.g. Ninomiya & Arai 1992). The relatively high contents of TiO₂ and Fe³⁺ of some plutonic spinels are possibly due to such a secondary effect. Almost all sub-arc spinels have low values of Y_{Fe}, < 0.3 (Fig. 2b). As is well known, the Mg# shows roughly negative correlations with the Cr# (e.g. Irvine 1967; Dick & Bullen 1984) (Fig. 3b). Two Mg#-Cr# spinel trends can be recognized in mantle peridotites, especially harzburgites, corresponding to two different derivations of the samples

treated here, namely the fore-arc rocks and xenoliths in arc magmas (Fig. 3b). This is due to the difference of equilibrium temperature between the two rock suites (e.g. Okamura *et al.* 2006), resulting from a decrease of Mg# of chromian spinel with decreasing the equilibrium temperature in peridotites (Irvine 1967; Evans & Frost 1975). Chromian spinel in some dunites, wehrlites and clinopyroxenites shows lower Mg#s at given Cr#s (Fig. 3b). The TiO₂ content is positively correlated with the Fe³⁺ ratio for the main cluster of sub-arc spinels, being 1 to 2 wt% at Y_{Fe} = 0.2 (Fig. 4b).

OCEANIC HOTSPOTS (PLUMES)

Ultramafic xenoliths have been extensively described from various oceanic hotspots on the Earth, especially from Hawaii and the French Polynesian (e.g. Nixon 1987). The Cr# of chromian spinel also changes from 0.1 to 0.8 with a lithological change from lherzolite to harzburgite (Fig. 1c). The TiO₂ content of the peridotite spinel is mostly lower than 4 wt% (Fig. 1c). The Cr# of chromian spinel shows almost the same range between the mantle peridotites and dunites. The TiO₂ content of spinel is generally higher in dunites than in mantle peridotites, and show the highest values, up to > 10 wt%, at the Cr# around 0.5 to 0.6 (Fig. 1c). Most of wehrlite spinel have relatively high

Cr#s, around 0.6, and TiO₂ content, up to 6 wt% (Fig. 1c). The Y_{Fe} of chromian spinel is highest around the intermediate Cr#, 0.5 to 0.6, and positively correlated to the TiO₂ content (Figs. 2c and 4c). The TiO₂ content of hotspot spinels varies at a given Y_{Fe}, ranging from 1 to 6 wt% at Y_{Fe} = 0.2 (Fig. 4c). Harzburgite spinels have higher Mg# at a given Cr# than dunite ones (Fig. 3c). As in the case of abyssal plutonic rocks, the Mg# is extended toward lower values at the highest Cr# of the whole range, 0.6 to 0.7, in the dunite spinels (Fig. 3c).

DISCUSSION

DISTINCTION OF THE THREE TECTONIC SETTINGS

Apart from the mantle peridotite, the plutonic rocks that bear chromian spinel are mainly dunite, troctolite and olivine gabbro from the ocean floor, but are dunite and wehrlite from the arc and the hotspot (Figs. 1 to 4). This indicates that the phase crystallizing next to olivine is mainly plagioclase in MORB but clinopyroxene in both arc and intraplate magmas. This is in turn related with the degree of partial melting in the mantle, which is lower, on average, in the mid-ocean ridge than in sub-arc and in

256 hotspot conditions (e.g. Arai 1994a,b).

257 The Cr# ranges of chromian spinel in plutonic rocks are overlapping with each
258 other around 0.1 to 0.6 for the three tectonic settings, i.e., the mid-ocean ridge, arc and
259 intraplate (Fig. 1). It is difficult, therefore, to distinguish the tectonic settings in terms of
260 Cr# of spinel alone. The Cr# of spinel is barely higher than 0.6 in plutonic rocks from
261 the mid-oceanic ridges. It is frequently over 0.6, and is up to 0.9 for the arc setting, and
262 up to over 0.7 for the intraplate (hotspot or plume) setting. The TiO₂ content in
263 chromian spinel combined with the Cr# is, however, convenient for distinction of
264 plutonic rocks between the three tectonic settings (Fig. 1). The TiO₂ content of
265 chromian spinel in plutonic rocks decreases on average from the intraplate setting to arc
266 via mid-ocean ridge setting (Fig. 1). It is concluded that deep-seated ultramafic rocks
267 can be distinguished as a group with each other in terms of spinel chemistry, especially
268 Cr# and Ti content (Fig. 1). This distinction is consistent with the diversity of chromian
269 spinel in volcanics depending on the tectonic setting (Arai 1992).

270

271 IMPLICATIONS FOR DEEP MAGMATIC PROCESSES

272 All kinds of plutonic rock have relatively low-Ti spinels from the arc setting, and

273 dunites are almost indistinguishable from harzburgites (or lherzolites) in terms of spinel
274 chemistry (see Arai 1994b). Chromian spinel in dunites, troctolites and
275 melt-impregnated harzburgites (plagioclase harzburgites) from the ocean floor is high
276 both in Cr# (around 0.6 to 0.7) and in TiO₂ (Fig. 1a). Dunites and wehrlites from the
277 oceanic hotspot also contain high-Cr# and high-Ti chromian spinels (Fig. 1c). The wide
278 range of TiO₂ content at a given Y_{Fe} for hotspot dunite spinels (Figs. 1 and 4) is possibly
279 due to a variety of dunites, from those related with older MORB genesis to those related
280 to younger hotspot magmatism. It is noteworthy that the hotspot plutonic spinels are
281 lower in Ti at a given Y_{Fe} than the mid-ocean ridge ones (Fig. 4), despite that the
282 relations are the reverse for volcanic spinels (Arai 1992). Abyssal plutonic rocks are
283 mostly troctolites (Fig. 4a), in which Ti and Fe³⁺ are partitioned to chromian spinel
284 because plagioclase is free of these components. In contrast, Ti and Fe³⁺ are partitioned
285 to both clinopyroxene and chromian spinel in wehrlites, which are common from
286 hotspot environments (Fig. 4c). This is also related with the redox condition;
287 deep-seated magmas are more oxidized for the hotspot environments than for the
288 mid-ocean ridge ones. This is concordant to the difference of oxidation states between
289 the hotspot magmas and MORB (e.g. Christie *et al.* 1986; Rhodes & Vollinger 2005).

Discrepancy in spinel chemistry between effusive rocks and related plutonic rocks is sometimes noticeable; volcanic spinels are sometimes more limited in Ti and Y_{Fe} ranges than plutonic spinels (Fig. 4). This is primarily due to effective magmatic evolution to concentrate these components within closed melt pools in deep parts (Arai *et al.* 1997). Difference in spinel chemistry is striking between MORB and abyssal plutonics (dunites, troctolites to olivine gabbros) (Fig. 4a). The TiO_2 and Y_{Fe} of chromian spinel are limited, mostly <1 wt% and <0.1, respectively in MORB (Arai 1992), as compared to the values in abyssal plutonics (Fig. 4a).

It is noteworthy that high-Ti chromian spinels are also high in Cr# (around 0.6 to 0.7) (Fig. 1). The high-Cr#, -Ti spinels are common to dunites and related rocks from the ocean floor and oceanic hotspot, part of which have been thought to be peridotite/magma reaction products (e.g. Arai & Matsukage 1996; Dick & Natland 1996). Some of dunite and wehrlite xenoliths from the island arc setting are also reaction products (e.g. Arai & Abe 1994), but the concerned arc magmas, which are initially low-Ti, have not increased the Ti contents of chromian spinel even through the peridotite/melt reaction processes.

307 EXAMPLES OF APPLICATION TO OPHIOLITES

308 The origin and nature of ophiolites have been controversial (e.g. Pearce *et al.* 1984;
 309 Nicolas 1989). Arai *et al.* (2006), for example, suggested polygenetic nature of the
 310 mantle part of the northern Oman ophiolite. We show two examples of discordant
 311 dunites from two ophiolites as below. The dunite is an important constituent of the
 312 Moho transition zone to upper mantle section of ophiolites, but the mineralogy is too
 313 simple to constrain its derivation. If we apply the systematics discussed here, the
 314 chromian spinel is indicative of the tectonic setting of the dunite formation.

315

316 DISCORDANT DUNITES FROM THE MANTLE SECTION OF THE

317 NORTHERN OMAN OPHIOLITE

318 It has been well recognized that the mantle section of the Oman ophiolite is dominated
 319 by harzburgites (e.g. Boudier & Coleman 1981; Lippard *et al.* 1986). Lherzolites are
 320 absent except at the base of the ophiolite (e.g. Lippard *et al.* 1986; Takazawa *et al.*
 321 2003). The harzburgites mainly constitute the mantle section, containing spinels with
 322 Cr#s <0.6 (Le Mée *et al.* 2004), similar to those obtained from the ocean floor of fast
 323 spreading ridge origin (Niu & Hekinian 1997). Tamura and Arai (2006) found a

324 harzburgite-orthopyroxenite-dunite suite of sub-arc chemical affinity from the northern
325 Oman ophiolite. Arai *et al.* (2006) examined chemical variations of detrital chromian
326 spinel particles derived from the mantle section from recent riverbeds in the Oman
327 ophiolite. They found more than 20 to 30 percent of the total detrital chromian spinel
328 grains examined have Cr#s higher than 0.6 (Arai *et al.* 2006).

329 Discordant dunites cutting foliated harzburgites are very common in the mantle
330 section (Fig. 5a). They form dikes or networks, and some of them contain chromian
331 spinel concentrations (podiform chromitites) (e.g. Augé 1987; Ahmed & Arai 2002).
332 They are massive in appearance and solely comprise olivine and euhedral to subhedral
333 chromian spinel (Fig. 5b). We examined chromian spinels in the discordant dunites from
334 Wadi Rajmi and Wadi Fizh areas of the northern Oman ophiolite (Fig. 5a). The Cr# of
335 chromian spinel ranges from 0.4 to 0.8 with very low amounts of TiO₂, mostly < 0.3
336 wt% (Fig. 6). Olivine associated with the chromian spinel is around Fo₉₀₋₉₂ in
337 composition. The Oman discordant dunites are most likely to have been related with arc
338 magmas (see Figs. 1, 2 and 4). The mantle section of the Oman ophiolite, therefore,
339 comprises the ocean-floor peridotites (mainly harzburgites) modified by addition of
340 dunites of sub-arc affinity. This suggests a switch of tectonic setting from mid-ocean

ridge to arc (= supra-subduction zone) for genesis of the Oman ophiolite (Arai *et al.* 2006). Alternatively, this characteristic can be obtained at a back-arc tectonic setting, where various magmas, from MORB-like to arc-type, are available (e.g. Pearce *et al.* 1984).

DISCORDANT DUNITES FROM THE LIZARD OPHIOLITE, CORNWALL

The mantle section of the Lizard ophiolite, Cornwall (Kirby 1979), is mainly composed of lherzolites and concordant dunites (Green 1964; Kadoshima & Arai 2001). This is very similar to a peridotite suite from the ocean floor of slow spreading ridge origin (Roberts *et al.* 1993) if we consider the abundance of lherzolite over harzburgite (e.g. Niu & Hekinian 1997). The Cr# ranges from 0.1 to 0.5 for the Lizard detrital spinels (Kadoshima & Arai 2001), exactly being the same as those of abyssal peridotites (lherzolites to harzburgites) (Fig. 6). This is consistent with the idea that the Lizard peridotite was representative of the uppermost sub-oceanic mantle of a slow spreading ridge, which may be composed of predominant lherzolites and subordinate harzburgites (Arai 2005).

Networks of younger discordant dunites are prominently cutting the concordant

358 lherzolites and dunites, especially around the central part of the ophiolite (Kadoshima &
359 Arai 2001). The younger discordant dunites are black in hand specimen, and seem
360 compact and hard on outcrop (Fig. 5c). This is in contrast to the concordant dunites that
361 are severely altered/weathered to be pale green in color and have been more strongly
362 eroded than the discordant ones due to mechanical weakness. This observation clearly
363 indicates different chemical and/or textural characteristics between the two types of
364 dunite. The discordant dunites are exclusively composed of olivine, highly serpentinized,
365 and euhedral to subhedral chromian spinel. The chromian spinel is opaque in thin
366 section and contains minute exsolution blebs of a Ti-rich phase (possibly Ti-rich
367 magnetite) (Fig. 5d). The olivine shows slightly lower Fo contents, 83 to 88, than in the
368 wall peridotite (89-90). The chromian spinel of this younger dunite is high in Cr# and
369 TiO₂, being around 0.6 and up to > 4 wt%, respectively (Fig. 6). It is low in both Cr#
370 and TiO₂ near the boundary with lherzolite (Fig. 6), suggesting fractional crystallization
371 (precipitation of minerals from the wall inward) or a reaction between the involved melt
372 and the lherzolite. The primary chromian spinel in the discordant dunite should have
373 contained higher TiO₂ contents before unmixing of the Ti-rich phase, being within the
374 chemical range of dunite spinels from oceanic hotspots (see Figs. 1 and 2). The magma

that produced the discordant dunite within the Lizard peridotite was of hotspot (intra-plate) origin (Figs. 1, 2, 4 and 6). It was most probably tholeiitic (cf. Arai 1992). The Lizard peridotite was, therefore, derived from the uppermost mantle that was initially generated at a slow spreading ridge and was later impacted by an intra-plate tholeiite magma.

CONCLUSIONS

The chromian spinel chemistry is highly useful to petrologically characterize ultramafic plutonic rocks, especially dunitic rocks and chromitites, where chromian spinel is often the only discriminating mineral. The trivalent cation ratio and TiO_2 content in chromian spinel are important parameters in discrimination of the plutonic rocks in terms of tectonic setting of formation. Discrimination diagrams based on spinel chemistry made from plutonic rocks derived from well-constrained settings should be applied to characterization of rocks from unknown origins. The spinel-based diagrams made for volcanic rocks or magmas should not been used for discrimination of plutonic rocks in tectonic setting of derivation, because chromian spinel shows different chemical ranges between effusive and plutonic rocks even of the same magmatic affinity as discussed

above. Discrimination in Mg# of chromian spinel is sometimes unreliable, because the Mg# in chromian spinel is strongly changeable depending on the thermal history in olivine-rich rocks. For example, the chromian spinel in possible abyssal peridotites suffered from low-temperature metamorphism at a subduction zone have lower Mg#s at a given Cr# than abyssal peridotites from the present-day ocean floor, which are mostly derived from near the spreading center and have not been cooled down sufficiently (e.g. Hirauchi *et al.* 2008).

ACKNOWLEDGEMENTS

We are grateful to T. Morishita and A. Ishiwatari for their discussions. S.A. thanks H. Hirai, N. Abe, M. Kida, Y. Kobayashi, M. Fujiwara, Y. Saeki, A. Ninomiya, S. Takada, N. Takahashi and K. Goto for their collaboration on mantle xenoliths derived from the mantle wedge. We appreciate suggestions made by S.H. Choi (associate editor), Y.J. Yu and J.W. Shervais, which were helpful in revision.

REFERENCES

ABE N., ARAI S. & SAEKI Y. 1992. Hydration processes in the arc mantle; petrology

- 409 of the Megata peridotite xenolith the Northeast Japan arc. *Journal of Mineralogy,*
 410 *Petrology and Economic Geology* **87**, 305-17. (in Japanese with English abstract)
- 411 ABE N., ARAI S. & NINOMIYA A. 1995. Peridotite xenoliths and essential ejecta from
 412 the Ninomegata crater, the Northeastern Japan arc. *Journal of Mineralogy, Petrology*
 413 *and Economic Geology* **90**, 41-9. (in Japanese with English abstract)
- 414 ABE N., ARAI S. & YURIMOTO H. 1998. Geochemical characteristics of the
 415 uppermost mantle beneath the Japan island arcs: implications for upper mantle
 416 evolution. *Physics of Earth and Planetary Interior* **107**, 233-47.
- 417 ABE N., ARAI S. & YURIMOTO H. 1999. Texture-dependent geochemical variations
 418 of sub-arc mantle peridotite from Japan island arcs. *Proceedings of VIIth*
 419 *International Kimberlite Conference J.B. Dawson Vol.*, 13-22.
- 420 AHMED A.H. & ARAI S. 2002. Unexpectedly high-PGE chromitite from the deeper
 421 mantle section of the northern Oman ophiolite and its tectonic implications.
 422 *Contributions to Mineralogy and Petrology* **143**, 263-78.
- 423 ALLAN J.F. & DICK H.J.B. 1996. Cr-rich spinel as a tracer for melt migration and
 424 melt-wall rock interaction in the mantle: Hess Deep, Leg 147. *Proceedings of ODP,*
 425 *Scientific Results* **147**, 157-72.

- 426 AOKI K. 1987. Japanese Island arc: xenoliths in alkali basalts, high-alumina basalts,
427 and calc-alkaline andesites and dacites. *In* Nixon P.H. (ed.) *Mantle Xenoliths*, pp
428 314-331, John Wiley & Sons, New York.
- 429 ARAI S. 1980. Dunite-harzburgite-chromitite complexes as refractory residue in the
430 Sangun-Yamaguchi zone, western Japan. *Journal of Petrology* **21**, 141-65.
- 431 ARAI S. 1992. Chemistry of chromian spinel in volcanic rocks as a potential guide to
432 magma chemistry. *Mineralogical Magazine* **56**, 173-84.
- 433 ARAI S. 1994a. Characterization of spinel peridotites by olivine-spinel compositional
434 relationships: Review and interpretation. *Chemical Geology* **113**, 191-204.
- 435 ARAI S. 1994b. Compositional variation of olivine-chromian spinel in Mg-rich
436 magmas as a guide to their residual spinel peridotites. *Journal of Volcanology and*
437 *Geothermal Research* **59**, 279-94.
- 438 ARAI S. 2005. Role of dunite in genesis of primitive MORB. *Proceedings of Japan*
439 *Academy. Series B* **81**, 14-9.
- 440 ARAI S. & ABE N. 1994. Podiform chromitite in the arc mantle: chromitite xenoliths
441 from the Takashima alkali basalt, southwest Japan arc. *Mineralium Deposita* **29**,
442 434-8.

- 443 ARAI S., ABE N. & HIRAI H. 1998. Petrological characteristics of the sub-arc mantle:
444 an overview on petrology of peridotite xenoliths from the Japan arcs. *Trends in*
445 *Mineralogy (India)* **2**, 39-55.
- 446 ARAI S., ABE N. & ISHIMARU S. 2007. Mantle peridotites from the Western Pacific.
447 *Gondwana Research*, **11**, 180-99.
- 448 ARAI S., HIRA H. & UTO K. 2000. Mantle peridotite xenoliths from the Southwest
449 Japan arc and a model for the sub-arc upper mantle structure and composition of the
450 Western Pacific rim. *Journal of Mineralogical and Petrological Sciences* **95**, 9-23.
- 451 ARAI S. & ISHIMARU S. 2008. Insights into petrological characteristics of the
452 lithosphere of mantle wedge beneath arcs through peridotite xenoliths: A review.
453 *Journal of Petrology* **49**, 665-95.
- 454 ARAI S., ISHIMARU S. & OKRUGIN V.M. 2003. Metasomatized harzburgite
455 xenoliths from Avacha volcano as fragments of mantle wedge of the Kamchatka arc:
456 an implication for the metasomatic agent. *Island Arc* **12**, 233-46.
- 457 ARAI S., KADOSHIMA K. & MORISHITA T. 2006. Widespread arc-related melting in
458 the mantle section of the northern Oman ophiolite as inferred from detrital chromian
459 spinels. *Journal of Geological Society, London* **163**, 869-79 .

- 460 ARAI S., KIDA M., ABE N., NINOMIYA A. & YUMUL G.P. JR. 1996. Classification
 461 of peridotite xenoliths in calc-alkaline andesite from Iraya volcano, Batan Island, the
 462 Philippines, and its genetical implications. *Science Reports of Kanazawa University*
 463 **41**, 25-45.
- 464 ARAI S. & MATSUKAGE K. 1996. Petrology of the gabbro-troctolite-peridotite
 465 complex from Hess Deep, equatorial Pacific: implications for mantle-melt
 466 interaction within the oceanic lithosphere. *Proceedings of ODP, Scientific Results*
 467 **147**, 135-55.
- 468 ARAI S., MATSUKAGE K., ISOBE E. & VYSOTSKIY S. 1997. Concentration of
 469 incompatible elements in oceanic mantle: Effect of melt/wall interaction in stagnant
 470 or failed conduits within peridotite. *Geochimica et Cosmochimica Acta* **61**, 671-175.
- 471 ARAI S., TAKADA S., MICHIBAYASHI K. & KIDA M. 2004. Petrology of peridotite
 472 xenoliths from Iraya Volcano, Philippines, and its implication for dynamic
 473 mantle-wedge processes. *Journal of Petrology* **45**, 369-89.
- 474 AUGÉ T. 1987. Chromite deposits in the northern Oman ophiolite: mineralogical
 475 constraints. *Mineralium Deposita*, **109** 301-4.
- 476 BARSDELL M & SMITH I.E.M. 1989. Petrology of recrystallized ultramafic xenoliths

- 477 from Merelava volcano, Vanuatu. *Contributions to Mineralogy and Petrology* **102**,
478 230-41.
- 479 BLOOMER S.H. 1983. Distribution and origin of igneous rocks from the landward
480 slopes of the Mariana Trench; Implications for its structure and evolution. *Journal of*
481 *Geophysical Research* **88**, 7411-28.
- 482 BLOOMER S.H. & FISHER R.L. 1987. Petrology and geochemistry of igneous rocks
483 from the Tonga trench — A non-accreting plate boundary. *Journal of Geology* **95**,
484 469-95.
- 485 BLOOMER S.H. & HAWKINS J.W. 1983. Gabbroic and ultramafic rocks from the
486 Mariana Trench: An island arc ophiolite. In Hayes D.E. (ed.) *The tectonic and*
487 *geologic evolution of southeast Asian seas and Islands (Part 2)*, pp 294-317,
488 Geophysical Monograph No. 27, American Geophysical Union, Washington, D.C..
- 489 BOUDIER F. & COLEMAN R.G. 1981. Cross section through the peridotite in the
490 Samail ophiolite, southeastern Oman mountains. *Journal Geophysical Research* **86**,
491 2573-92.
- 492 BRYANT J.A., YOGODZINSKI G.M. & CHURIKOVA T.G. 2007. Melt-mantle
493 interaction beneath the Kamchatka arc: Evidence from ultramafic xenoliths from

- 494 Shiveluch volcano. *Geochemistry, Geophysics, Geosystems* **8**, Q04007. doi:
495 10.1029200GC001443.
- 496 CANNAT M., CHATIN F., WHITECHURCH H. & CEULENEER G. 1997. Gabbroic
497 rocks trapped in the upper mantle at the mid-Atlantic ridge. *Proceedings of ODP,*
498 *Scientific Results* **153**, 243-64.
- 499 CHOI S.H., SHERVAIS J.W. & MUKASA S.B. 2008. Supra-subduction and abyssal
500 mantle peridotites of the Coast Range ophiolite, California. *Contributions to*
501 *Mineralogy and Petrology* **156**, 551-576.
- 502 CHRISTIE D.M., CARMICHAEL I.S.E. & LANGMUIR C.H. 1986. Oxidation states
503 of mid-ocean ridge basalt glasses. *Earth Planetary Science Letters* **79**, 397-411.
- 504 CLAGUE D.A. 1988. Petrology of ultramafic xenolith from Loihi seamount, Hawaii.
505 *Journal of Petrology* **29**, 1161-86.
- 506 COLEMAN R.G. 1977. *Ophiolites*. Springer, Berlin.
- 507 CONRAD W.K. & KAY R.W. 1984. Ultramafic and mafic inclusions from Adak Island:
508 crystallization history, and implications for the nature of primary magmas and
509 crustal evolution in the Aleutian arc. *Journal of Petrology* **25**, 88-125.
- 510 DEBARI S., KAY S.M. & KAY R.W. 1987. Ultramafic xenoliths from Adagdak

- 511 volcano, Adak, Aleutian Islands, Alaska: deformed igneous cumulates from the
512 Moho of an island arc. *Journal of Geology* **95**, 329-41.
- 513 DELONG S.E., HODGES F.N. & ARCULUS R.J. 1975. Ultramafic and mafic
514 inclusions, Kanaga Island, Alaska, and the occurrence of alkaline rocks in island
515 arcs. *Journal of Geology* **83**, 721-36.
- 516 DICK H.J.B. 1989. Abyssal peridotites, very slow spreading ridges and ocean ridge
517 magmatism. In Saunders A.D. and Norry M.J. (eds.) *Magmatism in the Ocean*
518 *Basins*, Geological Society Special Publications No. 42, pp 71-106, The Geological
519 Society, London.
- 520 DICK H.J.B. & BULLEN T. 1984. Chromian spinel as a petrogenetic indicator in
521 abyssal and alpine type peridotites and spatially associated lavas. *Contributions to*
522 *Mineralogy and Petrology* **86**, 54-76.
- 523 DICK H.J.B. & NATLAND J. 1996. Last-stage melt evolution and transport in the
524 shallow mantle beneath the East Pacific Ridge. *Proceedings of ODP, Scientific*
525 *Results* **147**, 103-34.
- 526 EVANS B.W. & FROST B.R. 1975. Chrome-spinel in progressive metamorphism: A
527 preliminary analysis. *Geochimica et Cosmochimica Acta* **39**, 959-72.

- 528 FISHER R.L. & ENGEL C.G. 1969. Ultramafic and basaltic rocks dredged from the
 529 nearshore flank of the Tonga trench. *Geological Society of America Bulletin* **80**,
 530 1373-78.
- 531 FUJII T. 1990. Petrology of peridotites from Hole 670A, Leg109. *Proceedings of ODP*,
 532 *Scientific Results* **106/109**, 19-25.
- 533 GREEN D.H. 1964. The petrogenesis of the high-temperature peridotite intrusion in the
 534 Lizard area, Cornwall. *Journal of Petrology* **5**, 134-88.
- 535 HIRAUCHI K., TAMURA A., ARAI S., YAMAGUCHI H. & HISADA K. 2008. Fertile
 536 abyssal peridotites within the Franciscan subduction complex, central California:
 537 possible origin as detached remnants of oceanic fracture zones located close to a
 538 slow-spreading ridge. *Lithos* **105**, 319-28.
- 539 IRVINE T.N. 1965. Chromian spinel as a petrogenetic indicator; part I, Theory.
 540 *Canadian Journal of Earth Sciences* **2**, 648-71.
- 541 IRVINE T.N. 1967. Chromian spinel as a petrogenetic indicator; part II, Petrologic
 542 applications. *Canadian Journal of Earth Sciences* **4**, 71-103.
- 543 ISHII T. 1985. Dredged samples from the Ogasawara fore-arc seamount or “Ogasawara
 544 Paleoland” – “fore-arc ophiolite”. In Nasu N., Kobayashi K., Uyeda S., Kushiro I.

- 545 and Kagami H. (eds.), *Formation of Active Ocean Margin*, pp 307-42, TERRAPUB,
546 Tokyo.
- 547 ISHII T., ROBINSON P.T., MAEKAWA H. & Fiske R. 1992. Petrological studies of
548 peridotites from diapiric serpentinite seamounts in the Izu-Ogasawara-Mariana
549 forearc, Leg 125. *Proceedings of ODP, Scientific Results* **125**, 445-85.
- 550 ISHII T., SATO H., HARAGUCHI S., FRAYER P., FUJIOKA K., BLOOMER S. &
551 YOKOSE H. 2000. Petrological characteristics of peridotites from serpentinite
552 seamounts in the Izu-Ogasawara-Mariana forearc. *Journal of Geography (Tokyo)*
553 **109**, 517-30. (in Japanese with English abstract)
- 554 ISHIMARU S. 2004. *The petrological characteristics of the mantle wedge beneath the*
555 *Kamchatka arc*. Unpublished MSc thesis, Kanazawa University, Kanazawa.
- 556 ISHIMARU S., ARAI S., ISHIDA Y., SHIRASAKA M. & OKRUGIN V. M. 2007.
557 Melting and multi-stage metasomatism in the mantle wedge beneath a frontal arc
558 inferred from highly depleted peridotite xenoliths from the Avacha volcano,
559 southern Kamchatka. *Journal of Petrology* **48**, 395-433.
- 560 JACKSON E.D. 1969. Chemical variation in co-existing chromite and olivine in
561 chromite zones of the Stillwater complex. In H.D.B. Wilson H.D.B. (ed.) *Magmatic*

- 562 *Ore Deposits*, Economic Geology Monograph No. 6, pp 41-71, The Economic
563 Geology Publishing Company, New Haven.
- 564 JACKSON E.D. & WRIGHT T.L. 1970. Xenoliths in the Honolulu Volcanic Series,
565 Hawaii. *Journal of Petrology* **11**, 405-30.
- 566 JEAN M.M., SHERVAIS J.W., CHOI S.H. & MUKASA S.B. 2010. Melt extraction and
567 melt refertilization in mantle peridotite of the Coast Range ophiolite: an La-ICP-MS
568 study. *Contributions to Mineralogy and Petrology* **159**, 113-136.
- 569 KADOSHIMA K. & ARAI S. 2001. Chemical analysis of detrital chromian spinels
570 from the Lizard area, Cornwall, England: an attempt for lithological and petrological
571 survey of the Lizard peridotite. *Neues Jahrbuch für Mineralogie Monatshefte* **2001**,
572 193-209.
- 573 KAMENETSKY V.S., CRAWFORD A.J. & MEFFRE S. 2001. Factors controlling
574 chemistry of magmatic spinel: an empirical study of associated olivine, Cr-spinel
575 and melt inclusions from primitive rocks. *Journal of Petrology* **42**, 655-71.
- 576 KELEMEN P.B., SHIMIZU N. & SALTERS V.J.M. 1995. Extraction of
577 mid-ocean-ridge basalt from the upwelling mantle by focused flow of melt in dunite
578 channels. *Nature* **375**, 747-753.

- 579 KIRBY G.A. 1979. The Lizard Complex as an ophiolite. *Nature* **282**, 58-61.
- 580 KOGA K.T., KELEMEN P.B. & SHIMIZU N. 2001. Petrogenesis of the crust-mantle
581 transition zone and the origin of lower crustal wehrlite in the Oman ophiolite.
582 *Geochemistry, Geophysics, Geosystems* **2**, 2000GC000132.
- 583 KOMOR S.C., GROVE T.L. & HÉBERT R. 1990. Abyssal peridotites from ODP Hole
584 670A (21°10'N, 45°02'W): Residues of mantle melting exposed by non-constructive
585 axial divergence. *Proceedings of ODP, Scientific Results* **106/109**, 85-101.
- 586 LE MÉE L., GIRARDEAU J. & MONNIER C. 2004. Mantle segmentation along the
587 Oman ophiolite fossil mid-ridge. *Nature* **432**, 167-72.
- 588 LIPPARD S.J., SHELTON A.W. & GASS I.G. 1986. *The Ophiolite of Northern Oman*.
589 Geological Society Memoir No. 11, Geological Society, London.
- 590 MIYASHIRO A. 1973. The Troodos ophiolitic complex was probably formed in an
591 island arc. *Earth and Planetary Science Letters* **19**, 218-224.
- 592 MOORES E.M., ROBINSON P.T., MALPAS J. & XENOPHONTOS C. 1984. Model
593 for origin of the Troodos massif, Cyprus, and other mideast ophiolites. *Geology* **12**,
594 500-503.

- 595 NAKAMURA E., CAMPBELL I.H., McCULLOCH M.T. 1989. Chemical
596 geodynamics in a back arc region around the Sea of Japan: implications for the
597 genesis of alkaline basalts in Japan, Korea, and China. *Journal of Geophysical*
598 *Research* **94**, 4634-4654.
- 599 NICOLAS A. 1989. *Structure of Ophiolites and Dynamics of Oceanic Lithosphere*.
600 Kluwer, Dordrecht.
- 601 NIIDA K. 1997. Mineralogy of Mark peridotites: Replacement through magma
602 channeling examined from Hole920D, MARK area. *Proceedings of ODP, Scientific*
603 *Results* **153**, 265-75.
- 604 NINOMIYA A. & ARAI S. 1992. Harzburgite fragment in a composite xenolith from an
605 Oshima-Oshima andesite, the Northeast Japan arc. *Bulletin of Volcanological*
606 *Society of Japan* **37**, 269-73 (in Japanese).
- 607 NIU Y. & HÉKINIAN R. 1997. Spreading-rate dependence of the extent of mantle
608 melting beneath ocean ridges. *Nature* **385**, 326-29.
- 609 NIXON P.H. 1987. *Mantle Xenoliths*. John Wiley & Sons, New York.
- 610 OHARA Y. & ISHII T. 1998. Peridotites from the southern Mariana forearc:
611 Heterogeneous fluid supply in the mantle wedge. *Island Arc* **7**, 541-58.

- 612 OKAMURA H., ARAI S. & KIM Y.U. 2006. Petrology of forearc peridotite from the
613 Hahajima Seamount, the Izu-Bonin arc, with special reference to chemical
614 characteristics of chromian spinel. *Mineralogical Magazine* **70**, 15-26.
- 615 OTOFUJI Y., MATSUDA T. & NOHDA S. 1985. Paleomagnetic evidence from the
616 Miocene counter-clockwise rotation of Northeast Japan – rifting process of the
617 Japan Sea. *Earth and Planetary Science Letters* **75**, 265-77.
- 618 PARKINSON I.J. & PEARCE J.A. 1998. Peridotites from the Izu-Bonin-Mariana
619 forearc (ODP Leg 125): Evidence for mantle melting and melt-mantle interaction in
620 a supra-subduction zone setting. *Journal of Petrology* **39**, 1577-618.
- 621 PEARCE J.A. 1975. Basalt geochemistry used to investigate past tectonic environments
622 on Cyprus. *Tectonophysics* **25**, 41-67.
- 623 PEARCE J.A., LIPPARD S.J. & ROBERTS S. 1984. Characteristics and tectonic
624 significance of supra subduction zone ophiolites. In Kokelaar B.P. and Howells M.R.
625 (eds.) *Marginal Basin Geology*, Geological Society Special Publications No. 16, pp
626 77-94, The Geological Society, London.
- 627 PRINZ M., KEIL K., GREEN J.A., REID A.M., BONATTI E. & HONNOREZ J. 1976.
628 Ultramafic and mafic dredge samples from the equatorial mid-Atlantic and fracture

- 629 zones. *Journal of Geophysical Research* **81**, 4087-103.
- 630 RHODES J.M. & VOLLINGER M.J. 2005. Ferric/ferrous ratios in 1984 Mauna Loa
- 631 lavas: a contribution to understanding the oxidation state of Hawaiian magmas.
- 632 *Contributions to Mineralogy and Petrology* **149**, 666-74.
- 633 ROBERTS S., ANDERS J.R., BULL J.M. & SANDERSON D.J. 1993. Slow-spreading
- 634 ridge-axis tectonics: evidence from Lizard Complex, U.K. *Earth and Planetary*
- 635 *Science Letters* **116**, 101-12.
- 636 ROEDER P.L. 1994. Chromite: From the fiery rain of chondrules to the Kilauea Iki lava
- 637 lake. *Canadian Mineralogist* **32**, 729-46.
- 638 SEN G. & PRESNALL D.C. 1986. Petrogenesis of dunite xenoliths from Koolau
- 639 volcano, Oahu, Hawaii: Implications for Hawaiian volcanism. *Journal of Petrology*
- 640 **27**, 197-217.
- 641 TAKAHASHI E. 1978. Petrological model of the crust and upper mantle of the
- 642 Japanese Island Arcs. *Bulletin Volcanologique* **41**, 529-47.
- 643 TAKAZAWA E., OKAYASU T. & SATOH K. 2003. Geochemistry and origin of the
- 644 basal lherzolites from the northern Oman ophiolite (northern Fizh block).
- 645 *Geochemistry, Geophysics, Geosystems* **4**, 1021, doi:10.1029/2001GC000232.

- 646 TAMURA A. & ARAI S. 2006. Harzburgite-dunite-orthopyroxenite suite as a record of
647 supra-subduction zone setting for the Oman ophiolite mantle. *Lithos* **90**, 43-56.
- 648 TANAKA C. 1999. *Upper mantle beneath hotspots inferred from peridotite xenoliths*.
649 Unpublished MSc thesis, Kanazawa University, Kanazawa.
- 650 TARTAROTTI P., SUSINI S., NIMIS P. & OTTOLINI L. 2002. Melt migration in the
651 upper mantle along the Romanche Fracture Zone (Equatorial Atlantic). *Lithos* **63**,
652 125-49.
- 653 TATSUMI Y. & EGGINS S. 1995. *Subduction Zone Magmatism*. Blackwell,
654 Cambridge.
- 655 TRACY R.J. 1980. Petrology and genetic significance of an ultramafic xenolith suite
656 from Tahiti. *Earth and Planetary Science Letters* **48**, 80-96.
- 657 UTO K. 1990. Neogene volcanism of Southwest Japan: Its time and space on K-Ar
658 dating. Ph.D. thesis, University of Tokyo, Tokyo.
- 659 YAMAMOTO M. 1984. Origin of calc-alkaline andesite from Oshima-Ōshima volcano,
660 north Japan. *Journal of Faculty of Science, Hokkaido University, Series 4*, **21**, 1,
661 77-131.
- 662
- 663

Figure Captions

Fig. 1. Relationships between TiO_2 contents and $\text{Cr}/(\text{Cr} + \text{Al})$ atomic ratios of chromian spinels in plutonic rocks. Upper and lower panels are for dunites and other possible cumulates, and for peridotites, respectively. (a) Mid-ocean ridges. Data source: Allan and Dick (1996), Arai and Matsukage (1996), Cannat *et al.* (1997), Dick (1989), Dick and Natland (1996), Fujii (1990), Niida (1997), Komor *et al.* (1990), Prinz *et al.* (1976), and Tartarotti *et al.* (2002). (b) Arcs. Data source: Abe *et al.* (1992, 1995), Arai *et al.* (2004), Barsdell and Smith (1989), Bloomer and Hawkins (1983), Bloomer and Fisher (1987), Conrad and Kay (1984), Debari *et al.* (1987), Delong *et al.* (1975), Ishii (1985), Ishii *et al.* (1992, 2000), Ishimaru (2004), Ninomiya and Arai (1992), Ohara and Ishii (1998), and Yamamoto (1984). (c) Oceanic hotspots. Data source: Clague (1988), Sen and Presnall (1986), Tracy (1980), and Tanaka (1999). Peridotites, lherzolites and harzburgites. Pl, plagioclase. Ol, olivine. Note the different compositional ranges between the three tectonic settings. Scales of vertical axis are different between (a) and (b), and (c).

Fig. 2. Cr-Al-Fe³⁺ atomic relationships of chromian spinels in plutonic rocks. Right and left panels are for dunites and other possible cumulates, and for peridotites, respectively. (a) Mid-ocean ridges. (b) Arcs. (c) Oceanic hotspots. Peridotites, lherzolites and harzburgites. Pl, plagioclase. Ol, olivine. Data source as in Fig. 1.

Fig. 3. Relationships between Mg/(Mg + Fe²⁺) and Cr/(Cr + Al) atomic ratios of chromian spinels in plutonic rocks. Upper and lower panels are for dunites and other possible cumulates, and for peridotites, respectively. (a) Mid-ocean ridges. (b) Arcs. (c) Oceanic hotspots. Peridotites, lherzolites and harzburgites. Pl, plagioclase. Ol, olivine. Data source as in Fig. 1.

Fig. 4. Relationships between TiO₂ contents and Fe³⁺/(Cr + Al + Fe³⁺) atomic ratios of chromian spinels in plutonic rocks. Upper and lower panels are for dunites and other possible cumulates, and for peridotites, respectively. (a) Mid-ocean ridges. (b) Arcs. (c) Oceanic hotspots. Peridotites, lherzolites and harzburgites. Pl, plagioclase. Ol, olivine. The fields for MOR (a) and hotspot (c) plutonics are shown in the panel (b). The field for MORB spinels (Arai, 1992) is shown in the panel (a) for

comparison. Data source as in Fig. 1. Fields for the main clusters of mid-ocean ridge (MOR) and hotspot spinels are shown in the panel (b).

Fig. 5. Photographs of discordant dunites. (a) Outcrop of discordant dunites (D) within foliated mantle harzburgite from Wadi Rajmi, the northern Oman ophiolite. (b) Photomicrograph of a partially serpentinized discordant dunite from Wadi Rajmi. Plane-polarized light. (c) Outcrop of a discordant dunite (D; selectively eroded) from the Lizard ophiolite. (d) Photomicrograph of a chromian spinel grain with high-Ti exsolution blebs (brighter) in a partially serpentinized discordant dunite from the Lizard ophiolite. Reflected plane-polarized light. Note the bright band fringing the right-side margin is remnants of carbon coating.

Fig. 6. Chromian spinel compositions in discordant dunites from the Oman and Lizard ophiolites. Note the different compositional characteristics between the two dunite spinels. (a) TiO_2 vs. $\text{Cr}/(\text{Cr} + \text{Al})$ atomic ratio. Compare with Figure 1. (b) TiO_2 vs. $\text{Fe}^{3+}/(\text{Cr} + \text{Al} + \text{Fe}^{3+})$ atomic ratios. Compare with Figure 4. (c) Cr-Al- Fe^{3+} atomic ratios. The fields for MOR and hotspot dunites (Figure 2) are shown in the

715 panel (b).

716

Fig.1 Arai et al.

Fig. 2 Arai et al.

Fig. 3 Arai et al.

Fig. 4 Arai et al.

Fig. 5 Arai et al.

Fig. 6 Arai et al.