Sphingosine-1-phosphate signaling in physiology and diseases

著者	Takuwaa Yoh, Okamoto Yasuo, Yoshioka Kazuaki, Takuwa Noriko
journal or publication title	BioFactors
volume	38
number	5
page range	329-337
year	2012-09-01
URL	http://hdl.handle.net/2297/31980

doi: 10.1002/biof.1030

Sphingosine-1-phosphate signaling in physiology and diseases

Yoh Takuwa^a, Yasuo Okamoto^a, Kazuaki Yoshioka^a, Noriko Takuwa^{a,b}

^aDepartment of Physiology, Kanazawa University School of Medicine, 13-1 Takara-machi, Kanazawa, Ishikawa 920-8640, Japan, and ^bDepartment of Health and Medical Sciences, Ishikawa Prefectural Nursing University, 1-1 Gakuendai, Kahoku, Ishikawa 929-1210, Japan

Short Title: S1P signaling

Correspondence to: Yoh Takuwa, M.D., Ph.D., Department of Physiology, Kanazawa University School of Medicine, 13-1 Takara-machi, Kanazawa, Ishikawa 920-8640, Japan ytakuwa@med.kanazawa-u.ac.jp

TEL: +81-76-265-2165

FAX: +81-76-234-4223

Abstract

Sphingosine-1-phosphate (S1P), which acts as both the extracellular and intracellular messenger, exerts pleiotropic biological activities including regulation of embryonic development, formation of the vasculature, vascular barrier integrity, vascular tonus and lymphocyte trafficking. Many of these S1P actions are mediated by five members of the G protein-coupled S1P receptors (S1P₁~S1P₅) with overlapping but distinct coupling to heterotrimeric G proteins. S1P₁ couples exclusively to G_i whereas S1P₂ and S1P₃ couple to multiple G proteins. S1P₂ and S1P₃ prefer G_{12/13} and G_q, respectively, among others. The biological activities of S1P are based largely on the cellular actions of S1P on migration, adhesion and proliferation. Notably, S1P often exhibits bimodal effects in these cellular actions in a receptor subtype-specific manner. For example, S1P₁ mediates cell migration toward S1P, i.e. chemotaxis, via G_i/Rac pathway whereas S1P₂ mediates inhibition of migration toward a chemoattractant, i.e. chemorepulsion, via G_{12/13}/Rho pathway which induces Rac inhibition. In addition, S1P₁ mediates stimulation of cell proliferation through the G_i-mediated signaling pathways including phosphatidylinositol 3-kinase (PI3K)/Akt and ERK whereas S1P₂ mediates inhibition of cell proliferation through mechanisms involving G_{12/13}/Rho/Rho kinase/PTEN-dependent Akt inhibition. These differential effects of S1P receptor subtypes on migration and proliferation lead to

bimodal regulation of various biological responses. An observed biological response is likely determined by an integrated outcome of the counteracting signals input by S1P receptor subtypes expressed in the cells. More recent studies identified the new intracellular targets of S1P; S1P acts as the intracellular messenger to bind to the inflammatory signaling molecule TRAF2 downstream of TNF receptor and to histone deacetylases HDAC1 and HDAC2, resulting in activation of NF- κ B and inhibition of histone deacetylation, respectively. Development of S1P receptor agonists and antagonists with improved receptor subtype-selectivity and their optimal drug delivery system augments useful actions and attenuates deleterious effects of S1P, thus providing novel therapeutic tactics. Inhibitors or modulators of S1P-synthesizing and -metabolizing enzymes also could be potential therapeutic tools.

Key words: sphingosine-1-phosphate, lysophospholipid, G protein-coupled receptor

Introduction

Sphingosine-1-phosphate (S1P), lysophosphatidic acid (LPA), lysophosphatidylserine and lysophosphatidylinositol constitute lysophospholipid mediators and are attracting increasing interest in cell signaling for the past decade. Among these lysophospholipid mediators, S1P and LPA are much better characterized than the others. It is now recognized that the S1P signaling system comprises S1P synthesizing/degrading enzymes [1,2], membrane S1P transporters [3,4], S1P carrier proteins in the plasma [5], and five members of the G protein-coupled S1P-specific receptor subtypes, S1P₁~S1P₅ [6-8]. S1P plays crucial roles in embryonic development and post-natal homeostasis in the cardiovascular, immune and nervous systems [9-13]. The S1P signaling system is also implicated as the target of therapeutic intervention in a variety of human diseases; multiple sclerosis, a debilitating autoimmune disease, is now treated with the S1P receptor agonist prodrug FTY720 [14], whose phosphorylation product downregulates S1P₁ in lymphocytes to inhibit their recirculation, thus resulting in lymphopenia and attenuated immune reaction [12]. In addition, animal studies suggest that targeting the S1P signaling system is a promising strategy for inhibiting vascular hyperpermeability and modulating angiogenesis [13, 15, 16]. Here, we will overview the signaling mechanisms underlying S1P regulation of biological functions

and the roles of S1P in diseases.

S1P synthesis and metabolism

S1P is generated within cells through phosphorylation of sphingosine by sphingosine kinase 1 (SphK1) and sphingosine kinase 2 (SphK2) (Figure 1) [1], which share a conserved catalytic domain, but are distinct in other aspects including their structures of non-catalytic domains and expression patterns. S1P is either dephosphorylated by S1P phosphatases (SPP1 & SPP2) [17] and lipid phosphate phosphatases (LPP1~LPP3) [18] to convert into sphingosine, or degraded by S1P lyase (SPL) to ethanolamine phosphate and hexadecenal [2], the latter reaction serving as the exit from sphingolipid metabolic pathway (Fig. 1). SPPs and SPL, both of which are highly specific for S1P, reside predominantly in the endoplasmic reticulum, while LPPs dephosphorylate a range of substrates including S1P.

SphK1-knockout (KO) mice are phenotypically normal except for a 60 % reduction in plasma and serum S1P concentrations compared to wild-type mice [19]. However, tissue S1P levels in SphK1-KO mice are similar to wild-type mice. SphK2-KO mice are also phenotypically normal and exhibit a 25% reduction in plasma S1P concentrations [20]. Thus, SphK1 plays a major role in maintaining plasma and serum S1P tone, and

SphK2 compensates for SphK1 in maintaining tissue S1P in the absence of SphK1.

SPL-KO mice display markedly high levels of S1P in tissues and serum with accumulation of ceramide and long chain bases, resulting in multi-organ damages with pro-inflammatory responses and altered lymphocyte and neutrophil distribution [21, 22].

LPP3 maintains S1P level at a low level in the thymus, thus playing a key role in T lymphocyte exit to the blood from the thymus [23].

SphKs and S1P-metabolizing enzymes play important roles not only in the production and degradation of S1P but also in controlling cellular levels of sphingolipids including sphingosine and its metabolic precursor ceramide. In contrast to S1P, these sphingolipid species exert growth inhibitory and proapoptotic effects when their cellular levels rise, through multiple mechanisms including activation of protein phosphatases and inhibition of Akt [1, 2, 24]. Since S1P exhibits anti-apoptotic or survival effects on a variety of cell types, the balance of S1P and ceramide levels is implicated in the determination of cell fate, i.e. survival or death under certain conditions.

Blood S1P and S1P transporters

In the mammalian body, there is a steep S1P concentration gradient across the

capillary wall [25]: the plasma S1P concentration is around 500 nM, which is considered to be markedly higher than that in the tissue interstitial fluid. The majority of plasma S1P derives from red blood cells [26], which express SphK1 but lack S1P degrading enzymes and thus serve as a supplier of S1P in blood, while the remaining of plasma S1P is released from other cells, particularly endothelial cells [27]. Indeed, anemia causes a reduction in the plasma S1P level. Release of S1P from erythrocytes strictly requires acceptor plasma proteins, mostly HDL and albumin [5]. The major part of plasma S1P is bound to HDL (~60%), albumin (~30%) and other plasma proteins, with only a few percentages of total S1P circulating in a free form. Plasma S1P is crucial in maintaining vascular integrity, which is achieved by endothelial S1P₁-mediated stabilization of adherence junctions [16, 28]. At least a part of beneficial effects of HDL, including activation of eNOS, atheroprotection and myocardial protection from ischemia/reperfusion injury, are suggested to be mediated by HDL-bound S1P [29].

S1P is released out of erythrocytes via a transmembrane S1P transporter. Although ABC family transporters have been implicated in S1P export from erythrocytes, the exact molecular entity of an S1P transporter remains inconclusive [3]. In a zebrafish mutant *miles apart*, loss of function mutation of S1P₂ results in an anomaly termed

cardia bifida (two primordial heart tissues remaining separated) [30]. In a different zebrafish mutant *ko157*, which also shows *cardia bifida*, the major facilitator superfamily type transporter, *Spns2*, was mutated and cardiac defects in *ko157* mutant was rescued by S1P injection [4]. Zebrafish *Spns2* and its mammalian counterpart were found to function as a transporter for S1P and FTY720 phosphate.

Expression of S1P receptors

S1P₁, S1P₂ and S1P₃ are broadly expressed in most of organs and mediate diverse actions of S1P (Table 1) [6-8, 11]. Detailed expression patterns of S1P receptors in tissues were defined by analyzing mice in which β -galactosidase (LacZ) reporter gene was knocked into the receptor gene loci. S1P₁, which was originally cloned from vascular endothelial cells, is detected in the endothelium in the lung, heart and liver of S1P₁^{+LacZ} mice [9]. In the lung, S1P₁ is also expressed in vascular smooth muscle. Unexpectedly, LacZ activity is undetectable in the endothelium of the kidney, spleen and testis. The non-vascular cells, including neuronal cells including Purkinje cells and neurons in the molecular layer, astrocytes, cardiomyocytes, cells in the marginal zone in spleen, and epithelial cells in the renal collecting duct express S1P₁.

In normal tissues of S1P₂^{+LacZ} mice, LacZ activity is detected in various sizes of

blood vessels in a variety of organs, which include lung, brain, skeletal muscle, kidney, and liver [31]. Vascular cells are the major cell types that express S1P₂ in many organs. Histological analysis using combined immunohistochemistry and X-gal staining showed that the endothelium in microvessels and both the endothelium and smooth muscle of larger vessels express S1P₂. In addition, a limited population of bone marrow cells and a small number of non-vascular cells in the brain express S1P₂.

In contrast to S1P₁, S1P₂ and S1P₃, the expression of the other two S1P receptors S1P₄ and S1P₅ is restricted: S1P₄ and S1P₅ are primarily expressed in lymphoid tissues and the lung, and the brain (especially oligodendrocytes), leukocytes and spleen, respectively [8, 11].

Distinct signaling mechanisms of S1P₁, S1P₂ and S1P₃

The signaling mechanisms of S1P₁~ S1P₃ are better characterized compared with S1P₄ and S1P₅. S1P₁, S1P₂ and S1P₃ activate overlapping yet distinctive intracellular signaling pathways, as analyzed by expressing cloned receptors in Chinese hamster ovary (CHO) cells and other cells (Fig. 2) [6-8, 11, 32, 33]. S1P₁ couples exclusively to heterotrimeric G_i to activate Ras/ERK, PI 3-kinase/Akt, and Rho family small GTPase Rac. S1P₁ also moderately activates phospholipase C (PLC) and consequently induces

Ca²⁺ mobilization [6, 7]. In contrast to S1P₁, S1P₂ and S1P₃ couple to multiple G proteins, i.e. G_q, G_i and G_{12/13} [11, 32, 33]. S1P₂ stimulates small GTPase Rho via G_{12/13}, PLC mainly via G_q, ERK via G_i, and JNK and p38 mitogen-activated protein kinase (MAPK) via pertussis toxin (PTX)-insensitive G protein [32]. S1P₂ mediates ERK activation obviously less potently compared with S1P₁ and S1P₃ [7, 33], suggesting inefficient G_i-coupling of S1P₂. Regardless of the G_i-coupling of S1P₂, S1P₂ increases cyclic AMP. This was found to be mediated via G₁₃ [34]. Like S1P₂, S1P₃ also couples to G_q-mediated PLC stimulation, G_{12/13}-mediated Rho stimulation, and G_i-mediated ERK and Rac stimulation [11, 33]. S1P₃ decreases or increases cyclic AMP level, depending on experimental conditions. Although S1P₂ and S1P₃ similarly can couple to G_q, G_i and G_{12/13} when overexpressed, obvious difference in the two receptor subtypes exists in primary cells: mouse embryonic fibroblasts (MEFs) from S1P₂-null mice exhibit impaired Rho activation while PLC activation is not compromised compared with wild-type MEFs [11]. On the other hand, MEFs from S1P₃-null mice show impaired PLC activation with Rho activation and adenylate cyclase inhibition unaffected. Although S1P₃ deletion does not impair Rho activation in MEFs, S1P₂- and S1P₃-double null MEFs completely lack Rho activation response, suggesting that there is partial functional redundancy between S1P₂ and S1P₃. S1P₄ was reported to couple

to G_i and $G_{12/13}$, which mediates ERK activation, PLC stimulation and Rho activation [11]. $S1P_5$ couples to G_i and $G_{12/13}$, resulting in adenylate cyclase inhibition and Ca^{2+} mobilization.

Since $S1P_1$, $S1P_2$ and $S1P_3$ are widely expressed, an integrated outcome of S1P signaling in a given cell type largely depends upon relative expression levels of the S1P receptor subtypes. In addition, ever growing numbers of examples of cross-talks between S1P receptor signaling and growth factor or cytokine receptor signaling have been reported. For example, under certain conditions $S1P_3$ activation leads to activation of TGF β signaling pathway and fibrosis. Update information regarding detailed cross-talk mechanisms is available in recently published excellent reviews [8, 35].

Regulation of cell migration by S1P receptor signaling

Cell migration is a fundamental biological process essential for morphogenesis, angiogenesis, immune surveillance, inflammation, tumor cell invasion and metastasis [36]. It is regulated through receptor-mediated processes in response to a variety of ligands, which are either soluble, bound to extracellular matrix or expressed on cell surface.

One of outstanding biological activities of S1P is the ability to regulate cell

migration either negatively or positively, which was first recognized to be apparently cell type-dependent [37]. S1P potently inhibits cell migration in a variety of tumor cells including B16 melanoma, breast cancer, and glioblastoma cells, as well as vascular smooth muscle cells. By contrast, S1P induces chemotaxis in vascular endothelial cells (ECs) [28], MEFs [11], and T and B lymphocytes [10, 12].

CHO cells are an excellent model for studying mechanism of cell migration [38]. They vigorously exhibit stimulation or inhibition of cell migration, depending on stimuli. In a Boyden chamber assay in which cells are placed in the upper well, either S1P₁ or S1P₃ mediate migration of CHO cells toward S1P in the lower well, i.e. chemotaxis, with typical bell-shaped dose-response curves [38]. In contrast, S1P₂ mediates inhibition of cell migration directed toward a chemoattractant. This S1P₂ effect is dependent on a concentration gradient of S1P: S1P₂ mediates inhibition of cell migration toward a chemoattractant in the lower well, when S1P is placed only in the lower well. If S1P is placed only in the upper well or in both the upper and lower wells, chemotaxis is not suppressed. Therefore, S1P₂ mediates chemorepulsion. Prostaglandin E₂ (PGE₂) and isoproterenol, which elevate the intracellular cyclic AMP level via G_s, also inhibit chemotaxis in CHO cells. However, the inhibitory effects of PGE₂ and isoproterenol are distinct from the S1P₂-mediated effect in that PGE₂ and isoproterenol

effectively inhibit chemotaxis, whether these ligands are placed in the upper well or in both the upper and lower wells [39]. Thus, cell migration inhibition induced by PGE₂ or isoproterenol is not dependent on their concentration gradients and therefore differs from chemorepulsion. Rho family GTPase Rac promotes actin polymerization to induce lamellipodia formation and plays a pivotal role in cell migration. The chemoattractant receptors S1P₁ and S1P₃ mediate Rac activation via G_i, whereas chemorepellant receptor S1P₂ does not [38]. Importantly, S1P₂ but not S1P₁ or S1P₃ inhibits Rac activation induced by a chemoattractant. S1P₂-mediated inhibition of Rac activation and cell migration in response to a chemoattractant is abolished by the expression of dominant negative Rho mutant N¹⁹Rho and inhibition of S1P₂-G_{12/13} coupling, indicating that Rho mediates Rac inhibition in S1P₂-expressing cells. Detailed analysis suggests the involvement of stimulation of Rac GTPase-activating protein (GAP) in Rac inhibition in a manner independent of a Rho kinase [38]. Rac activation by a chemoattractant, whether it is a ligand for a GPCR or a receptor tyrosine kinase, is at least in part dependent on PI 3-kinase. The product of PI 3-kinase produces PI-3,4,5-P₃, which mediates recruitment and activation of signaling molecules including a Rac-guanine nucleotide exchange factor such as Tiam-1. PI-3,4,5-P₃ is de-phosphorylated by the 3'-specific phosphoinositide phosphatase "Phosphatase and

Tensin Homolog Deleted from Chromosome 10" (PTEN). PTEN was found to be stimulated by S1P₂ [40]. However, S1P₂-mediated inhibition of Rac and migration does not seem to involve inhibition of PI 3-kinase or stimulation of PTEN [41]. S1P₂ mediates elevation of cyclic AMP, which could mediate inhibition of cell migration. However, this is also unlikely because different from the cases of PGE₂ and isoproterenol as stated above, S1P₂ activation induces chemorepulsion. Various cells, e.g. endothelial cells and smooth muscle cells, express multiple S1P receptor subtypes. A net effect of S1P on cell migration is likely determined by integration of the counteracting signals input by the chemoattractant receptors S1P₁ and S1P₃ and the chemorepellent receptor S1P₂.

Regulation of vascular formation by S1P receptor signaling

Angiogenesis is a complex process comprising EC proliferation and migration, cell-cell adhesion, and mural cell recruitment [36]. The first discovery of an in vivo angiogenic activity of S1P came from the observation that S1P stimulated angiogenesis in the Matrigel implants in mice. S1P induced directed migration of endothelial cells via G_i and proliferation [28]. S1P also facilitates adherens junction assembly in an S1P₁-G_i-Rac- and S1P₃-G_{12/13}-Rho-dependent manner, leading to stimulation of

capillary-like tube formation. S1P₁-null mouse embryo is defective in recruiting pericytes and SMCs to vessels, i.e. vascular maturation or stabilization [9] (see below for more detail). Conditional EC-specific deletion of S1P₁ results in the similar vascular maturation defect to global S1P₁ deletion, indicating that vessel coverage by mural cells is directed by S1P₁ in ECs. In contrast to S1P₁-null mice, either S1P₂- or S1P₃-single null mice are alive without a vascular formation defect. However, compared with mice null for S1P₁ alone, embryos null for both S1P₁ and S1P₂, null for both S1P₁ and S1P₃, and null for all of S1P₁, S1P₂ and S1P₃ exhibit more severe vascular phenotypes including a vascular maturation defect and hemorrhage with earlier intrauterine death [42]. S1P₁ is the most important receptor for vascular development while S1P₂ and S1P₃ possess partially redundant and cooperative functions in S1P regulation of vascular formation.

S1P signaling is involved in pathological angiogenesis including tumor neovascularization. In a tumor cell implantation model in mice, S1P₁ is upregulated in vessels at sites of tumor implantation [15]. S1P₁ silencing by repeated local injections of S1P₁-specific siRNA suppresses tumor angiogenesis and vascular maturation. Administration of monoclonal anti-S1P neutralizing antibody inhibits tumor growth [43]. The effectiveness of anti-S1P antibody is substantial and more than that obtained

with monoclonal anti-VEGF antibody. This anti-tumor effect is likely due to inhibition of both angiogenesis and tumor cell motility, survival and proliferation [44]. Interestingly, anti-S1P antibody suppresses VEGF- and FGF-induced angiogenesis in Matrigel plugs in mice, suggesting that endogenous S1P plays a permissive role in angiogenesis or functions downstream of VEGF and FGF.

In contrast to S1P₁, S1P₂, which is also expressed in ECs, inhibits growth factor-induced Rac activation, cell migration and capillary-like tube formation via a G_{12/13}/Rho-dependent mechanism [38]. The S1P₂-selective antagonist JTE-013 enhances S1P-induced angiogenesis in Matrigel plugs in mice [36]. In murine retinal angiogenesis model, S1P₂ inhibits post-natal physiological angiogenesis in avascular areas of the retina [45]. Thus, different from S1P₁, S1P₂ is a negative regulator of angiogenesis. S1P₂ deletion enhances angiogenesis in implanted tumors with accelerated tumor growth [31]. In tumors, S1P₂ is expressed in ECs and mural cells in tumor vessels. In S1P₂-null mice, the coverage of tumor neovessels with pericytes and SMCs is enhanced compared with wild-type mice. VEGF- and FGF2-induced microvascular formation and mural cell coverage in matrigel plugs are also enhanced in S1P₂-null mice, suggesting that angiogenesis induced by these growth factors is negatively affected by S1P₂.

The ECs isolated from S1P₂-null mice display altered phenotypes compared with wild-type ECs: S1P₂-null ECs show increased cell proliferation, migration and the formation of tube-like tube structures in response to growth factors compared with wild-type ECs [31]. In S1P₂-null MLECs, two major changes in the intracellular signals are noted. Both the basal and S1P-stimulated activities of Rac are greater in S1P₂-null ECs compared with wild-type ECs. Secondly, in wild-type ECs S1P inhibits VEGF-induced activation of Akt but not ERK whereas S1P fails to inhibit Akt activation in S1P₂-null ECs. Thus, S1P₂ seems to mediate S1P-induced Akt inhibition in wild-type ECs. The Akt inhibition is probably mediated through PTEN stimulation, which reduces amount of PI-3,4,5-P₃ [40]. Thus, S1P₂ inhibition of angiogenesis involves the G_{12/13}-Rho-Rac/PTEN signaling pathway in ECs.

In addition to ECs, S1P₂ is also expressed in CD11b⁺ positive bone marrow-derived cells (BMDCs) in the tumor stroma [31]. Myeloid cells including CD11b⁺ cells participate in tumor angiogenesis through multiple mechanisms [36]. Infiltrating myeloid cells in tumors release pro-angiogenic factors including VEGFs, FGF-2, PDGFs and matrix metalloproteases (MMPs), the enzymes that contribute to angiogenesis through degradation of the extracellular matrix proteins and resultant release of VEGFs and TGFβ that has been deposited in the matrix. A subpopulation of

BMDCs is capable of transdifferentiating into vascular ECs and become incorporated into the new blood vessels in tumors. In S1P₂-null mice, CD11b⁺ cells infiltrating into tumors are increased compared with wild-type mice [31]. Bone marrow chimera experiments document that S1P₂ in BMDCs exerts an inhibitory effect on tumor angiogenesis.

Thus, S1P₂ exerts inhibitory effects on tumor angiogenesis through both the EC-autonomous and myeloid cell-dependent actions. These S1P₂ actions open the possibility of a novel anti-angiogenic therapy to target S1P₂. It is an interesting possibility that S1P receptor subtype-selective pharmacological targeting strategies, i.e. S1P₁ inhibition in combination with S1P₂ activation, could lead to more effective inhibition of tumor angiogenesis. In addition to an expected anti-angiogenic action of S1P₂-selective agonist, S1P₂ stimulation in tumor cells could directly inhibit tumor progression in vivo, leading to inhibition of invasion and metastasis, as previously demonstrated [44, 46].

Regulation of vascular homeostasis by S1P receptor signaling

S1P regulates vascular tone by acting on both the endothelium and smooth muscle through multiple S1P receptors. In ECs, S1P₁ is most abundant with S1P₂ and S1P₃

being expressed at much lower levels, whereas in smooth muscle the expression of S1P₂ and S1P₃ are abundant with S1P₁ expression being very low [13]. S1P-induced relaxation is mediated through its action on the endothelium whereas S1P directly contracts smooth muscle. In ECs, S1P stimulates a calmodulin-dependent enzyme, eNOS, which produces nitric oxide (NO). NO diffuses into the underlying smooth muscle to induce relaxation through generating cyclic GMP. This S1P action is mediated via S1P₁ and S1P₃, which activate Akt through PI 3-kinase to phosphorylate eNOS [47]. S1P₁ and S1P₃ also activate PLC to mobilize Ca²⁺, which fully activates eNOS in concert with Akt-mediated phosphorylation. Although G_q-coupled S1P₃ more robustly activates PLC compared with G_i-coupled S1P₁, the contribution of S1P₁ seems to dominate in eNOS stimulation because S1P₁ expression is higher in ECs compared with S1P₃. In smooth muscle, S1P activates Rho and Rho kinase via S1P₂/S1P₃ and G_{12/13} [41]. Rho kinase phosphorylates the myosin targeting subunit, MYPT1, of myosin phosphatase and the myosin phosphatase inhibitor protein, CPI-17, to inhibit myosin phosphatase. The myosin light chain kinase activation by PLC-Ca²⁺, together with myosin phosphatase inhibition by Rho-Rho kinase, efficiently increases myosin light chain phosphorylation and, thereby, vascular contraction. S1P₂ is also suggested to contribute to vascular tone through a mechanism involving the action on the

endothelium although the precise mechanism remains to be defined [48].

S1P contributes to vascular barrier integrity. Initially, S1P was found to enhance barrier function of an EC monolayer and to protect barrier disruption induced by the edemagenic agent thrombin [49]. This effect is mediated by S1P₁ and, to the lesser extent, S1P₃ through G_i-PI 3-kinase-Rac. In contrast to S1P₁, S1P₂, when overexpressed in vitro in ECs, disrupts barrier integrity via Rho-Rho kinase-PTEN pathway [50].

Endothelial barrier dysfunction, which increases vascular permeability, occurs in inflammation, tumor neovessels and atherosclerotic lesions. Challenge with lipopolysaccharide (LPS) or thrombin induces an increase in pulmonary microvascular permeability. S1P₁^{+/-} mice exhibited reductions in barrier protection by administering a moderate dose of S1P or the S1P₁-selective agonist SEW-2871, after LPS challenge [51]. In contrast, S1P₂^{-/-} mice were protected from LPS-induced barrier disruption compared with wild-type mice. Barrier disruption is also enhanced in SphK1-null mice. Adenoviral transduction of SphK1 into the lung protects mice from barrier disruption whereas that of SphK2 rather augments it, indicating the distinct roles of SphK1 and SphK2 [52]. The intravenous or intratracheal administration of S1P is protective against LPS-induced barrier disruption [51]. However, a higher dose of S1P or repeated administration of S1P₁ agonists (FTY720 and AUY954) rather exacerbates barrier

disruption by stimulating internalization and degradation of S1P₁ protein in a lung injury model [53], highlighting the importance of S1P₁ agonist concentration.

Plasma S1P concentration is another critical determinant for maintaining barrier integrity. In inducibly SphK1-deleted mice with SphK1^{fl/-}:SphK2^{-/-}/Mx1-Cre Tg+ (S1Pless mice), which show approximately 30 nM plasma S1P compared with 2.5 μM in control mice, vascular leak on anaphylaxis and administration of platelet-activating factor or histamine is augmented with impaired survival [16]. Transfusion of erythrocytes, which restores plasma S1P levels, or acute administration of an S1P₁ agonist reverse vascular leak and prevent death. In contrast, SphK2-null mice have a rapid recovery from anaphylaxis [54]. S1P₂- but not S1P₃- null mice also show poor recovery from anaphylaxis. S1P infusion fails to promote recovery of S1P₂-null mice from anaphylaxis.

Physiological levels of endothelial S1P₁ and SphK1-produced S1P serve a constitutive maintaining role for vascular barrier function. Exogenous supraphysiological S1P₁ agonists impair this mechanism by downregulating S1P₁. Furthermore, S1P₂ participates in the vascular protection from anaphylaxis although the precise mechanism of the S1P₂ action remains to be fully defined.

Modulation of leukocyte functions and inflammation by S1P signaling

The role of S1P signaling as significant modulator of leukocyte functions and inflammation has emerged. SphK1-derived S1P regulates pro-inflammatory signaling pathways, including activation of nuclear factor- κ B [55]. S1P₁ regulates endothelial barrier integrity as stated above [49-52, 56]; cytokine and adhesion molecule expression, lymphocyte maturation, differentiation and trafficking, and mast cell migration. S1P₂ also regulates B lymphocyte survival and confinement in lymph node follicles [57]. S1P₃ modulate dendritic cell trafficking and activation. In addition, S1P₅ regulates NK cell trafficking [12].

S1P₁-G_i signaling pathway regulates trafficking of lymphocytes and other immune cells by directing migration of immune cells toward a compartment with a relatively higher S1P concentration. Therefore, the existence of a S1P concentration gradient between compartments, e.g. lymphoid tissue parenchyma and blood plasma/lymphatic fluid, which is created and maintained by the SphK-catalyzed S1P production by erythrocytes and vascular/lymphatic endothelial cells and SPL- and LPP3-catalyzed S1P degradation in lymphoid tissue parenchyma, is critical. S1P₁ expression on the cell surface of lymphocytes and other immune cells is maintained in a low S1P environment in the thymus and lymph nodes, through its inhibited internalization/degradation or

upregulation as a result of lymphocyte maturation and interaction with other immune cells within lymphoid tissues [12]. S1P₁ also participates in the regulation of lymphocyte recirculation through tightening the cell-cell junction of sinus-lining ECs [56]. S1P₂-G_{12/13} pathway ensures the localization of S1P₂-expressing B cells in a follicular center in lymph nodes [57]: S1P concentration is higher at the follicle perimeter than the follicular center due to S1P production by stromal cells abundant at the perimeter and rapid S1P degradation by follicular B cells in the center. In the presence of this S1P concentration gradient, migration of S1P₂-expressing B cells from the center to the perimeter of a follicle is impeded by the chemorepellent activity of S1P₂ through Rho-induced Rac inhibition. The low S1P environment at the follicular center also favors survival and proliferation of S1P₂-expressing B cells because the mitogenic and survival signaling molecule Akt, which is negatively regulated by S1P₂-G_{12/13}-Rho-PTEN, is spared from suppression.

SphK1 are involved in inflammation through both the extracellular messenger and intracellular messenger actions of S1P [12]. In a septic model due to bacterial peritonitis, thrombin, which is produced by coagulation reaction, binds to and activates the GPCR propease-activated receptor-1 (PAR1) on dendritic cells involved in innate immunity [58]. The activation of PAR1 in turn stimulates SphK1, S1P export to the cell exterior,

and S1P₃ activation, which induces amplification of inflammation by stimulating the production of IL-1 and tissue factor from dendritic cells and disrupting EC barrier function. SphK1 is also implicated in the actions of tumor necrosis factor (TNF) and other cytokines, in which intracellular S1P produced by SphK1 binds to TRAF2 and thereby activates NF- κ B [55]. Disruption of SphK1 gene alleviates inflammatory diseases including colitis and arthritis, providing further support for the involvement of SphK1 in inflammatory responses [12]. In addition to the intracellular action of SphK1-generated S1P, a recent study [59] showed that S1P produced by SphK2 in the nucleus bound to the histone deacetylases HDAC1 and HDAC2 and inhibited their enzymatic activity, which suggested that HDACs are direct intracellular targets of S1P. Furthermore, S1P generated by SphK2 in mitochondria plays the important role in cytochrome-c oxidase assembly and respiration [60].

Conclusion

There is now broad consensus that S1P signaling plays a crucial role in the physiology and pathophysiology of the cardiovascular, immune and other systems. Observations obtained with gene-engineered mice and pharmacological tools to target

receptors and enzymes rapidly promote our understanding S1P functions. Investigation in more depth into involvements of S1P signaling in various diseases, in combination with development of drugs with improved specificity and efficacy and their optimal drug delivery system, will provide new treatment strategies.

Acknowledgements

This study was supported by Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science (Y.T., N.T. Y.O., K.Y.), Grant-in-Aid for Scientific Research on Priority Areas (Y.T.) from the Ministry of Education, Culture, Sports, Science and Technology in Japan, funds from the Kanazawa University Strategic Research Development Program (Y.T.), and the IPNU Research Promotion Program (N.T.).

References

- [1] S. Spiegel, S. Milstien (2007) Functions of the multifaceted family of sphingosine kinases and some close relatives, *J. Biol. Chem.* 282, 2125-2129.
- [2] H.Fyrst, J. Saba (2008) Sphingosine-1-phosphate lyase in developmental and disease: Sphingolipid metabolism takes flight, *Biochim. Biophys. Acta.* 1781, 448-458.
- [3] R.H. Kim, K. Takabe, S. Milstien, S. Spiegel (2009) Export and functions of sphingosine-1-phosphate, *Biochim. Biophys. Acta.* 1791, 692-699.
- [4] A. Kawahara, T. Nishi, Y. Hisano, H. Fukui, A. Yamaguchi, N. Mochizuki (2009) The sphingolipid transporter spns2 functions in migration of zebrafish myocardial precursors , *Science.* 323, 524-527.
- [5] F. Okajima (2002) Plasma lipoproteins behave as carriers of extracellular sphingosine 1-phosphate: is this an atherogenic mediator or an anti-atherogenic mediator?, *Biochim Biophys Acta.* 1582, 132-137.
- [6] M. J. Lee, J. R. Van Brocklyn, S. Thangada, C. H. Liu, A. R. Hand, R. Menzeleev, S. Spiegel, T. Hla, Sphingosine-1-phosphate as a ligand for the G protein-coupled receptor EDG-1. *Science.* 279, 1552-5.
- [7] H. Okamoto, N. Takuwa, K. Gonda, H. Okazaki, K. Chang, Y. Yatomi, H. Shigematsu and Y. Takuwa (1998) EDG1 is a functional sphingosine-1-phosphate receptor that is linked via a Gi/o to multiple signaling pathways, including phospholipase C activation, Ca²⁺ mobilization, Ras-mitogen-activated protein kinase activation, and adenylate cyclase inhibition. *J. Biol. Chem.* 273, 27104-10.
- [8] J. Chun, T. Hla, K. R. Lynch, S. Spiegel, W. H. Moolenaar (2010) International Union of Basic and Clinical Pharmacology. LXXVIII. Lysophospholipid receptor nomenclature. *Pharmacol. Rev.* 62, 579-587.
- [9] Y. Liu, R. Wada, T. Yamashita, Y. Mi, C. X. Deng, J. P. Hobson, H. M. Rosenfeldt , V. E. Nava, S. S. Chae, M. J. Lee, C. H. Liu, T. Hla, S. Spiegel, R. L. Proia (2000) Edg-1, the G protein-coupled receptor for sphingosine-1-phosphate, is essential for vascular maturation. *J. Clin. Invest.* 106, 951-961.

- [10] M. Matloubian, C. G. Lo, G. Cinamon, M. J. Lesneski, Y. Xu, V. Brinkmann, A. L. Allende, R. L. Proia, J. G. Cyster (2004) Lymphocyte egress from thymus and peripheral lymphoid organs is dependent on S1P receptor 1. *Nature*. 427, 355-360.
- [11] I. Ishii, N. Fukushima, X. Ye, J. Chun (2004) Lysophospholipid receptors: signaling and biology, 73, 321-354.
- [12] S. Spiegel, S. Milstien (2011) The outs and the ins of sphingosine-1-phosphate in immunity. *Nat. Rev. Immunol.* 11, 403-415.
- [13] C. K. Means, J. H. Brown (2009) Sphingosine-1-phosphate receptor signaling in the heart. *Cardiovasc. Res.* 82, 193-200.
- [14] L. Kappos, J. Antel, G. Comi, X. Montalban, P. O'Connor, C. H. Polman, T. Haas, A. A. Korn, G. Karlsson, E. W. Radue; FTY720 D2201 Study Group (2006) Oral fingolimod (FTY720) for relapsing multiple sclerosis. *N Engl. J. Med.* 355, 1124-1140.
- [15] S. S. Chae, J. H. Paik, H. Furneaux, T. Hla (2004) Requirement for sphingosine 1-phosphate receptor-1 in tumor angiogenesis demonstrated by in vivo RNA interference. *J. Clin. Invest.* 114, 1082-1089.
- [16] E. Camerer, J. B. Regard, I. Cornelissen, Y. Srinivasan, D. N. Duong, D. Palmer, T. H. Pham, J. S. Wong, R. Pappu, S. R. Coughlin. (2009) Sphingosine-1-phosphate in the plasma compartment regulates basal and inflammation-induced vascular leak in mice. *J Clin Invest.* 119, 1871-1879.
- [17] S. M. Mandala, R. Thornton, I. Galve-Roperh, S. Poulton, C. Peterson, A. Olivera, J. Bergstrom, M. B. Kurtz, S. Spiegel (2000) Molecular cloning and characterization of a lipid phosphohydrolase that degrades sphingosine-1-phosphate and induces cell death. *Proc. Natl. Acad. Sci. U. S. A.* 97, 7859-7864.
- [18] D. W. Waggoner, J. Xu, I. Singh, R. Jasinska, Q. X. Zhang, D. N. Brindley (1999) Structural organization of mammalian lipid phosphate phosphatases: implications for signal transduction. *Biochim. Biophys. Acta.* 1439, 299-316.

- [19] M. L. Allende, T. Sasaki, H. Kawai, A. Olivera, Y. Mi, G. van Echten-Deckert, R. Hajdu, M. Rosenbach, C. A. Keohane, S. Mandala, S. Spiegel, R. L. Proia RL. (2004) Mice deficient in sphingosine kinase 1 are rendered lymphopenic by FTY720. *J. Biol. Chem.* 279, 52487-5292.
- [20] Y. Kharel, S. Lee, A. H. Snyder, S. L. Sheasley-O'Neill, M. A. Morris, Y. Setiady, R. Zhu, M. A. Zigler, T. L. Burcin, K. Ley, K. S. Tung, V. H. Engelhard, Macdonald, S. Pearson-White S, K. R. Lynch. (2005) Sphingosine kinase 2 is required for modulation of lymphocyte traffic by FTY720, *J. Biol. Chem.* 280, 36865-36872.
- [21] P. Vogel, M. S. Donoviel, R. Read, G. M. Hansen, J. Hazlewood, S. J. Anderson, W. Sun, J. Swaffield, T. Oravec (2009) Incomplete inhibition of sphingosine 1-phosphate lyase modulates immune system function yet prevents early lethality and non-lymphoid lesions, *PLoS One.* 4, e4112.
- [22] M. Bektas, M. L. Allende, B. G. Lee, W. Chen, M. J. Amar, A. T. Remaley, J. D. Saba, R. L. Proia. (2010) Sphingosine 1-phosphate lyase deficiency disrupts lipid homeostasis in liver. *J. Biol. Chem.* 285, 10880-10889.
- [23] B. Bréart, W. D. Ramos-Perez, A. Mendoza, A. K. Salous, M. Gobert, Y. Huang, R. H. Adams, J. J. Lafaille, D. Escalante-Alcalde, A. J. Morris, S. R. Schwab. (2011) Lipid phosphate phosphatase 3 enables efficient thymic egress, *J. Exp. Med.* 208, 1267-1278.
- [24] N. Bartke, Y. A. Hannun. Bioactive sphingolipids: metabolism and function. (2009) *J. Lipid Res.* 50, Suppl:S91-96.
- [25] S. R. Schwab, J. P. Pereira, M. Matloubian, Y. Xu, Y. Huang, J. G. Cyster JG. (2005) Lymphocyte sequestration through S1P lyase inhibition and disruption of S1P gradients. *Science.* 309, 1735-1739.
- [26] R. Pappu, S. R. Schwab, I. Cornelissen, J. P. Pereira, J. B. Regard, Y. Xu, E. Camerer, Y. W. Zheng, Y. Huang, J. G. Cyster, S. R. Coughlin SR. (2007) Promotion of lymphocyte egress into blood and lymph by distinct sources of sphingosine-1-phosphate. *Science.* 316, 295-298.

- [27] K. Venkataraman, Y. M. Lee, J. Michaud, S. Thangada, Y. Ai, H. L. Bonkovsky, N. S. Parikh, C. Habrukowich, T. Hla (2008) Vascular endothelium as a contributor of plasma sphingosine 1-phosphate, *Circ. Res.* 102, 669-676.
- [28] M. J. Lee, S. Thangada, K. P. Claffey, N. Ancellin, C. H. Liu, M. Kluk, M. Volpi, R. I. Sha'afi, T. Hla (1999) Vascular endothelial cell adherens junction assembly and morphogenesis induced by sphingosine-1-phosphate. *Cell.* 99, 301-312.
- [29] J. R. Nofer, M. van der Giet, M. Tölle, I. Wolinska, K. von Wnuck Lipinski, H. A. Baba, U. J. Tietge, A. Gödecke, I. Ishii, B. Kleuser, M. Schäfers, M. Fobker, W. Zidek, G. Assmann, J. Chun, B. Levkau. (2004) HDL induces NO-dependent vasorelaxation via the lysophospholipid receptor S1P3. *J. Clin. Invest.* 113, 569-581.
- [30] E. Kupperman, S. An, N. Osborne, S. Waldron, D. Y. Stainier. A (2000) Sphingosine-1-phosphate receptor regulates cell migration during vertebrate heart development. *Nature.* 406, 192-195.
- [31] W. Du, N. Takuwa, K. Yoshioka, Y. Okamoto, K. Gonda, K. Sugihara, A. Fukamizu, M. Asano, Y. Takuwa (2010) S1P2, the G protein-coupled receptor for sphingosine-1-phosphate, negatively regulates tumor angiogenesis and tumor growth in vivo in mice, *Cancer Res.* 70, 772-781.
- [32] K. Gonda, H. Okamoto, N. Takuwa, Y. Yatomi, H. Okazaki, T. Sakurai, S. Kimura, R. Sillard, K. Harii and Y. Takuwa. (1999) The novel sphingosine 1-phosphate receptor AGR16 is coupled via pertussis toxin-sensitive and -insensitive G-proteins to multiple signalling pathways. *Biochem. J.* 337, 67-75.
- [33] H. Okamoto, N. Takuwa, Y. Yatomi, K. Gonda, H. Shigematsu and Y. Takuwa. (1999) EDG3 is a functional receptor specific for sphingosine-1-phosphate and sphingosylphosphorylcholine with signaling characteristics distinct from EDG1 and AGR16. *Biochem. Biophys. Res. Commun.* 260, 203-208.
- [34] L. I. Jiang, J. Collins, R. Davis, K. M. Lin, D. DeCamp, T. Roach, R. Hsueh, R. A. Rebres, E. M. Ross, R. Taussig, I. Fraser, P. C. Sternweis. (2007) Use of a cAMP BRET sensor to characterize a novel regulation of cAMP by the sphingosine 1-phosphate/G13 pathway. *J. Biol. Chem.* 282, 10576-10584.

- [35] N. J. Pyne, S. Pyne. (2010) Sphingosine 1-phosphate and cancer. *Nat. Rev. Cancer.* 10, 489-503.
- [36] N. Takuwa, W. Du, E. Kaneko, Y. Okamoto, K. Yoshioka, Y. Takuwa (2011) Tumor-suppressive Sphingosine-1-phosphate Receptor-2 Counteracting Tumor-promoting Sphingosine-1-phosphate Receptor-1 and Sphingosine Kinase 1-Jekyll Hidden behind Hyde. *Am. J. Cancer Res.* 1, 460-481.
- [37] Y. Sadahira, F. Ruan, S. Hakomori, Y. Igarashi (1992) Sphingosine 1-phosphate, a specific endogenous signaling molecule controlling cell motility and tumor cell invasiveness. *Proc. Natl. Acad. Sci. U. S. A.* 89, 9686-9690.
- [38] H. Okamoto, N. Takuwa, T. Yokomizo, N. Sugimoto, S. Sakurada, H. Shigematsu and Y. Takuwa (2000) Inhibitory Regulation of Rac Activation, Membrane Ruffling and Cell Migration by Sphingosine-1-Phosphate Receptor EDG5, but not EDG1 or EDG3, *Mol. Cell. Biol.* 20, 9247-9261.
- [39] S. Nagasawa, N. Takuwa, N. Sugimoto, H. Mabuchi and Y. Takuwa (2005) Inhibition of Rac activation as a mechanism for negativeregulation of actin cytoskeletal reorganization and cell motility by cyclic AMP, *Biochem. J.* 385, 737-744.
- [40] T. Sanchez, S. Thangada, M. T. Wu, C. D. Kontos, D. Wu, H. Wu, T. Hla. (2005) PTEN as an effector in the signaling of antimigratory G protein-coupled receptor. *Proc. Natl. Acad. Sci. U. S. A.* 102, 4312-4317.
- [41] Y. Takuwa, Y. Okamoto, K. Yoshioka, N. Takuwa (2008) Sphingosine-1-phosphate signaling and biological activities in the cardiovascular system *Biochim. Biophys. Acta.* 1781, 483-488.
- [42] M. Kono, Y. Mi, Y. Liu, T. Sasaki, M. L. Allende, Y. P. Wu, T. Yamashita, R. L. Proia (2004) The sphingosine-1-phosphate receptors S1P1, S1P2, and S1P3 function coordinately during embryonic angiogenesis. *J. Biol. Chem.* 279, 29367-29373.
- [43] B. Visentin, J. A. Vekich, B. J. Sibbald, A. L. Cavalli, K. M. Moreno, R. G. Matteo, W. A. Garland, Y. Lu, S. Yu, H. S. Hall, V. Kundra, G. B. Mills, R. A. Sabbadini (2006)

Validation of an anti-sphingosine-1-phosphate antibody as a potential therapeutic in reducing growth, invasion, and angiogenesis in multiple tumor lineages. *Cancer Cell.* 9, 225-238.

[44] H. Yamaguchi, J. Kitayama, N. Takuwa, K. Arikawa, I. Inoki, K. Takehara, H. Nagawa and Y. Takuwa. (2003) Sphingosine-1-phosphate receptor subtype-specific positive and negative regulation of Rac and hematogenous metastasis of melanoma cells *Biochem. J.* 374, 715-722.

[45] A. Skoura, T. Sanchez, K. Claffey, S.M. Mandala, R.L. Proia, T. Hla (2007) Essential role of sphingosine 1-phosphate receptor 2 in pathological angiogenesis of the mouse retina, *J Clin Invest.* 117, 2506-2516.

[46] G. Cattoretti, J. Mandelbaum, N. Lee, A. H. Chaves, A. M. Mahler, A. Chadburn, R. Dalla-Favera, L. Pasqualucci, A. J. MacLennan. (2009) Targeted disruption of the S1P2 sphingosine 1-phosphate receptor gene leads to diffuse large B-cell lymphoma formation, *Cancer Res.* 69, 8686-8692.

[47] J. Igarashi, T. Michel (2009) Sphingosine-1-phosphate and modulation of vascular tone, *Cardiovasc. Res.* 82, 212-220.

[48] J. N. Lorenz, L. J. Arend, R. Robitz, R. J. Paul RJ, A. J. MacLennan (2007) Vascular dysfunction in S1P2 sphingosine 1-phosphate receptor knockout mice. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 292, R440-446.

[49] J. G. Garcia, F. Liu, A. D. Verin, A. Birukova, M. A. Dechert, W. T. Gerthoffer, J. R. Bamberg, D. English (2001) Sphingosine 1-phosphate promotes endothelial cell barrier integrity by Edg-dependent cytoskeletal rearrangement. *J. Clin. Invest.* 108, 689-701.

- [50] T. Sanchez, A. Skoura, M. T. Wu, B. Casserly, E. O. Harrington, T. Hla (2007) Induction of vascular permeability by the sphingosine-1-phosphate receptor-2 (S1P2R) and its downstream effectors ROCK and PTEN. *Arterioscler. Thromb. Vasc. Biol.* 27, 1312-1318.
- [51] S. Sammani, L. Moreno-Vinasco, T. Mirzapoiazova, P. A. Singleton, E. T. Chiang, C. L. Evenoski, T. Wang, B. Mathew, A. Husain, J. Moitra, X. Sun, L. Nunez, J. R. Jacobson, S. M. Dudek, V. Natarajan, J. G. Garcia (2010) Differential effects of sphingosine 1-phosphate receptors on airway and vascular barrier function in the murine lung. *Am. J. Respir. Cell. Mol. Biol.* 43, 394-402.
- [52] R. Wadgaonkar, V. Patel, N. Grinkina, C. Romano, J. Liu, Y. Zhao, S. Sammani, J. G. Garcia, V. Natarajan (2009) Differential regulation of sphingosine kinases 1 and 2 in lung injury. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 296, L603-613.
- [53] M. L. Oo, S. H. Chang, S. Thangada, M. T. Wu, K. Rezaul, V. Blaho, S. I. Hwang, D. K. Han, T. Hla (2011) Engagement of S1P1-degradative mechanisms leads to vascular leak in mice. *J Clin Invest.* 121, 2290-2300.
- [54] A. Olivera, C. Eisner, Y. Kitamura, S. Dillahun, L. Allende, G. Tuymetova, W. Watford, F. Meylan, S. C. Diesner, L. Li, J. Schnermann, R. L. Proia, J. Rivera (2010) Sphingosine kinase 1 and sphingosine-1-phosphate receptor 2 are vital to recovery from anaphylactic shock in mice. *J. Clin. Invest.* 120, 1429-1440.
- [55] Alvarez SE, Harikumar KB, Hait NC, Allegood J, Strub GM, Kim EY, Maceyka M, Jiang H, Luo C, Kordula T, Milstien S, Spiegel S. Sphingosine-1-phosphate is a missing cofactor for the E3 ubiquitin ligase TRAF2. *Nature.* 2010; 465: 1084-8.
- [56] H. Rosen, M. G. Sanna, S. M. Cahalan, P. J. Gonzalez-Cabrera (2007) Tipping the gatekeeper: S1P regulation of endothelial barrier function. *Trends Immunol.* 28, 102-107.

- [57] J. A. Green, K. Suzuki, B. Cho, L. D. Willison, D. Palmer, C. D. Allen, T. H. Schmidt, Y. Xu, R. L. Proia, S. R. Coughlin, J. G. Cyster (2011) The sphingosine 1-phosphate receptor S1P₂ maintains the homeostasis of germinal center B cells and promotes niche confinement. *Nat. Immunol.* 12, 672-680.
- [58] F. Niessen, F. Schaffner, C. Furlan-Freguia, R. Pawlinski, G. Bhattacharjee, J. Chun, C. K. Derian, P. Andrade-Gordon, H. Rosen, W. Ruf (2008) Dendritic cell PAR1-S1P3 signalling couples coagulation and inflammation. *Nature.* 452, 654-658.
- [59] N. C. Hait, J. Allegood, M. Maceyka, G. M. Strub, K. B. Harikumar, S. K. Singh, C. Luo, R. Marmorstein, T. Kordula, S. Milstien, S. Spiegel (2009) Regulation of histone acetylation in the nucleus by sphingosine-1-phosphate. *Science* 325:1254-1257.
- [60] G. M. Strub, M. Paillard, J. Liang, L. Gomez, J. C. Allegood, N. C. Hait, M. Maceyka, M. M. Price, Q. Chen, D. C. Simpson, T. Kordula, S. Milstien, E. J. Lesnefsky, S. Spiegel (2011) Sphingosine-1-phosphate produced by sphingosine kinase 2 in mitochondria interacts with prohibitin 2 to regulate complex IV assembly and respiration. *Faseb J* 25:600-612.

Figure legends

Figure 1. Sphingolipid metabolism in various subcellular compartments

Ceramide (Cer) is produced either by de novo synthesis from palmitoyl CoA (palmCoA) and serine with sequential enzymatic reactions in endoplasmic reticulum (ER) or through degradation of sphingomyelin (SM) by the action of sphingomyelinases in the plasma membrane and intracellular membranes including lysosomes. Cer is deacylated by ceramidase to yield sphingosine (Sph), which is then phosphorylated by SphK1/2 to generate S1P. S1P is exported through a plasma membrane S1P transporter, leading to activation of the G protein-coupled S1P receptor subtypes (S1P₁~S1P₅). S1P could be either dephosphorylated by S1P phosphatase1/2 (SPP) and lipid phosphate phosphatase1-3 (LPP) back to Sph or degraded to ethanolamine-phosphate (Eth-P) and hexadecenal (hxdcnl) by S1P lyase (SPL) to leave sphingolipid metabolic pathway. SphK1 is present in both cytosolic and membrane-bound fractions, both being enzymatically active. SPPs and SPL are located in ER. At least, a subtype of LPPs exists on the plasma membrane. Intracellular transfer of Cer from ER to Golgi is facilitated by transfer proteins such as CERT, and both Cer and SM traffic between membrane compartments via vesicular transport.

Figure 2. S1P receptor subtype-specific heterotrimeric G protein coupling and intracellular signaling mechanisms

S1P₁ couples exclusively to G_i to activate Ras-ERK and PI 3-kinase-Akt/Rac pathways, leading to stimulation of chemotaxis and cell proliferation. S1P₂ couples to multiple G proteins, especially to G_{12/13} to induce robust Rho activation, leading to inhibition of Rac and cell migration, and also inhibition of cell proliferation via inhibition of Akt. S1P₂ also couples to stimulation of adenylate cyclase via G₁₃. S1P₃ activates G_q-PLC-Ca²⁺ pathway, and G_i-Ras-ERK and G_i-PI 3-kinase-Akt/Rac pathways. S1P₃-G_{12/13}-Rho pathway becomes evident only when G_i is inhibited by pertussis toxin.

Table 1. Characteristics of S1P receptors

Receptor	G protein coupling	Expression	Agonist	Antagonist	Phenotypes of knockout mice
S1P₁	G _i	Widespread	FTY720-P SEW2871 AUY954 AAL-R CYM5442 KRP-203 RP-002	W146 VPC-23019 VPC-44116	Embryonic lethal Impaired vascular maturation Abnormal brain development Abnormal lymphocyte trafficking
S1P₂	G _{12/13} G _q G _i	Widespread	None	JTE-013	Seizure Hearing loss Vestibular ataxia Decreased blood pressure High incidence of B cell lymphoma Small litter size
S1P₃	G _q G _{12/13} G _i	Widespread	FTY720-P AAL-R KRP-203	VPC-23019 VPC-44116	Attenuated inflammation in sepsis Decreased nitric oxide production in vascular endothelium Disappearance of pressor response and bradycardia to S1P
S1P₄	G _i G _{12/13}	Lymphoid tissues, lung	FTY720-P AAL-R KRP-203	None	Unknown
S1P₅	G _i G _{12/13}	Brain, spleen	FTY720-P AAL-R KRP-203	None	Abnormal trafficking of NK cells