The fast emergency vehicle pre-emption system improved the outcomes of out-of-hospital cardiac arrest

著者	Tanaka Yoshio, Yamada Hiroshi, Tamasaku Shuji, Inaba Hideo
journal or publication title	American Journal of Emergency Medicine
volume	31
number	10
page range	1466-1471
year	2013-10-01
URL	http://hdl.handle.net/2297/35643

doi: 10.1016/j.ajem.2013.07.031

The Fast Emergency Vehicle Pre-emption System Improved the Outcomes of Out-of-hospital

Cardiac Arrest

Short title: OHCA outcomes improved by FAST™

Yoshio Tanaka, MD, MSc

Department of Emergency Medical Science, Kanazawa University Graduate School of Medicine

Hiroshi Yamada

Kanazawa city Fire Department

Shuji Tamasaku

Kanazawa city Fire Department

Hideo Inaba, MD, PhD

Professor and Chairman

Department of Emergency Medical Science, Kanazawa University Graduate School of Medicine

Reprint requests: Hideo Inaba, Department of Emergency Medical Science,

Kanazawa University Graduate School of Medicine 13-1 Takara-machi, Kanazawa, Ishikawa,

920-8641, Japan

Phone +81 76-265-2825 Fax +81 76-234-4243 Email: hidinaba@med.kanazawa-u.ac.jp

Name of grant: None.

A part of the work described in the manuscript has been formally poster presented in English at the “31st International Symposium on Intensive Care and Emergency Medicine” March 22–25, 2011 and oral presented in Japanese at “The 39th annual meeting of the Japanese Association for Acute Medicine” October 20, 2011.

Keywords: ambulance response time; out-of-hospital cardiac arrest; traffic control system; emergency medical service

Total word count: 2639, Number of tables: 3, Number of figures: 2, Number of references: 33

Corresponding author:

Hideo Inaba, MD, PhD Professor and Chairman

Department of Emergency Medical Science,

Kanazawa University Graduate School of Medicine

13-1 Takara-machi, Kanazawa, 920-8641, Japan

Phone +81 76-265-2825

Fax +81 76-234-4243

Email: hidinaba@med.kanazawa-u.ac.jp

Abstract

Purposes: Ambulance response time is a major factor associated with survival in out-of-hospital cardiac arrests (OHCAs); the fast emergency vehicle pre-emption system (FASTTM) aids response time by controlling traffic signals. This eight-year observational study investigated whether FASTTM implementation reduced response times and improved OHCA outcomes.

Methods: Data was prospectively collected from 1161 OHCAs that were not witnessed by emergency medical technicians (EMTs) from April 1, 2003 to March 31, 2011. The study took place in Kanazawa city, where ambulances without FASTTM (non- FASTTM -equipped) were being progressively replaced by new FASTTM -equipped ambulances. OHCA data, including the response times recorded in seconds, were collected and compared between the FASTTM -equipped and non- FASTTM -equipped ambulances. OHCA outcomes were subsequently compared in the subgroup of OHCAs managed by EMTs without tracheal intubation or epinephrine administration. The primary end-point of this study was one-year (1-Y) survival.

Results: The median response time significantly differed between the FASTTM -equipped and non-FASTTM -equipped groups at 327 and 381 s, respectively. The 1-Y survival rates were 7.0% in the FASTTM -equipped group and 2.8% in the non- FASTTM -equipped group. Logistic regression analysis revealed that the dispatch of a FASTTM -equipped ambulance was an independent factor for 1-Y survival (adjusted odds ratio = 3.077, 95% confidence interval = 1.180–9.350).

Conclusions: The FASTTM implementation significantly reduced ambulance response times and improved OHCA outcomes in Kanazawa city. (Word count: 231)

1. Introduction

The response time, defined as the interval between call for and arrival of an ambulance, is one of the major factors associated with favorable outcomes of out-of-hospital cardiac arrests (OHCAs) [1, 2]. Although reducing the response time may improve OHCA outcomes [3-7], there are only a few ways to achieve it. For example, response times can be improved by increasing the number of ambulance teams and fire department offices, or by equipping additional first-line responders with defibrillators, such as the fire fighters and police services [8, 9].

The fast emergency vehicle pre-emption system is a component of the universal traffic control system (UTMS) used in Japan, and is officially termed FASTTM by the UTMS society of Japan [10]. FASTTM minimizes emergency vehicle transit time by controlling traffic signals [11], thereby offering a potential approach to reduce ambulance response times. However, the effect of FASTTM on OHCA outcomes has not been investigated to date. This study therefore aimed to determine whether the implementation of FASTTM reduced ambulance response times and, in turn, improved outcomes.

2. Methods

An eight-year prospective, observational study was designed to evaluate the impact of FASTTM implementation on the emergency medical service (EMS) response times and OHCA outcomes. All data were collected in accordance with the national guideline of ethics for epidemiological surveys [12]. This study was approved by a review board at the Ishikawa Medical Control Council.

Patient data

Kanazawa city Fire Department prospectively collected data in accordance with the Utstein recommendation [13, 14]. In central Kanazawa city, data for all OHCA with attempted resuscitation and those who were transported to hospitals were collected from April 1, 2003 to March 31, 2011. The following data were collected: region; arrest location; patient age and gender; witness of arrest; arrest etiology; cardiopulmonary resuscitation (CPR) before emergency medical technician (EMT) arrival; initial cardiac rhythm; various time factors including response time and duration of transportation to hospital (defined as the time interval between ambulance departure from the arrest scene and arrival at hospital); sustained return of spontaneous circulation (SROSC); one-year (1-Y) survival; and 1-Y survival with a favorable neurological outcome (cerebral performance score = 1 or 2) [15]. SROSC was defined as the continuous presence of palpable pulses for at least 20 min [13, 14]. Survival at 1-Y was defined as being alive in the hospital at 1-Y or discharged alive from the hospital to home or to care and rehabilitation facilities within 1-Y. The primary end point was 1-Y survival. The secondary end points were SROSC and 1-Y survival with a favorable neurological outcome.

Populations and setting

Kanazawa city, the capital of Ishikawa Prefecture, covers 468 km² on the western coast of Honshu, the main island of Japan, and has a population of 461,700. The city is a historical castle town, and the streets in the central area are often congested. The city has eight ambulance stations, each with a one-tiered ambulance system controlled by a single dispatch center and the same level of EMT team is dispatched to all emergency cases. The number of dispatched cases (number of dispatch to

OHCAs/total dispatch) during the study period was 224/11951 (1.9%) in 2003 (fiscal year beginning on April 1); 204/12870 (1.6%) in 2004; 206/12894 (1.6%) in 2005; 210/13328 (1.6%) in 2006; 223/14155 (1.6%) in 2007; 247/13694 (1.8%) in 2008; 302/13890 (2.1%) in 2009; and 273/13942 (2.0%) in 2010.

Telephone-assisted CPR instruction was regularly and strictly conducted by a dispatcher. EMTs resuscitated OHCA patients according to the protocol developed by the Ishikawa Medical Control Council from the guidelines of the American Heart Association and the Japan Resuscitation Council, unless OHCA patients had post-mortem changes. Paramedics were included in all ambulance teams and were authorized to perform the following procedures during resuscitation: (a) use of airway adjuncts, including the supraglottic airway or laryngeal mask airway, (b) infusion of Ringer's lactate through a peripheral vein, and (c) use of semi-automated external defibrillators. Since July 2004, specially trained paramedics have been permitted to insert endotracheal tubes, and since April 2006, they have been permitted to administer intravenous epinephrine. Strict criteria limited the use of these pre-hospital advanced cardiac life support (ACLS) procedures (Table 1) [16]. EMTs are not permitted to terminate resuscitation in the field.

Fast emergency vehicle pre-emption system

FASTTM is one component of the UTMS [10] that minimizes the transit time of emergency vehicles by controlling traffic signals; an animated presentation, demonstrating FASTTM can be seen on the UTMS website [11]. The system includes an infrared beacon that recognizes emergency vehicles on the road and a traffic signal control unit. These components were installed on trunk roads

in the central Kanazawa city at the beginning of 2003. However, to activate FASTTM, emergency vehicles need to be equipped with an infrared beacon. In response to the increase in the length of FASTTM-implemented trunk roads, the fire department progressively replaced old non-FASTTM-equipped ambulances with newer FASTTM-equipped ambulances (Table 1). All ambulances in the central Kanazawa city had been loaded with the FASTTM equipment for the observation period. The new ambulances had lower horsepower to weight ratios than the old ambulances due to an increased demand to reduce fuel costs. In total, 48 traffic signals on trunk roads, at a total length of 12.6 km, were under the control of FASTTM in the central Kanazawa city and most major emergency hospitals were located in this area (Figure 1).

FASTTM does not modify traffic signals every time an ambulance passes; its function is controlled by integrated traffic control systems that are informed by current traffic conditions. Previous traffic engineering studies [17] revealed that FASTTM activated at a rate of 91.2% when ambulances passed FASTTM-controlled signals. Of the ambulances dispatched from the central area, 68.8% passed FASTTM-controlled signals.

Statistical analysis

The data for all OHCA cases unwitnessed by EMTs in the central area were compared between the FASTTM- and non-FASTTM-equipped ambulances, which individually comprised two groups. The control group was OHCA cases to which non-FASTTM-equipped ambulances were dispatched. We analyzed the effect of FASTTM installation on OHCA outcomes managed prior to hospital arrival without tracheal intubation or epinephrine administration because the incidences of these procedures

widely differed between the two groups (Table 2).

We analyzed the data using the Joint Medical Program (JMP), version 9, for Windows [Statistical Analysis System (SAS) Institute, Cary, NC, USA]. The chi-squared test with or without Pearson's correction was applied for univariate analyses. The Wilcoxon rank sums test and the Kruskal–Wallis test were used for non-parametric comparisons. One-way ANOVA was used for parametric comparison. We used multiple logistic regression analysis to elucidate the factors associated with the outcomes. In all analyses, $p < 0.05$ was considered statistically significant.

3. Results

Annual changes in FASTTM installation, traffic conditions, and critical parameters (Table 1)

The annual changes in OHCA outcomes, together with the various parameters related to them, were compared over the eight years of the observational study. Table 1 demonstrates the significant changes in the incidences of tracheal intubation, epinephrine administration, response time, and 1-Y survival rates for OHCA. An annual traffic survey at four intersections revealed that there were small, but significant changes in traffic volume (less than 10%), over the study period.

In accordance with the introductions of tracheal intubation and epinephrine administration, the EMT protocol was revised. However, during the study period, it was continuously emphasized that EMTs should provide high-quality basic life support (BLS) for all OHCA patients.

Background and time factor differences (Table 2)

There were significant differences in both arrest location (home versus others) and patient gender between the FASTTM- and non-FASTTM-equipped groups. However, no significant difference existed in the duration of transportation to hospital between FASTTM- and non-FASTTM-equipped ambulances. The response time was significantly shorter and the incidence of response time under 300 s was significantly higher in the FASTTM-equipped group. The median response times (25–75%) were 327 (244–429) s in the FASTTM-equipped group and 381 (291–487) s in the non-FASTTM-equipped group. ACLS procedures, including tracheal intubation and epinephrine administration, occurred more frequently in OHCA patients transported by FASTTM-equipped ambulances.

Comparisons of OHCA outcomes managed without prehospital ACLS procedures between FASTTM- and non-FASTTM-equipped ambulances (Figure 2)

Because tracheal intubation and epinephrine administration have been shown to affect OHCA outcomes [16, 18–22] and because the incidences of these procedures widely differed between the two groups (Table 2), we analyzed the impact of FASTTM on the outcomes of OHCA managed prior to hospital arrival without tracheal intubation or epinephrine administration.

As demonstrated in Figure 2, incidences of SROSC and 1-Y survival were significantly higher in the group with dispatch of FASTTM-equipped ambulances for all OHCA, OHCA with a presumed cardiac etiology, and witnessed OHCA with a presumed cardiac etiology.

Factors associated with 1-Y survival from OHCA managed without prehospital ACLS procedures

(Table 3)

As shown in Table 3, univariate analysis identified several factors associated with 1-Y survival: dispatch with FASTTM-equipped ambulance, patient age, location of cardiac arrest, witnessed cardiac arrest, arrests of a presumed cardiac etiology, and response time. Traffic volume, estimated from Table 1, was not a significant factor associated with 1-Y survival.

Multiple logistic analysis revealed that dispatch with a FASTTM-equipped ambulance, patient age, a witnessed cardiac arrest, and cardiac arrests with a presumed cardiac etiology were independent factors associated with 1-Y survival. The location of arrest or the response time was not an independent factor associate with 1-Y survival. However, adjusted odds ratio (OR) of response time for 1-Y survivals was 0.998 (95% confidence interval = 0.996–0.999), when the factor of dispatch with FASTTM-equipped ambulance was excluded from the logistic regression analysis,

4. Discussion

Because the incidences of tracheal intubation and epinephrine administration significantly increased during the study period and differed between the two groups, we compared the outcomes between the two groups for those OHCA managed without either tracheal intubation or epinephrine administration. This eight-year observational study in the central area of a single city showed that FASTTM implementation in ambulances successfully reduced the median response time by 54 s in EMT-unwitnessed OHCA and improved OHCA outcomes in the subgroup not receiving ACLS procedures. When assessing the OHCA subcategory of presumed cardiac etiology, the

FASTTM-equipped ambulance was significantly associated with greater incidences of an initial shockable rhythm, SROSC, 1-Y survival, and 1-Y survival with a favorable neurological outcome.

Multiple logistic regression analysis followed by univariate analysis revealed that FASTTM-equipped ambulance dispatch, patient age, cardiac etiology, and arrest witness were independent factors associated with 1-Y survival. Although univariate analysis disclosed a significant difference in response time between 1-Y survivors and non-survivors, multiple logistic regression analysis revealed that the response time was not an independent factor with 1-Y survival. However, the response time was an independent factor associated with 1-Y survival when dispatch with FASTTM-equipped ambulance was excluded from multiple logistic regression analysis. This difference was mostly due to the dependence of response time on the dispatch with FASTTM-equipped ambulances, which was shown as the significantly reduced response time and augmented incidence of early arrival (response time < 300 s (5 min)) by the dispatch with FASTTM-equipped ambulance in Table 2. Thus, the benefit of FASTTM implementation seems to be attributed, at least in part, to the reduced response time and/or incidence of arrival delay.

In contrast to response time, the transportation time widely varied and there was no significant difference in the transportation time between FASTTM- and non-FASTTM-equipped ambulances. The nearest ambulance team is always dispatched to the scene, while the transportation is not always made to the closest hospital. This may explain the difference between the effects of FASTTM-equipped ambulance on the median values of two time intervals.

The absolute difference in the median response time between the FASTTM- and

non-FASTTM-equipped ambulances was 54 s. This reduction may be too small to explain the improvement seen in outcomes. However, the adjusted unit OR of response time for 1-Y survival was 0.998 (95% confidence interval = 0.996–0.999) when the factor of dispatch with FASTTM-equipped ambulances was excluded from the logistic regression analysis, indicating that the reduction in response time by 1 s increases the 1-Y survival rate by 0.2%. It has been shown that the effect of reducing the response time on survival from OHCA is prominent when the response time does not exceed 5 to 6 min [5, 23]. Furthermore, a previous study showed that a short response time (less than 6 min) could lead to a high survival rate [24]. Reduced response time may be associated with an early application of first defibrillation that is related to the survival of OHCA patients having a shockable initial rhythm [25]. In this study, we showed that the incidence of response time less than 300 s (5 min) was significantly increased when transported by FASTTM-equipped ambulances (Table 2). Thus, a large improvement in the 1-Y survival seemed to be achievable by dispatch with FASTTM-equipped ambulances in the central area.

FASTTM has been implemented in Kanazawa city by the Police Department of Ishikawa Prefecture as a public enterprise. FASTTM has also been introduced in nine other cities and in the Tokyo metropolitan area in Japan. The exact cost of this initiative is unclear, but the Police Department of Ishikawa Prefecture has estimated that approximately 180 million yen (1.8 million USD) was spent to install the FASTTM communication system on trunk roads. In addition to the cost of installation on roads and in ambulances, the UTMS requires a traffic control center. The cost benefit of FASTTM implementation remains to be clarified.

Limitations

Immeasurable or unpredictable changes might have occurred during the study period, which might have affected our interpretation. Nevertheless, their impact might have been minimized by the gradual introduction of FAST™ equipment to ambulances between 2003 and 2011 (see Figure 1 and Table 1). All new ambulances equipped with a FAST™ beacon had a lower horsepower and weight ratio than the old ambulances due to an increased demand to reduce fuel costs. Therefore, it is unlikely that the newer FAST™-equipped ambulances benefitted from performance improvements. We also considered both the changes in traffic volume and the EMS level provided.

The EMS protocol changed several times during the study period, with major revisions made following the introduction of tracheal intubation and epinephrine administration. However, during the study period, it was continuously emphasized that EMTs should provide high-quality BLS for all OHCA patients. To remove the confounding effect of these changes, we also determined the significant factors associated with 1-Y survival in OHCA patients managed without ACLS procedures (Table 3). This exclusion appeared acceptable but bias may have been introduced according to literature suggesting unfavorable influences on OHCA outcomes [26-29].

The traffic volume estimated from Table 1 did not significantly differ between survivors and non-survivors. No data on the quality of bystander CPR were collected in this study. Bystander CPR is a recognized factor in achieving good outcomes for OHCA patients [30, 31], and the lack of record could negatively affect our results.

A previous traffic engineering study estimated that installing FAST™ equipment on

ambulances increased driving speeds by 17.9 km/h on FASTTM-implemented roads [17]. However, the actual distance of the FASTTM-implemented roads on which each ambulance drove was not known in this study. If this information were available for the entire observational period, a more in-depth analysis of this effect might be possible.

This study was a retrospective analysis of prospectively collected data from a single city in Japan. The data set analyzed might have been too small for accurate multiple logistic regression analysis. This limits the universality of this study, although the independent factors associated with survival in the present study were clearly comparable with those in previous, larger studies reported by us and others [32, 33].

5. Conclusions

This observational study in Kanazawa city showed that the FASTTM implementation significantly reduced ambulance response times and improved OHCA outcomes. However, a large, multi-region study is necessary to confirm the cost-benefit relationship for FASTTM implementation.

6. References

1. Vukmir RB. Survival from prehospital cardiac arrest is critically dependent upon response time. Resuscitation. 2006;69:229-234.
2. Pons PT, Haukoos JS, Bludworth W, et al. Paramedic response time: does it affect patient survival? Acad Emerg Med. 2005;12:594-600.

3. Pell JP, Sirel JM, Marsden AK, et al. Effect of reducing ambulance response times on deaths from out of hospital cardiac arrest: cohort study. *BMJ*. 2001;322:1385-8.
4. Do HQ, Nielsen SL, Rasmussen LS. Response interval is important for survival until admission after prehospital cardiac arrest. *Dan Med Bull*. 2010;57:A4203.
5. Sladjana A, Gordana P, Ana S. Emergency response time after out-of-hospital cardiac arrest. *Eur J Intern Med*. 201;22:386-93.
6. Neukamm J, Gräsner JT, Schewe JC, et al. The impact of response time reliability on CPR incidence and resuscitation success: a benchmark study from the German Resuscitation Registry. *Crit Care*. 2011;15:282.
7. Kitamura T, Iwami T, Kawamura T, et al. Nationwide improvements in survival from out-of-hospital cardiac arrest in Japan. *Circulation*. 2012;126:2834-43.
8. Nichol G, Laupacis A, Stiell IG, et al. Cost-effectiveness analysis of potential improvements to emergency medical services for victims of out-of-hospital cardiac arrest. *Ann Intern Med*. 1996;27:711-20.
9. Lin CS, Chang H, Shyu KG, et al. A method to reduce response times in prehospital care: the motorcycle experience. *Am J Emerg Med*. 1998;16:711-3.
10. UTMS society of Japan.
(Accessed 5 March 2013, at <http://www.utms.or.jp/english/system/index.html>)
11. FAST (Fast Emergency Vehicle Preemption Systems).
(Accessed 5 March 2013, at <http://www.utms.or.jp/english/system/fast.html>)

12. The Ministry of Health, Labour and Welfare in Japan. (Accessed 5 March 2013, at

<http://www.mhlw.go.jp/seisakunitsuite/bunya/hokabunya/kenkyujigyou/i-kenkyu/index.html>)

13. Cummins RO, Chamberlain DA, Abramson NS, et al. Recommended guidelines for uniform reporting of data from out-of-hospital cardiac arrest: the Utstein Style. A statement for health professionals from a task force of the American Heart Association, the European Resuscitation Council, the Heart and Stroke Foundation of Canada, and the Australian Resuscitation Council. *Circulation*. 1991;84:960–75.

14. Jacobs I, Nadkarni V, Bahr J, et al. Cardiac arrest and cardiopulmonary resuscitation outcome reports: update and simplification of the Utstein templates for resuscitation registries: a statement for healthcare professionals from a task force of the International Liaison Committee on Resuscitation (American Heart Association, European Resuscitation Council, Australian Resuscitation Council, New Zealand Resuscitation Council, Heart and Stroke Foundation of Canada, Inter American Heart Foundation, Resuscitation Councils of Southern Africa). *Circulation*. 2004;23:3385–97.

15. A. Safar P. Resuscitation after Brain Ischemia, in Grenvik A and Safar P Eds: *Brain Failure and Resuscitation*. Churchill Livingstone, New York 1981;155-84.

16. Takei Y, Enami M, Yachida T, et al. Tracheal intubation by paramedics under limited intubation criteria may improve the short-term outcome of out-of-hospital cardiac arrests with noncardiac origin. *J Anesth*. 2010;24:716-25.

17. Takada K, Inaba H, Nanbu S. Effect of introducing a FAST Emergency Vehicle Preemption system in Kanazawa (in Japanese). *IATSS Review*. 2009;34:47-54.

18. Stiell IG, Wells GA, Field B, et al. Advanced cardiac life support in out-of-hospital cardiac arrest. *N Engl J Med*. 2004;351:647-56.
19. Olasveengen TM, Sunde K, Brunborg C, et al. Intravenous drug administration during out-of-hospital cardiac arrest: a randomized trial. *JAMA*. 2009;302:2222-9.
20. Studnek JR, Thestrup L, Vandevender S, et al. The association between prehospital endotracheal intubation attempts and survival to hospital discharge among out-of-hospital cardiac arrest patient. *Acad Emerg Med*. 2010;17:918-25.
21. Jacobs IG, Finn JC, Jelinek GA, et al. Effect of adrenalin on survival in out-of-hospital cardiac arrest: A randomised double-blind placebo-controlled trial. *Resuscitation*. 2011;82:1138-43.
22. Nakahara S, Tomio J, Nishida M, et al. Association between timing of epinephrine administration and intact neurologic survival following out-of-hospital cardiac arrest in Japan: a population-based prospective observational study. *Acad Emerg Med*. 2012;19:782-92.
23. 2005 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Part 4: Adult Basic Life Support. *Circulation*. 2005;112:IV-19-IV-34.
24. O'Keeffe C, Nicholl J, Turner J, et al. Role of ambulance response times in the survival of patients with out-of-hospital cardiac arrest. *Emerg Med J*. 2011;28:703-6.
25. De Maio VJ, Stiell IG, Wells GA, et al. Optimal defibrillation response intervals for maximum out-of-hospital cardiac arrest survival rates. *Ann Emerg Med*. 2003;42:242-50.
26. Advanced airway management does not improve outcome of out-of-hospital cardiac arrest. Hanif MA, Kaji AH, Niemann JT. *Acad Emerg Med*. 2010;17:926-31.

27. Hasegawa K, Hiraide A, Chang Y, Brown DF. Association of prehospital advanced airway management with neurologic outcome and survival in patients with out-of-hospital cardiac arrest. JAMA. 2013;309:257-66.
28. Holmberg M, Holmberg S, Herlitz J. Low chance of survival among patients requiring adrenaline (epinephrine) or intubation after out-of-hospital cardiac arrest in Sweden. Resuscitation. 2002;54:37-45.
29. Hagihara A, Hasegawa M, Abe T, et al. Prehospital epinephrine use and survival among patients with out-of-hospital cardiac arrest. JAMA. 2012;307:1161-8.
30. Gallagher EJ, Lombardi G, Gennis P. Effectiveness of bystander cardiopulmonary resuscitation and survival following out-of-hospital cardiac arrest. JAMA. 1995;274:1922-1925.
31. Wik L, Steen PA, Bircher NG. Quality of bystander cardiopulmonary resuscitation influences outcome after prehospital cardiac arrest. Resuscitation. 1994;28:195-203.
32. Takei Y, Inaba H, Yachida T, et al. Analysis of reasons for emergency call delays in Japan in relation to location: High incidence of correctable causes and the impact of delays on patient outcomes. Resuscitation. 2010;81:1492-8.
33. Holmberg M, Holmberg S, Herlitz J, et al. Survival after cardiac arrest outside hospital in Sweden. Resuscitation. 1998;36:29-36.

7. Legends for Figures/Tables

Figure legends

Figure 1 Trunk roads with infrared beacons installed, and the location of ambulance teams and major emergency hospitals in Kanazawa city.

<u>Ambulance Team Number</u>	<u>Name of Team</u>	<u>Dispatch Area</u>	<u>Date of FASTTM Installation</u>
AT1	Ekinishi	Central	July 28, 2003
AT2	Chuo	Central	December 21, 2004
AT3	Misogura	Central	March 16, 2007
AT4	Izumino	Central	January 27, 2011

AT-(number): ambulance team number.

H1: Ishikawa Prefecture Central Hospital. H2: Kanazawa Medical Center. H3: Kanazawa University Hospital

Figure 2 Crucial comparisons of outcomes between FASTTM- and non-FASTTM-equipped ambulances for OHCAs managed without ACLS procedures.

The panels represent all OHCAs, OHCAs with a presumed cardiac etiology, and witnessed OHCAs with a presumed cardiac etiology.

★ Significant difference between the dispatch with FASTTM- and non-FASTTM-equipped ambulances (by chi-squared test with Person's correction).

Table legends

Table 1 Annual changes in FASTTM installation, traffic conditions, and critical parameters.

* At the beginning of fiscal year.

Table 2 Differences in backgrounds, time factors, and the management of OHCAs between the dispatch with FASTTM- and non-FASTTM-equipped ambulances.

* CPR first performed by either citizen or EMT.

Table 3 Factors associated with 1-Y survival of all OHCAs managed without ACLS procedures.

* Multiple logistic regression analysis.

** CPR first performed by either citizen or EMT.

CI: confidence interval.

Table 1 Annual changes in FAST™ installation, traffic conditions, and critical parameters

Fiscal year	Total length of FAST™ - implemented roads (km)	Number of ambulances equipped with FAST™ beacon unit*	Traffic volume, median (cars/day)	Number of OHCA's (Dispatch with FAST™-equipped ambulances/total), N (%)	Prehospital ACLS		Response time, second (25–75%)	1-Y survival rate, N (%)
					Tracheal intubation, N (%)	Epinephrine administration, N (%)		
2003	2.3	0	1440	22/138 (15.9%)	0	0	360 (240–480)	1 (0.7%)
2004	6.0	1	1368	45/128 (35.2%)	5 (3.9%)	0	300 (240–420)	6 (4.7%)
2005	8.9	2	1444	51/127 (40.2%)	18 (14.2%)	0	360 (240–480)	4 (3.1%)
2006	12.6	2	1431	100/130 (76.9%)	15 (11.5%)	6 (4.6%)	360 (240–420)	7 (5.4%)
2007	12.6	2	1393	104/141 (73.8%)	12 (8.5%)	8 (5.7%)	293 (234–400)	9 (6.4%)
2008	12.6	3	1379	105/135 (77.8%)	35 (25.9%)	6 (4.4%)	340 (269–428)	10 (7.4%)
2009	12.6	3	1376	131/192 (68.2%)	31 (16.1%)	23 (12.0%)	386 (280–500)	5 (2.6%)
2010	12.6	3	1395	129/170 (75.9%)	19 (11.2%)	34 (21.8%)	377 (301–485)	14 (8.2%)
Statics	Undefined	Undefined	One-way ANOVA	Chi-squared analysis with Pearson's correction	Chi-squared analysis with Pearson's correction		Kruskal–Wallis test	Chi-squared analysis with Pearson's correction
<i>p</i>			<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	0.0360

* At the beginning of fiscal year.

Table 2 Differences in backgrounds, time factors, and the management of OHCA between the dispatch with FASTTM- and non-FASTTM-equipped ambulances

	Dispatch with FAST TM -equipped ambulances	Dispatch with non- FAST TM -equipped ambulances	Wilcoxon or chi-squared test
Number	687	474	
Etiology: cardiac, N (%)	317 (46.1%)	235 (49.6%)	0.2493
Arrests: witnessed, N (%)	320 (46.6%)	219 (46.2%)	0.8993
Location: home, N (%)	449 (65.4%)	276 (58.2%)	0.0137
CPR before arrival, N (%)	371 (54.0%)	250 (52.7%)	0.6722
Age, median (25–75%)	75 (60–83)	76 (61–84)	0.3591
Sex: female, N (%)	256 (37.3%)	210 (44.3%)	0.0162
Collapse/arrest recognition–call (min), median (10–25–75–90%)	2 (0–1–4–10)	2 (0–0–4–10)	0.0534
Response time (s), median (10–25–75–90%)	327 (197–244–429–571)	381 (238–291–487–704)	<0.0001
Response time <300 s, N (%)	262/687 (38.1%)	123/474 (25.9%)	<0.0001
Call-first CPR* (min), median (10–25–75–90%)	3 (-1–0–7–9)	4 (-2–0–7–10)	0.3328
Duration of transportation to hospitals (s), median (10–25–75–90%)	420 (224–300–558–718)	420 (240–300–600–810)	0.0768
Tracheal intubation, N (%)	102 (14.8 %)	33 (7.0%)	<0.0001
Epinephrin administration, N (%)	64 (9.3%)	16 (3.4 %)	<0.0001

* CPR first performed by either citizen or EMT.

Table 3 Factors associated with 1-Y survival of all OHCA managed without ACLS procedures

Factors	1-Y survival		<i>p</i>	Odds ratio (95% CI)	Adjusted odds ratio* (95% CI)
	Yes	No			
	N = 49	N = 909			
Dispatch with FAST™-equipped ambulance, N (%)	37 (75.5%)	493 (54.2%)	0.0035	2.602 (1.339–5.054)	3.077 (1.180–9.350)
Ambulance team, N			0.1848		
Ekinishi EMS	9	194		Undefined	Reference
Chuo EMS	15	215		Undefined	0.814 (0.308–2.051)
Misogura EMS	17	240		Undefined	0.461 (0.175–1.143)
Izumino EMS	8	260		Undefined	0.394 (0.091–1.604)
Patient age (years), median (25–75%)	63 (48–73)	74 (60–83)	0.0003	Undefined	0.976 (0.962–0.990)
Patient sex: female, N (%)	15 (30.6%)	372 (40.9%)	0.1519	1.579 (0.843–2.924)	0.933 (0.467–1.803)
Location: home, N (%)	23 (46.9%)	569 (62.6%)	0.0280	0.529 (0.297–0.941)	0.633 (0.337–1.186)
Arrest: witnessed, N (%)	42 (85.7%)	400 (44.0%)	<0.0001	7.635 (3.394–17.176)	6.798 (3.125–17.061)
Etiology: cardiac, N (%)	33 (67.4%)	442 (48.6%)	0.0107	2.179 (1.183–4.015)	2.593 (1.366–5.130)
CPR before EMT arrival, N (%)	29 (59.2%)	461 (50.7%)	0.2480	1.409 (0.786–2.528)	1.603 (0.754–3.119)
Collapse/arrest recognition–call (min), median (25–75%)	2 (1–3)	2 (1–4)	0.8312	Undefined	0.983 (0.943–1.013)
Response time (s), median (25–75%)	284 (240–343)	357 (253–472)	0.0010	Undefined	0.998 (0.996–1.000)
Call–first CPR** (min), median (25–75%)	1 (0–6)	4 (0–7)	0.0877	Undefined	0.999 (0.940–1.005)
Transportation to hospitals (s), median (25–75%)	397 (300–486)	420 (300–552)	0.2679	Undefined	0.999 (0.999–1.001)
Traffic volume estimated from Table 1, cars/day, (25–75%)	1376 (1341–1395)	1376 (1341–1395)	0.1281	Undefined	1.007 (0.996–1.019)

* Multiple logistic regression analysis. ** CPR first performed by either citizen or EMT. CI: confidence interval.