Repeated total en bloc spondylectomy for spinal metastases at different sites in one patient

著者	Sugita Shurei, Murakami Hideki, Demura Satoru, Kato Satoshi, Yoshioka Katsuhito, Yokogawa Noriaki, Tanaka Sakae, Tsuchiya Hiroyuki
journal or publication title	European Spine Journal
volume	24
number	10
page range	2196-2200
year	2015-10-01
URL	http://hdl.handle.net/2297/43036

doi: 10.1007/s00586-015-4091-y

Title: Repeated total *en bloc* spondylectomy for spinal metastases at different sites in one patient.

Shurei Sugita, M.D., Ph.D.,¹ Hideki Murakami M.D., Ph.D.,¹ Satoru Demura, M.D., Ph.D.,¹

Satoshi Kato M.D., Ph.D.,¹ Katsuhito Yoshioka M.D., Ph.D.,¹ Noriaki Yokogawa M.D.,¹

Sakae Tanaka M.D., Ph.D.,² and Hiroyuki Tsuchiya M.D., Ph.D.¹

1) Department of Orthopaedic Surgery, Kanazawa University, Kanazawa, Japan

2) Department of Orthopaedic Surgery, Faculty of Medicine, the University of Tokyo,
Tokyo, Japan

Correspondence should be addressed to:

Shurei Sugita, M.D., Ph.D.

13-1, Takaramachi, Kanazawa, Ishikawa, 920-8641, Japan

Tel: +81762652374 Fax: +81762344261

email: ssugita-tky@umin.ac.jp

Conflict of interest; none

Title: Repeated total *en bloc* spondylectomy for spinal metastases at different sites in one patient.

Keywords:

spinal metastases; total en bloc spondylectomy; preoperative artery embolization; nerve root ligation and cut; repeated times

Abstract

Purpose: Total *en bloc* spondylectomy (TES) is accompanied by preoperative embolization of segmental arteries, which is limited to three consecutive levels to avoid the risk of spinal cord ischemia. We retrospectively examined the efficacy and safety of repeated TES with embolization of more than three levels of segmental arteries.

Methods: Seven patients underwent TES twice for spinal metastases at different levels.

Every patient underwent embolization of the bilateral segmental arteries before each

surgery. We assessed the total number of segmental arteries embolized, the existence of

Adamkiewicz arteries during the embolization procedure, intraoperative blood loss and the

motor function of the lower limbs, using the American Spinal Injury Association (ASIA)

motor score.

Results: No patient experienced any motor deficit after embolization. During the

1
2
3
4
5
6
7
8 embolization procedure, an Adamkiewicz artery was depicted in five patients, which
9
10 precluded embolization at that level. The median number of segmental arteries embolized
11
12 in total was 9 (9-11). Intraoperative blood loss (median, IQR) was 480 (420 – 630) g during
13
14 the first surgery and 520 (280 – 600) g during the second surgery. The ASIA motor scores
15
16 (median, IQR) were as follows; 100 (98 – 100) (first admission), 100 (100 – 100) (first
17
18 discharge), 100 (98 – 100) (second admission), and 97 (94 – 100) (second discharge). No
19
20 patients had developed statistically significant neurological deterioration, and there had
21
22 been no local recurrence after a median follow-up of 17.8 months (range 1 - 51 months).
23
24
25
26
27
28
29
30
31
32
33 Conclusion: Repeated TES procedures can be performed safely even if more than three
34
35
36 levels of segmental arteries are embolized.
37
38
39
40
41

42 Introduction

43
44
45 Spinal metastases is a clinically challenging condition, with a morbidity rate of
46
47
48 approximately 30% among cancer patients [1]. Additionally, 5% of cancer patients will
49
50
51 develop neurological deterioration during their clinical course [2-5]. The resulting clinical
52
53
54 burden of these conditions is substantial due to the increasing life expectancy
55
56
57
58 accompanying the recent advances in oncological therapies.
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8 The purposes of treatment for spinal metastases are to prevent tumor recurrence,
9
10 to decompress the spinal cord to avoid paralysis, and to restore the mechanical stability of
11
12 the spine. To achieve these purposes, we developed a new procedure, total *en bloc*
13
14 spondylectomy, which aims to provide a the local cure for the spinal metastases [6]. This
15
16 technique of *en bloc* resection has proved to be effective in improving the prognosis of both
17
18 primary and metastatic bone tumors of the spine [7, 8]. To prevent massive intraoperative
19
20 bleeding, we routinely embolize the segmental artery at the level of the affected vertebrae
21
22 one or two days before surgery [9]. However, the extent of embolization is limited up to
23
24 three levels to help prevent spinal cord ischemia [10, 11].
25
26
27
28
29
30
31
32
33
34
35

36 In some cases, after the first TES, another spinal metastasis emerges at a different
37
38 level, and it is sometimes necessary to perform another TES procedure. As part of this
39
40 procedure, we have also had to embolize the segmental arteries, corresponding to the
41
42 newly-emerged spinal metastases. As a result, the total number of embolized arteries
43
44 exceeded three levels, which could potentially induce spinal cord ischemia, leading to
45
46 neurologic compromise
47
48
49
50
51
52
53
54

55 The purpose of this study was to retrospectively examine the efficacy and safety
56
57 of repeated TES, with embolization of more than three levels of segmental arteries.
58
59
60
61
62
63
64
65

Materials and methods

Patients

We retrospectively reviewed the records of seven patients (four males and three females, median age: 51 years old, (IQR 42 – 57) Table 1) who underwent TES twice for spinal metastases at different levels and times within study period between April 2010 and December 2014. The cases consisted of four patients with renal cell carcinoma, two with thyroid carcinoma and one with chondrosarcoma.

We collected the data regarding arterial embolization (the total number of segmental arteries embolized, the existence of an Adamkiewicz artery depicted during the embolization procedure), the surgical procedure (intraoperative blood loss, months between the first and second operation) and the clinical outcome (motor function of the lower limbs, using the American Spinal Injury Association (ASIA) motor score, local recurrence during the study period)

Embolization procedure

Before each TES procedure was accompanied by preoperative embolization of segmental arteries. The patients underwent diagnostic angiography and embolization under local anesthesia using a femoral approach. Selective catheterization of the corresponding

1
2
3
4
5
6
7
8 segmental arteries, including one level above and one below the tumor site, was performed
9
10 followed by selective angiography of suspected tumor feeder arteries to the tumor. Coils
11
12
13
14 were used in combination with polyvinyl alcohol as the embolic material. If an
15
16
17 Adamkiewicz artery was depicted during angiography procedure embolization at the origin
18
19
20 of this segmental artery was avoided.
21
22

23 Surgical procedure

24
25
26 All patients underwent the routine surgical procedure as has been described in the literature
27
28
29 [6]. Decompression of the spinal cord was accomplished by *en bloc* corpectomy following
30
31
32
33 *en bloc* laminectomy. The anterior column was then reconstructed by inserting a vertebral
34
35
36 spacer packed with autograft bone. During each surgery, posterior fixation extended two
37
38
39 levels above and two below the tumor site. If the proposed levels of fixation in the second
40
41
42 surgery were to overlap the fixation placed during the first surgery, rod connectors or
43
44
45 converting rods were used.
46
47

48 Statistical analysis

49
50
51
52 Statistical analyses were performed using the JMP® version 11 software program (SAS,
53
54
55 Cary, NC) Descriptive statistics were used to analyze clinical information, demographic
56
57
58 factors and other test data. Continuous variables are expressed as the medians and
59
60
61
62
63
64
65

interquartile range (IQR). The differences in the motor scores before and after treatment were examined using Mann-Whitney *U* test. A *P* value < 0.05 was considered to be statistically significant.

Results

Embolization

Every patient underwent preoperative embolization of the segmental arteries before each surgery. The mean total number of segmental arteries embolized was nine (range 8-11). Detailed data about embolization procedure are shown in Table 2. During the embolization procedure, an Adamkiewicz artery was depicted in five patients (Table 2), and embolization at the origin of this segmental artery was avoided in these cases. No patient experienced deterioration of his or her baseline neurologic status post-embolization.

Surgery

Detailed data about the surgical procedure are shown in Table 3. The median interval between the first and second surgeries (median, IQR) was 21 (5 – 49.1) months. In the thoracic spine, the nerve roots corresponding to the tumor level were ligated and cut; as necessary, nerve roots one level above and/or below were also ligated and cut. In the lumbar spine, the nerve roots below the L3 level were not cut, but only dissected from

1
2
3
4
5
6
7
8 surrounding tissues. If the corresponding nerve roots were at L1 or L2, we also ligated and
9
10 cut these roots. Based on our experience, sacrificing the nerve roots above the L1 or L2
11
12
13 level does not lead to a significant loss of neurologic function.
14
15

16
17 The intraoperative blood loss (median, IQR) was 480 (420 – 630) g during the
18
19 first surgery, and 520 (280 – 600) g during the second surgery. Only one patient needed a
20
21 blood transfusion postoperatively.
22
23

24 Outcomes

25
26 The ASIA motor scores of the lower limbs are shown in Fig. 1. No patients had developed
27
28 statistically significant neurological deterioration and all were free from local recurrence
29
30 after a median follow-up of 17.8 months (range 1- 51 months). Significant surgical
31
32 complications were two hardware failures which required revision surgery, and a case of
33
34 postural hypotension of unknown etiology.
35
36
37
38
39
40
41
42
43
44

45 Illustrative case

46
47 Case.7 was a 68-year-old male with spinal metastases from renal cell carcinoma to T7 who
48
49 presented at our hospital. After embolization of the bilateral segmental arteries from T6 to
50
51 T8, he underwent TES of T7 and experienced no perioperative complications (Fig. 2a).
52
53
54
55
56
57
58 However, 20 months later, rod breakage was found on follow-up radiographs and newly
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8 emerged metastases at T8 and T9 were discovered on MRI (Fig. 2b). A second TES
9
10 procedure was planned and during preoperative embolization, a communicating branch had
11
12 developed from the T9 segmental artery to the T8 segmental artery (Fig. 2c). The patient
13
14 underwent a second TES procedure at T8 and T9 without complications (Fig. 2d). Since
15
16 surgery, no local recurrence has been observed during the 17-month follow-up period.
17
18
19
20
21
22

23 Discussion

24
25
26 In our study, repeated TES and preoperative embolization caused no neurological
27
28 deterioration arising from spinal cord ischemia or the surgical procedure. The local control
29
30 of the affected spine was favorable.
31
32
33
34
35

36 Preoperative embolization to prevent excessive intraoperative bleeding when
37
38 surgically treating metastatic spinal tumors is now widely used [12]. At our institution, the
39
40 extent of embolization has been limited up to three levels to prevent spinal cord ischemia
41
42 [10, 11]. In this case series, though embolization was performed twice at separate times, the
43
44 total number of embolized arteries exceeded three levels, which may increase the risk of
45
46 neurological deterioration due to decreased blood flow to the spinal cord. However the
47
48 outcomes with regard to the neurological function did not worsen even after the second
49
50 TES. There have been several studies which have indicated that occlusion of the segmental
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8 arteries leads to the development of collateral pathways of intraspinal and paraspinal
9
10 arteries and arterioles [13-16], and these collateral pathways developed within five days
11
12 after occlusion in those cases. These studies may explain the reasons for the preservation of
13
14 the neurological function after the multiple-level embolization of the segmental arteries in
15
16 our case series. The time interval between the first and second embolizations was long
17
18 enough to allow the development of a collateral artery network. An example of this is seen
19
20 in case 7, where collateral blood flow from the T9 segmental artery to the T8 segmental
21
22 artery was depicted during the second embolization procedure (Fig. 2).
23
24
25
26
27
28
29
30
31
32

33 Regarding the surgical procedure, nerve roots at the operative level had to be
34
35 sacrificed if the level was above L3. In this case series, the number of sacrificed nerve roots
36
37 was multiple and often exceeded three levels. Despite this, neurological function remained
38
39 preserved. We previously reported that ligation of nerve roots in the thoracic region did not
40
41 lead to worsening of the neurological function [17]. We have also observed that neurologic
42
43 function is also preserved following the ligation and transection of the L1 or L2 nerve roots.
44
45
46
47
48
49
50
51 Further studies will be needed to confirm these studies, but they support our experiences
52
53
54
55 indicating that repeat TES is safe.
56
57

58 The TES procedures in this case series were without any perioperative
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8 complications. The intraoperative blood loss was minimized, and only one of the seven
9
10 patients required a blood transfusion, which indicates that the embolization had succeeded
11
12 in limiting the intraoperative blood loss, as reported previously[9, 12, 18, 19].
13
14

15
16
17 There are several limitations associated with this study that should be kept in mind when
18
19 interpreting the results. First, the number of cases in this study was relatively small (seven)
20
21 and we are unable to extrapolate the result of this study to future cases. Secondly, we had
22
23 two cases of hardware failure in these seven cases, both of which required revision surgery.
24
25
26 Furthermore, improvements in reconstruction after the resection of tumor-containing
27
28 vertebra are needed.
29
30
31
32
33
34

35 36 Conclusion

37
38
39 Repeated TES procedures can be performed safely and can provide good surgical outcomes,
40
41 even if preceded by embolization of segmental arteries in more than three levels.
42
43
44
45
46
47

48 49 References

- 50
51
52
53
54 1. Ortiz Gomez JA (1995) The incidence of vertebral body metastases. International
55 orthopaedics 19:309-311
56
57 2. Black P (1979) Spinal metastasis: current status and recommended guidelines for
58
59
60
61
62
63
64
65

management. *Neurosurgery* 5:726-746

3. Rades D, Blach M, Nerreter V, Bremer M, Karstens JH (1999) Metastatic spinal cord compression. Influence of time between onset of motoric deficits and start of irradiation on therapeutic effect. *Strahlentherapie und Onkologie : Organ der Deutschen Rontgengesellschaft [et al]* 175:378-381

4. Constans JP, de Divitiis E, Donzelli R, Spaziante R, Meder JF, Haye C (1983) Spinal metastases with neurological manifestations. Review of 600 cases. *J Neurosurg* 59:111-118. doi: 10.3171/jns.1983.59.1.0111

5. Bach F, Larsen BH, Rohde K, Borgesen SE, Gjerris F, Boge-Rasmussen T, Agerlin N, Rasmussen B, Stjernholm P, Sorensen PS (1990) Metastatic spinal cord compression. Occurrence, symptoms, clinical presentations and prognosis in 398 patients with spinal cord compression. *Acta neurochirurgica* 107:37-43

6. Tomita K, Kawahara N, Baba H, Tsuchiya H, Nagata S, Toribatake Y (1994) Total en bloc spondylectomy for solitary spinal metastases. *Int Orthop* 18:291-298

7. Boriani S, Bandiera S, Donthineni R, Amendola L, Cappuccio M, De Iure F, Gasbarrini A (2010) Morbidity of en bloc resections in the spine. *Eur Spine J* 19:231-241. doi: 10.1007/s00586-009-1137-z

8. Kato S, Murakami H, Demura S, Yoshioka K, Kawahara N, Tomita K, Tsuchiya H (2014) More than 10-year follow-up after total en bloc spondylectomy for spinal tumors. *Annals of surgical oncology* 21:1330-1336. doi: 10.1245/s10434-013-3333-7

9. Guzman R, Dubach-Schwizer S, Heini P, Lovblad KO, Kalbermatten D, Schroth G, Remonda L (2005) Preoperative transarterial embolization of vertebral metastases. *Eur Spine J* 14:263-268. doi: 10.1007/s00586-004-0757-6

10. Fujimaki Y, Kawahara N, Tomita K, Murakami H, Ueda Y (2006) How many ligations of bilateral segmental arteries cause ischemic spinal cord dysfunction? An experimental study using a dog model. *Spine (Phila Pa 1976)* 31:E781-789. doi: 10.1097/01.brs.0000238717.51102.79

11. Kato S, Kawahara N, Tomita K, Murakami H, Demura S, Fujimaki Y (2008) Effects on spinal cord blood flow and neurologic function secondary to interruption of bilateral segmental arteries which supply the artery of Adamkiewicz: an experimental study using a dog model. *Spine (Phila Pa 1976)* 33:1533-1541. doi: 10.1097/BRS.0b013e318178e5af

12. Pikis S, Itshayek E, Barzilay Y, Hasharoni A, Kaplan L, Gomori M, Cohen JE (2014) Preoperative embolization of hypervascular spinal tumors: current practice and center experience. *Neurological research* 36:502-509. doi: 10.1179/1743132814y.0000000361

13. Etz DC, Luehr M, Aspern KV, Misfeld M, Gudehus S, Ender J, Koelbel T, Debus ES, Mohr FW (2014) Spinal cord ischemia in open and endovascular thoracoabdominal aortic aneurysm repair: new concepts. *The Journal of cardiovascular surgery* 55:159-168
14. Luehr M, Salameh A, Haunschild J, Hoyer A, Grrbach FF, von Aspern K, Dhein S, Mohr FW, Etz CD (2014) Minimally invasive segmental artery coil embolization for preconditioning of the spinal cord collateral network before one-stage descending and thoracoabdominal aneurysm repair. *Innovations (Philadelphia, Pa)* 9:60-65. doi: 10.1097/imi.0000000000000038
15. Etz CD, Kari FA, Mueller CS, Brenner RM, Lin HM, Griep RB (2011) The collateral network concept: remodeling of the arterial collateral network after experimental segmental artery sacrifice. *The Journal of thoracic and cardiovascular surgery* 141:1029-1036. doi: 10.1016/j.jtcvs.2010.06.017
16. Geisbusch S, Schray D, Bischoff MS, Lin HM, Griep RB, Di Luozzo G (2012) Imaging of vascular remodeling after simulated thoracoabdominal aneurysm repair. *The Journal of thoracic and cardiovascular surgery* 144:1471-1478. doi: 10.1016/j.jtcvs.2012.08.062
17. Murakami H, Kawahara N, Demura S, Kato S, Yoshioka K, Tomita K (2010) Neurological function after total en bloc spondylectomy for thoracic spinal tumors. *Journal of neurosurgery Spine* 12:253-256. doi: 10.3171/2009.9.spine09506
18. Kato S, Murakami H, Minami T, Demura S, Yoshioka K, Matsui O, Tsuchiya H (2012) Preoperative embolization significantly decreases intraoperative blood loss during palliative surgery for spinal metastasis. *Orthopedics* 35:e1389-1395. doi: 10.3928/01477447-20120822-27
19. Kato S, Hozumi T, Takaki Y, Yamakawa K, Goto T, Kondo T (2013) Optimal schedule of preoperative embolization for spinal metastasis surgery. *Spine (Phila Pa 1976)* 38:1964-1969. doi: 10.1097/BRS.0b013e3182a46576

Figure legends.

Figure 1.

The ASIA motor scores in this case series.

Figure 2.

(a) A postoperative radiograph of the thoracic spine after first surgery.

(b) MRI images showing recurring metastasis at T8,9 (a high intensity area around the pedicle screw in the axial image)

(c) The angiography findings before the second operation. The right T8 segmental artery with a collateral pathway to T9 was depicted during angiography of the T9 artery (arrow).

(d) A postoperative radiograph of the thoracic spine after the second surgery.

Fig.1

ASIA
Motor
Score

100

75

50

Before
1st
surgery

After
1st
surgery

Before
2nd
surgery

After
2nd
surgery

- ◆— Case1
- Case2
- ▲— Case3
- Case4
- Case5
- Case6
- +— Case7

Fig.2

Table 1. The demographic data of the seven patients included in this study

Case		Age	Gender	Tumor
	1	48	M	RCC
	2	39	M	Thyroid CA.
	3	57	F	Thyroid CA.
	4	42	M	RCC
	5	51	F	RCC
	6	54	F	Chondrosarcoma
	7	68	F	RCC

RCC: Renal Cell Carcinoma, Thyroid CA: Thyroid Carcinoma

Table 2. Detailed data about the embolization procedures

Case	Embolized segmental arteries (1st)	Embolized segmental arteries (2nd)	Total number of embolized arteries	Origin of Adamkiewicz artery
1	Lt T3, bil T4, 5	Bil T11,12, L1	11	Rt T3
2	Lt T3, bil T4, Lt T5	Bil L3, 4	8	Not visualized
3	Bil T4, 5, 6	Bil L1, rt L3, L4	10	Not visualized
4	Bil T7, 8, rt T9	Bil L3, 4	9	Lt T9
5	Bil T9, 10, 11	Bil L1, 3, rt L2	11	Lt L2
6	Bil T7, T8, lt T9	Bil T12, L2	9	Rt T9
7	Bil T6, 7, 8	Bil T9, rt T10	9	Lt T10

Table 3. Detailed data about surgical procedure.

Case	First operation			Second operation			Months between surgeries
	Excised vertebrae	Sacrificed roots	Blood loss (g)	Excised vertebra	Sacrificed roots	Blood loss (g)	
1	T4	Bil T4	480	T12, L1	Bil T12, L1	420	12.9
2	T4	Bil T3, T4	860	L4	none	950	1.3
3	T5, 6	Bil T5, T6	630	L1	Bil L1	280	30.0
4	T7, 8	Bil T7, T8	480	L4	none	600	5.0
5	T9, 10, 11	Bil T9, T10, T11	340	L2, 3	Bil L2	570	54.2
6	T7, 8, 9	Bil T7, T8, T9	420	L1	Bil L1	180	49.1
7	T7	Bil T7	550	T8, 9	Bil T8, T9	520	21.0