Polybrominated diphenyl ethers in human serum and sperm quality

著者	Akutsu Kazuhiko, Takatori S., Nozawa S., Yoshiike M., Nakazawa H., Hayakawa Kazuichi, Makino T., Iwamoto T.
journal or publication title	Bulletin of Environmental Contamination and Toxicology
volume	80
number	4
page range	345-350
year	2008-04-01
URL	http://hdl.handle.net/2297/9975

doi: 10.1007/s00128-008-9370-4

Polybrominated Diphenyl Ethers in Human Serum and Sperm Quality

K. Akutsu,^{1,2*} S. Takatori,¹ S. Nozawa,³ M. Yoshiike,³ H. Nakazawa,⁴ K. Hayakawa,² T. Makino,⁵ T. Iwamoto^{3†}

¹ Division of Food Chemistry, Osaka Prefectural Institute of Public Health, 1-3-69 Nakamichi, Higashinari-ku, Osaka 537-0025, Japan

² Graduate School of Natural Science and Technology, Kanazawa University, Kakumamachi, Kanazawa, Ishikawa 920-1192, Japan

³ Department of Urology, St. Marianna University School of Medicine, 2-16-1 Sugao, Miyamae, Kawasaki, Kanagawa 216-8511, Japan

⁴ Department of Analytical Chemistry, Hoshi University, 2-4-41 Ebara, Shinagawa-ku, Tokyo 142-8501, Japan

⁵ Department of Obstetrics and Gynecology, School of Medicine, Tokai University, 143 Shimokasuya, Isehara, Kanagawa 259-1143, Japan

[†]Present address: Center for Infertility and IVF, International University of Health and Welfare, 2600-1 Kitakanemaru, Ohtawara, Tochigi 324-8501, Japan.

Abstract: Polybrominated diphenyl ethers (PBDEs) are widely used flame retardants; currently, they are identified as ubiquitous environmental contaminants. Several studies indicate that PBDEs might affect male fertility. We present the results of a pilot study on the relationship between human serum PBDEs and sperm quality. The PBDE levels in Japan are comparable to those found in European countries. Strong inverse correlations were observed between the serum concentration of 2,2',4,4',5,5'-hexabromodiphenyl ether and sperm concentration ($r = -0.841$, $p = 0.002$) and testis size ($r = -0.764$, $p = 0.01$). Extensive studies on the relationship between PBDEs and sperm quality are required.

Keywords: polybrominated diphenyl ethers; flame retardants; human serum; sperm.

* Correspondence to: Kazuhiko Akutsu Phone: +81-6-6972-1321 FAX: +81-6-6972-2393 E-mail: akutu@iph.pref.osaka.jp

1 Polybrominated diphenyl ethers (PBDEs) are used as flame retardants in the
2 production of common consumer products such as electronics, furniture, and
3 textiles. PBDEs are currently recognized as environmental pollutants of global
4 concern because their levels in the environment and in humans have increased
5 markedly over the past several decades (Meironyté et al., 1999; Ikononou et al.,
6 2002; Akutsu et al., 2003). Since PBDEs are somewhat structurally similar to
7 thyroid hormones such as thyroxine (T4), it was speculated that PBDEs might
8 mimic thyroid hormones and disrupt thyroid homeostasis. Several studies indicate
9 that exposure to PBDEs can decrease the circulating levels of T4 in laboratory
10 animals (Fowles et al., 1994; Zhou et al., 2002) and can cause permanent
11 neurological effects similar to those associated with thyroid hormone deficiencies
12 (Eriksson et al., 2001; Viberg et al., 2004). In addition, several PBDEs possess
13 weak estrogenic/antiestrogenic activities (Meerts et al., 2001). The proliferation
14 and differentiation of Sertoli cells and sperm production are regulated by thyroid
15 and sex hormones. Thus, PBDEs might affect male reproductive health by
16 interfering with the thyroid- and sex-hormone functions. Kuriyama et al. (2005)
17 have reported that developmental exposure to a single low dose (60 µg/kg body
18 weight) of 2,2',4,4',5-pentabromodiphenyl ether (PeBDE-99) decreased the sperm
19 count in male Wistar rats. However, no previous studies have examined the
20 relationship between human PBDE levels and sperm quality.

21
22 We participated in an international project examining the sperm quality of fertile
23 males and found that the sperm concentration of Japanese males was lower than
24 that of European males (Iwamoto et al., 2006). The examination of sperm quality
25 and an estimation of the concentration of chemicals in the serum would be
26 required to reveal the correlation between chemical exposure and the sperm
27 quality in Japanese males. The aim of this pilot study was to measure PBDEs in
28 serum samples from young Japanese males and to examine the relationship
29 between serum PBDE levels and sperm quality.

30 31 **MATERIALS AND METHODS**

32 This study was performed in accordance with the protocols which were approved
33 by the ethical committees of the St. Marianna University School of Medicine and
34 Osaka Prefectural Institute of Public Health. Written informed consent was
35 obtained from all study participants. Blood serum and sperm samples were
36 collected on a monthly basis in the year 2003 from 45 young Japanese males at
37 the Department of Urology, St. Marianna University School of Medicine. The
38 participants were instructed to abstain from ejaculation for at least 48 h prior to
39 sperm collection. The blood samples were collected in vacuum tubes, and the
40 serum fractions were separated by centrifugation. The serum samples were stored
41 at -80°C until analysis. Of the 45 sample sets, 10 were randomly selected for this
42 study. For PBDE analysis, 10 pooled serum samples (0.5 g × 12 months; total, 6 g
43 per person) were prepared, and each pool was regarded as a representative sample
44 of each set. The mean ± standard deviation (SD) of the age of the 10 participants
45 was 22 ± 1 years (range, 18–22 years). The mean ± SD abstinence period was 3.1
46 ± 0.4 days (range, 2.6–3.8 days). In addition, 2 brands of commercially pooled

human serum ("L-Consera N" and "L-Suitrol I," Nissui Pharmaceutical, Tokyo, Japan) were used as in-house reference materials.

Standard mixture solutions of native PBDEs (BDE-AAP-A-15X) were purchased from AccuStandard (New Haven, CT, USA), and $^{13}\text{C}_{12}$ -labeled PBDEs (MBDE-MXC) were purchased from Wellington Laboratories (Ontario, Canada). In this study, 29 PBDE congeners with 3 to 7 bromine atoms were monitored. The PBDE numbers are assigned according to the International Union of Pure and Applied Chemistry nomenclature for polychlorinated biphenyls. Acetone, acetonitrile, and *n*-hexane of pesticide analysis grade; ammonium sulfate of biochemistry grade; 44% sulfuric acid-impregnated silica gel; and *n*-nonane of dioxin analysis grade were purchased from Wako Pure Chemical Industries (Osaka, Japan). Water was deionized and purified using a Milli-Q cartridge system (Millipore, Bedford, MA, USA).

Sperm analyses were performed at the Department of Urology, St. Marianna University School of Medicine, according to World Health Organization criteria as described elsewhere (World Health Organization, 1999; Iwamoto et al., 2006).

Serum samples were analyzed at Osaka Prefectural Institute of Public Health. The serum sample (6 g) was extracted using ethanol/*n*-hexane (1:3 v/v, 14 mL) in a 50 mL test tube after adding $^{13}\text{C}_{12}$ -labeled surrogate standards ($^{13}\text{C}_{12}$ -2,4,4'-tribromodiphenyl ether ($^{13}\text{C}_{12}$ -TrBDE-28), $^{13}\text{C}_{12}$ -2,2',4,4'-tetrabromodiphenyl ether ($^{13}\text{C}_{12}$ -TeBDE-47), $^{13}\text{C}_{12}$ -2,2',4,4',5-pentabromodiphenyl ether ($^{13}\text{C}_{12}$ -PeBDE-99), $^{13}\text{C}_{12}$ -2,2',4,4',5,5'-hexabromodiphenyl ether ($^{13}\text{C}_{12}$ -HxBDE-153), $^{13}\text{C}_{12}$ -2,2',4,4',5,6'-HxBDE ($^{13}\text{C}_{12}$ -HxBDE-154), and $^{13}\text{C}_{12}$ -2,2',3,4,4',5',6-heptabromodiphenyl ether ($^{13}\text{C}_{12}$ -HpBDE-183); 10 pg for each congener) and 3.6 mL saturated ammonium sulfate solution. The test tube was shaken for 30 min and then centrifuged for 10 min at 3000 rpm. The *n*-hexane phase was collected, and the aqueous phase was re-extracted twice with 12 mL *n*-hexane. The 3 *n*-hexane phases were combined and washed with 12 mL water. After evaporation of the solvent, the lipid content was determined gravimetrically using a semimicro balance (Sartorius RC210P, Goettingen, Germany). The lipid was dissolved in *n*-hexane and transferred to a column of 44% sulfuric acid-impregnated silica gel (3 g). The column was eluted with 30 mL *n*-hexane, and the eluate was evaporated to 2 mL. The *n*-hexane solution was transferred to a test tube and partitioned with *n*-hexane-saturated acetonitrile (4 mL) 3 times by shaking the test tube for 10 min and then centrifuging for 10 min at 3000 rpm. The acetonitrile phase was combined and then evaporated to dryness. The residue was redissolved in *n*-hexane and transferred to a microconcentration tube. After addition of the injection standard ($^{13}\text{C}_{12}$ -3,3',4,4',5-PeBDE ($^{13}\text{C}_{12}$ -PeBDE-126)) and keeper solvent (10 μL *n*-nonane), the extract was finally evaporated to approximately 10 μL under a gentle stream of nitrogen. The serum extract was assayed by a gas chromatography/mass spectrometry (GC/MS) system (Agilent 6890A GC coupled with JEOL JMS-GCmateII, Tokyo, Japan) using a fused silica capillary column (Rtx-1MS, 15 m length, 0.25 mm i.d., 0.1 μm film thickness; Restek, Bellefonte,

PA, USA). For each compound, 2 ions of the molecular ion or fragment ion cluster were monitored. Quantitation was based on the isotope dilution method using $^{13}\text{C}_{12}$ -labeled internal standards. The PBDE concentrations were adjusted for total serum lipids and were expressed in units of nanogram per gram lipid weight (ng/g lw). TeBDE-47, PeBDE-99, PeBDE-100, and HxBDE-153 were of interest because they are dominant in human serum.

We validated the serum extraction procedure prior to beginning sample analysis by analyzing 4 replicate samples of pooled serum fortified with target analytes at 0.04–0.1 ng/g serum. The mean percent recovery of 7 representative PBDE congeners (TrBDE-28, TeBDE-47, PeBDE-99, PeBDE-100, HxBDE-153, HxBDE-154, and HpBDE-183) ranged from 91% to 107%, and the relative standard deviation (RSD) ranged from 2% to 10%. The limit of detection (LOD) and limit of quantification (LOQ) were defined as 3 times and 10 times the SD values obtained from the analysis of the 7 procedural blank samples (6 g of water), respectively. However, for congeners that could not be detected in the blanks, the values that were 3 times and 10 times the SD values obtained from the analysis of 5 replicates of the lowest calibration standard were used as LOD and LOQ. The LOD values for all the PBDE congeners were below 0.3 ng/g lw. In the analysis of 3 split unfortified serum samples, the RSD values for all the detected congeners were below 10%.

RESULTS AND DISCUSSION

Of the 29 PBDE congeners monitored, 4 congeners (TeBDE-47, PeBDE-99, PeBDE-100, and HxBDE-153) were mainly detected in human serum samples (Figure 1). The concentrations of the detected PBDE congeners in the serum samples ($n = 10$) are shown in Table 1. The median levels of the individual PBDE congeners were as follows: TeBDE-47, 1.4 ng/g lw; PeBDE-99, 0.21 ng/g lw; PeBDE-100, 0.24 ng/g lw; and HxBDE-153, 0.72 ng/g lw. The levels of total PBDEs in Japanese human serum samples were almost the same as those reported in European countries but were 1 order of magnitude lower than those reported in USA (Hites, 2004). Significant positive correlations were observed between the concentrations of TeBDE-47 and PeBDE-99 ($r = 0.988$, $p < 0.001$), TeBDE-47 and PeBDE-100 ($r = 0.938$, $p < 0.001$), and PeBDE-99 and PeBDE-100 ($r = 0.915$, $p < 0.001$). In contrast, no significant correlations were observed between the concentration of HxBDE-153 and those of the other 3 congeners ($r = 0.306$ – 0.390 , $p = 0.26$ – 0.39). The absence of a significant correlation between HxBDE-153 and the other 3 dominant congeners (TeBDE-47, PeBDE-99, and PeBDE-100) implies that the main sources and/or biological properties of HxBDE-153 were different from those of the other 3 congeners. It has been reported that the technical mixtures of pentaBDE (DE-71 and Bromkal 70-5DE) and octaBDE (DE-79 and Bromkal 79-8DE) both contained HxBDE-153 in the range 5.32–5.44% w/w and 0.15–8.66% w/w, respectively (La Guardia et al., 2006). The congeners TeBDE-47, PeBDE-99, and PeBDE-100 have been found in pentaBDE as the major components, but they have not been found in octaBDE (La Guardia et al., 2006).

1 These 3 congeners and HxBDE-153 have never been found in a technical
2 decaBDE mixture (Saytex 102E and Bromkal 82-0DE) (La Guardia et al., 2006).
3 Therefore, TeBDE-47, PeBDE-99, and PeBDE-100 are mainly sourced from
4 pentaBDE, although HxBDE-153 is sourced from both pentaBDE and octaBDE.
5 In the early 1990s, Japanese manufacturers voluntarily stopped the production and
6 use of pentaBDE because its potency to accumulate in the biota and produce toxic
7 polybrominated dibenzofurans/dioxins under thermal stresses was a cause of
8 concern. However, the production and use of octaBDE were continued in Japan
9 until 2002 (Ministry of the Environment, Japan, 2006). Therefore, many consumer
10 products containing octaBDE in the Japanese indoor environment might continue
11 to exist. Thus, with regard to octaBDE components such as HxBDE-153 and
12 HpBDE-183, inhalation and dermal exposure might be important exposure routes
13 for the Japanese people. Geyer et al. (2004) have predicted elimination half-lives
14 of PBDEs in the human adipose tissue; the predicted half-lives of individual
15 congeners in an adult male were as follows: TeBDE-47, 1.9 years; PeBDE-99, 3.5
16 years; PeBDE-100, 2.4 years; and HxBDE-153, 7.8 years. It is expected that the
17 half-lives of Te-HxBDEs increase with the number of bromine atoms per
18 molecule, and the half-life of HxBDE-153 is much longer than those of other
19 dominant congeners detected in human serum. Further research is needed to
20 examine the difference between the elimination half-lives and toxicity of
21 individual PBDE congeners in animals and humans.

22
23 The sperm concentration and testis size of the 10 participants are shown in Table
24 2. The sperm concentration of these participants ranged from 25 to 115
25 million/mL. No participant had a sperm concentration below 20 million/mL, the
26 minimum fertility standard established by the World Health Organization (World
27 Health Organization, 1999). Strong inverse correlations were observed between
28 the serum HxBDE-153 concentration and sperm concentration ($r = -0.841$, $p =$
29 0.002 ; Figure 2) and testis size ($r = -0.764$, $p = 0.01$). However, no significant
30 relationships were observed between the serum concentrations of the other
31 congeners and the sperm concentration (r ranged from -0.187 to -0.099 , $p =$
32 0.605 – 0.786) or testis size (r ranged from -0.216 to -0.054 , $p = 0.548$ – 0.883).
33 Researchers have hypothesized that endocrine-disrupting chemicals with thyroid-
34 hormonal or sex-hormonal activities might adversely affect male fertility. The
35 thyroid-disrupting and estrogenic/antiestrogenic activities of PBDEs have been
36 reported in several studies (Meerts et al., 2001; Zhou et al., 2002). In addition,
37 considerable evidence regarding the reproductive effects of PBDEs is available
38 from *in vivo* studies. Kuriyama et al. (2005) have reported that developmental
39 exposure to a single low dose (60 $\mu\text{g/kg}$ body weight) of PeBDE-99 decreased the
40 sperm count in male Wistar rats. Although the levels of PBDEs found in our study
41 are relatively low, we observed significant inverse associations between the serum
42 concentration of HxBDE-153 and the sperm concentration and testis size; this
43 suggests an association between the serum HxBDE-153 concentration and human
44 sperm quality. The lack of a significant relationship among other individual PBDE
45 congeners and sperm parameters might indicate a difference in bioactivity
46 between the congeners. The relationship between PBDEs and sperm quality is a

1 complicated problem and needs further study.

2
3
4 *Acknowledgments* We thank the participants who donated their blood and semen
5 samples. This study was supported by grants from the Ministry of the
6 Environment and the Ministry of the Health, Labor and Welfare, Japan.

8 REFERENCES

- 9
10 Akutsu K, Kitagawa M, Nakazawa H, Makino T, Iwazaki K, Oda H, Hori S
11 (2003) Time-trend (1973-2000) of polybrominated diphenyl ethers in Japanese
12 mother's milk. *Chemosphere* 53:645-654.
13 Eriksson P, Jakobsson E, Fredriksson A (2001) Brominated flame retardants: a
14 novel class of developmental neurotoxicants in our environment? *Environ*
15 *Health Perspect* 109:903-908.
16 Fowles JR, Fairbrother A, Baecher-Steppan L, Kerkvliet NI (1994) Immunologic
17 and endocrine effects of the flame-retardant pentabromodiphenyl ether (DE-
18 71) in C57BL/6J mice. *Toxicology* 86:49-61.
19 Geyer HJ, Schramm K-W, Darnerud PO, Aune M, Feicht A, Fried KW,
20 Henkelmann B, Lenoir D, Schmid P, McDonald TA (2004) Terminal
21 elimination half-lives of the brominated flame retardants TBBPA, HBCD, and
22 lower brominated PBDEs in humans. *Organohalogen Comp* 66:3867-3872.
23 Hites RA (2004) Polybrominated diphenyl ethers in the environment and in
24 people: a meta-analysis of concentrations. *Environ Sci Technol* 38:945-956.
25 Ikononou MG, Rayne S, Addison RF (2002) Exponential increases of the
26 brominated flame retardants, polybrominated diphenyl ethers, in the Canadian
27 Arctic from 1981 to 2000. *Environ Sci Technol* 36:1886-1892.
28 Iwamoto T, Nozawa S, Yoshiike M, Hoshino T, Baba K, Matsushita T, Tanaka
29 SN, Naka M, Skakkebaek NE, Jorgensen N (2006) Semen quality of 324 fertile
30 Japanese men. *Hum Reprod* 21:760-765.
31 Kuriyama SN, Talsness CE, Grote K, Chahoud I (2005) Developmental exposure
32 to low dose PBDE 99: effects on male fertility and neurobehavior in rat
33 offspring. *Environ Health Perspect* 113:149-154.
34 La Guardia MJ, Hale RC, Harvey E (2006) Detailed polybrominated diphenyl
35 ether (PBDE) congener composition of the widely used penta-, octa-, and deca-
36 PBDE technical flame-retardant mixtures. *Environ Sci Technol* 40:6247-6254.
37 Meerts IA, Letcher RJ, Hoving S, Marsh G, Bergman Å, Lemmen JG, van der
38 Burg B, Brouwer A (2001) In vitro estrogenicity of polybrominated diphenyl
39 ethers, hydroxylated PBDEs, and polybrominated bisphenol A compounds.
40 *Environ Health Perspect* 109:399-407.
41 Meironyté D, Norén K, Bergman Å (1999) Analysis of polybrominated diphenyl
42 ethers in Swedish human milk. A time-related trend study, 1972-1997. *J*
43 *Toxicol Environ Health A* 58:329-341.
44 Ministry of the Environment, Japan (2006) Regarding brominated flame
45 retardants (reference data 3). In: report on the survey of emissions of
46 brominated dioxins and related compounds in fiscal year 2004. (in Japanese).

1 Available: <http://www.env.go.jp/air/report/h18-07/index.html> [accessed 20
2 June, 2007].
3 Viberg H, Fredriksson A, Eriksson P (2004) Investigations of strain and/or gender
4 differences in developmental neurotoxic effects of polybrominated diphenyl
5 ethers in mice. *Toxicol Sci* 81:344-353.
6 World Health Organization (1999) WHO Laboratory Manual for the Examination
7 of Human Semen and Sperm-Cervical Mucus Interaction (4th Edition).
8 Cambridge University Press, Cambridge.
9 Zhou T, Taylor MM, DeVito MJ, Crofton KM (2002) Developmental exposure to
10 brominated diphenyl ethers results in thyroid hormone disruption. *Toxicol Sci*
11 66:105-116.
12
13
14

1 Figure legends

2

3 Figure 1 Chromatograms of PBDEs in human serum (participant No.2) and
4 standard solution (1 to 2.5 ng/mL each)

5 Figure 2 Relationship between the serum HxBDE-153 concentration and
6 sperm concentration

7

Table 1 Concentrations of PBDEs in serum samples from 10 Japanese males (ng/g lw)

Congener	Participant No.									
	1	2	3	4	5	6	7	8	9	10
TrBDE-17	tr <0.04	tr <0.05	nd <0.01	nd <0.01	nd <0.02	nd <0.01	nd <0.02	nd <0.01	nd <0.02	nd <0.02
TrBDE-28/33	tr <0.2	0.37	0.16	tr <0.2	0.16	0.24	tr <0.2	0.17	tr <0.2	tr <0.2
TrBDE-37	tr <0.02	tr <0.03	nd <0.01	nd <0.01	nd <0.01	nd <0.01	nd <0.01	nd <0.01	nd <0.01	nd <0.01
TeBDE-49	nd <0.02	nd <0.03	0.09	tr <0.07	tr <0.08	0.07	nd <0.02	0.09	nd <0.03	tr <0.09
TeBDE-47	1.3	5.9	1.5	0.96	1.6	1.8	0.54	2.9	0.93	0.81
TeBDE-66	nd <0.04	nd <0.05	nd <0.04	nd <0.04	nd <0.04	nd <0.04	nd <0.04	tr <0.2	nd <0.05	nd <0.05
PeBDE-100	0.23	0.67	0.24	0.21	0.24	0.40	0.13	0.31	0.21	0.25
PeBDE-99	0.21	1.1	0.21	0.16	0.25	0.21	0.10	0.49	0.15	0.20
PeBDE-118	0.02	0.03	tr <0.02	tr <0.02	0.02	0.03	tr <0.02	tr <0.02	0.03	0.03
PeBDE-85	tr <0.07	tr <0.09	tr <0.07	nd <0.02	tr <0.08	nd <0.02	nd <0.02	tr <0.07	nd <0.03	nd <0.02
HxBDE-155	nd <0.02	tr <0.07	tr <0.05	tr <0.05	nd <0.02	tr <0.06	nd <0.02	nd <0.02	tr <0.07	nd <0.02
HxBDE-154	tr <0.06	0.08	0.05	0.05	tr <0.06	0.06	tr <0.06	tr <0.06	tr <0.07	tr <0.07
HxBDE-153	0.76	0.96	1.1	0.56	0.58	0.68	0.37	0.52	0.91	0.79
HpBDE-183	nd <0.1	nd <0.2	tr <0.4	tr <0.4	tr <0.4	tr <0.4	nd <0.1	nd <0.1	tr <0.5	nd <0.2
Sum of 4 PBDEs ^a	2.5	8.6	3.1	1.9	2.7	3.1	1.1	4.2	2.2	2.1

Abbreviations: tr, trace; nd, not detected. ^aSum of TeBDE-47, PeBDE-100, PeBDE-99, and HxBDE-153.

Table 2 Sperm concentration and testis size of 10 Japanese males

	Participant No.									
	1	2	3	4	5	6	7	8	9	10
Sperm concentration (million/mL)^a	49	55	38	108	83	74	115	78	25	30
Testis size (mL)^b	36	36	40	50	46	42	51	54	29	33

^aAnnual average of monthly data. ^bTotal of right and left testes.

Fig.1

Fig.2