The first total synthesis of bufobutanoic acid by two routes based on nucleophilic substitution reaction on indole nucleus

著者	Kurauchi Takashi, Nagahama Yoshiyuki, Hasegawa Masakazu, Yamada Koji, Somei Masanori
journal or publication title	Heterocycles
volume	53
number	5
page range	1017-1019
year	2000-05-01
URL	http://hdl.handle.net/2297/4358

doi: <https://doi.org/10.3987/com-00-8870>

THE FIRST TOTAL SYNTHESIS OF BUFOBUTANOIC ACID BY TWO ROUTES BASED ON NUCLEOPHILIC SUBSTITUTION REACTION ON INDOLE NUCLEUS¹

Takashi Kurauchi, Yoshiyuki Nagahama, Masakazu Hasegawa, Koji Yamada,
and Masanori Somei*

Faculty of Pharmaceutical Sciences, Kanazawa University,
13-1 Takara-machi, Kanazawa 920-0934, Japan

Abstract — Regioselective nucleophilic substitution reaction of 1-hydroxytryptamines led to establish two novel routes for the first synthesis of bufobutanoic acid. An effective synthesis of 5-benzyloxytryptamine from tryptamine is also reported.

In 1999, Kamano and co-workers² isolated bufobutanoic acid (**1a**, Scheme 1) as a cytotoxic substance against murine P388 lymphocytic leukemia cells from Ch'an Su and determined its structure. From our ongoing project for developing biologically active novel compounds,³ we have much interested in **1a** and intended to establish a methodology applicable for producing its various congeners. To meet our end, we initially needed simple synthesis of **1a**. Now, we have succeeded in developing two routes based on 1-hydroxyindole chemistry.⁴

The first route is the one utilizing 1-hydroxy-*Nb*-methoxycarbonyltryptamine (**3a**) as an intermediate, a potent inhibitor of platelet aggregation.⁵ Thus, **3a**, obtained in three steps from tryptamine (**2**) in 62% overall yield as described before,⁶ was converted to **4b** in 48% yield by the regioselective hydroxylation at the 5-position upon the reaction with 85% HCOOH at room temperature for 24 h. Interestingly, the corresponding 1-methoxy-*Nb*-methoxycarbonyltryptamine⁶ (**3b**) provided **4a** selectively in 69% yield by the similar treatment with 85% HCOOH at 80°C for 20 min. Subsequent reaction of **4a** with 85% HCOOH at room temperature for 2 days provided **4b** in 70% yield together with 10% yield of starting material.

The reaction of **4b** with benzyl bromide in the presence of K₂CO₃ in DMF afforded **4c** in 94% yield. Alkaline hydrolysis of **4c** with 10% NaOH in refluxing MeOH provided 96% yield of 5-benzyloxytryptamine (**5**).⁷ With an useful building block for preparing various serotonin derivatives in hand, it was converted to **6** in 96% yield by the reaction with succinic anhydride in THF. Catalytic hydrogenation of **6** over 10% Pd/C at room temperature produced **1a** in 99% yield. The spectra of **1a** are identical with those reported in the literature.² Thus, the first synthesis of **1a** was achieved in eight steps from **2** in 25% overall yield with 33% originality rate.⁸

As the second one, six-steps synthesis of **1a** in 13% overall yield with 43% originality rate was developed. Tryptamine (**2**) was initially reacted with succinic anhydride in THF at room temperature, followed by methylation with CH₂N₂ in one pot procedure to give *Nb*-methoxysuccinyltryptamine (**7**) in 89% yield. Subsequent reduction of **7** with Et₃SiH in CF₃COOH⁹ at 60 °C provided the corresponding 2,3-dihydroindole (**8**) in 99% yield. Our 1-hydroxyindole synthetic method using Na₂WO₄·2H₂O⁴ and

Scheme 1

1) CH₂N₂; 2) 85% HCOOH; 3) PhCH₂Br, K₂CO₃, DMF; 4) 10% NaOH, MeOH; 5) succinic anhydride, THF; 6) 10% Pd/C, H₂; 7) Et₃SiH, CF₃COOH; 8) Na₂WO₄·2H₂O, 30% H₂O₂; 9) 1_M K₂CO₃, MeOH.

30% H₂O₂ at room temperature was successfully applied to **8** giving the desired 1-hydroxytryptamine (**9a**) in 56% yield. Structure of **9a** was confirmed by converting it to 1-methoxytryptamine (**9b**) in 86% yield by the reaction with CH₂N₂. Then, **9a** was treated with 85% HCOOH at 50°C for 50 min to give serotonin derivative (**1b**) in 38% yield. Finally, ester part of **1b** was hydrolyzed with 1_M K₂CO₃ in MeOH at 50°C to provide **1a** in 70% yield.

In conclusion, we have disclosed that nucleophilic substitution reaction¹⁰ of 1-hydroxytryptamines¹¹ is a suitable methodology for the preparations of serotonin congeners.

ACKNOWLEDGMENT

This work is supported in part by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, Sports, and Culture, Japan, which is gratefully acknowledged.

REFERENCES AND NOTES

1. This is Part 96 of a series entitled "The Chemistry of Indoles". Part 95: M. Somei, K. Noguchi, R. Yamagami, Y. Kawada, K. Yamada, and F. Yamada, *Heterocycles*, 2000, **53**, 7. All new compounds

gave satisfactory spectral and elemental analysis or high-resolution MS data for crystals or gums, respectively. **1b**, gum; **4a**, gum; **4b**, gum; **4c**, gum; **5**, mp 97.5—99.5°C; **6**, mp 145—147°C; **7**, mp 118—120°C; **8**, mp 74—75°C; **9a**, mp 151.5—153.5°C.

2. Y. Kamano, H. Morita, R. Takano, A. Kotake, T. Nogawa, H. Hashima, K. Takeya, H. Itokawa, and G. R. Pettit, *Heterocycles*, 1999, **50**, 499.
3. K. Nakagawa and M. Somei, *Heterocycles*, 1991, **32**, 873; F. Yamada, K. Kobayashi, A. Shimizu, N. Aoki, and M. Somei, *ibid.*, 1993, **36**, 2783; F. Yamada, S. Hamabuchi, A. Shimizu, and M. Somei, *ibid.*, 1995, **41**, 1905; M. Somei, Y. Fukui, and M. Hasegawa, *ibid.*, 1995, **41**, 2157; M. Somei, H. Hayashi, T. Izumi, and S. Ohmoto, *ibid.*, 1995, **41**, 2161; M. Somei, K. Yamada, M. Hasegawa, M. Tabata, Y. Nagahama, H. Morikawa, and F. Yamada, *ibid.*, 1996, **43**, 1855; M. Somei, F. Yamada, T. Izumi, and M. Nakajou, *ibid.*, 1997, **45**, 2327; F. Yamada, M. Tamura, and M. Somei, *ibid.*, 1998, **49**, 451; M. Somei, N. Oshikiri, M. Hasegawa, and F. Yamada, *ibid.*, 1999, **51**, 1237. See also reference 11.
4. Review: M. Somei, *Heterocycles*, 1999, **50**, 1157 and references cited therein.
5. M. Somei, K. Yamada, M. Hasegawa, M. Tabata, Y. Nagahama, H. Morikawa, and F. Yamada, *Heterocycles*, 1996, **43**, 1855.
6. M. Somei, K. Kobayashi, K. Tanii, T. Mochizuki, Y. Kawada, and Y. Fukui, *Heterocycles*, 1995, **40**, 119.
7. Although hydrochloride of **5** is commercially available from Sigma, it is expensive and therefore not suitable as a common starting material for multi-gram scale production of serotonin congeners. Our present method seems to be better to obtain **5** at cheaper cost compared to the conventional one. Another choice is to utilize serotonin hydrochloride as a starting material.
8. M. Somei, *J. Synth. Org. Chem.*, 1982, **40**, 387; M. Somei, Y. Makita, and F. Yamada, The 3rd International Kyoto Conference on New Aspects of Organic Chemistry, Abstracts Papers, Nov., 1985, p. 128; M. Somei, *Yakugaku Zasshi*, 1988, **108**, 361.
Originality rate is the result of the following calculation:
Originality Rate (%) = 100 x [Number of Newly Developed Steps + 1] ÷ [Total Number of Synthetic Steps + 1]
9. A. E. Lanzilotti, R. Littell, W. J. Fanshawe, T. C. McKenzie, and F. M. Lovell, *J. Org. Chem.*, 1979, **44**, 4809.
10. J. A. Joule, "Progress in Heterocyclic Chemistry", Vol. 11, ed. by G. W. Gribble and T. L. Gilchrist, Elsevier Science Ltd., Oxford, 1999, pp. 45—65.
11. M. Somei, H. Morikawa, K. Yamada, and F. Yamada, *Heterocycles*, 1998, **48**, 1117; M. Hasegawa, K. Yamada, Y. Nagahama, and M. Somei, *ibid.*, 1999, **51**, 2815.

Received, 8th February, 2000