Complexation of Am with size-fractionated soil humic acids

著者	Nagao S., Aoyama M., Watanabe A., Tanaka T.
雑誌名	年報 金沢大学環日本海域環境研究センター = K-INET Kanazawa University
巻	平成20年度
号	2008
ページ	70-70
発行年	2010-01-29
URL	http://hdl.handle.net/2297/30152

Complexation of Am with size-fractionated soil humic acids

S. Nagao¹, M. Aoyama², A. Watanabe³, T. Tanaka⁴

¹Faculty of Environmental Earth Science, Hokkaido University,

²Faculty of Agriculture and Life Science, Hirosaki University ,

³Graduate school of Bioagricultural Sciences, Nagoya University,

⁴Tokai Research and Development Center, Japan Atomic Energy Agency

Humic substances (humic and fulvic acids) are ubiquitous in nature and have many beneficial effects on soils and aquatic environments. They play important role in geochemical behavior of trace elements such as actinides from fallout and local sources in aquatic environments. Therefore, characteristics of actinides bound to humic and fulvic acids can yield information on the geochemical role of humic substances in the migration of actinides. In this study, the association properties of Am with humic substances were studied on the basis of structural feature of humic substances. We separated two humic acids ('HA': fluorescence-poor and higher molecular size fraction, 'FL': fluorescence-rich and lower molecular size fraction) with different structural features from humic acids prepared from three types of soils (andisol, entisol and buried andisol) by Sephadex G-25. The association experiments were carried out in a medium of 0.01M NaClO₄ solution at a humus concentration of 10 mg/l and pH6-8. After finishing the association experiments, ultrafiltration was used to size-fractionate Am in the humus solutions. Figure 1 shows molecular size distribution of Am in the presence of size-fractionated HAs. The dominant size fractions of Am in the presence of 'HA' were 450nm-100k Da (48-53% for the andisol and entisol) and 100k-30k Da (60% for the buried andisol). On the other hand, 43-51% of Am was presented in the size of 30-10k Da in the 'FL' HA solutions from three soil types. These results indicate that characteristics of size-fractionated humic acids are related association properties with Am.

Fig. 1 Size distribution of Am in the presence of HAs.