Nursing care in multifunctional small group homes providing day, visiting and overnight services for older people living at home

著者	片平 伸子
著者別表示	Katahira Nobuko
journal or publication title	博士論文本文Full
学位授与番号	13301甲第4532号
学位名	博士(保健学)
学位授与年月日	2017-03-22
URL	http://hdl.handle.net/2297/46949

doi: <https://doi.org/10.1111/ijn.12482>

✧ RESEARCH PAPER ✧

Nursing care in multifunctional small group homes providing day, visiting and overnight services for older people living at home

Nobuko Katahira MLIS, PHN, RN

Division of Health Sciences, Graduate School of Medical Sciences, Kanazawa University, Kanazawa, Japan

Department of Nursing and Rehabilitation, Konan Women's University, Kobe, Japan

Keiko Tsukasaki DSN, PHN, RN

Faculty of Health Sciences, Institute of Medical, Pharmaceutical and Health Sciences, Kanazawa University, Kanazawa, Japan

Accepted for publication 13 August 2016

Katahira N, Tsukasaki K. *International Journal of Nursing Practice* 2016; 22: 605–615

Nursing care in multifunctional small group homes providing day, visiting and overnight services for older people living at home

Home care programs for older people have been developed around the world. Nurses are key to these programs. The aim of this study is to explore details of the nursing activities used in group homes to provide a basis for describing effective nursing practices in such facilities. A self-administered questionnaire was sent to 240 randomly selected facilities throughout Japan in 2013. Responses indicated that the activities of nurses in managing the health of older people included determining the need for medical consultations, dealing with emergencies, and making arrangements for the use of flexible care services. Nursing activities were directly related to the percentage of older people in each facility with diagnoses such as dementia or heart disease. Nurses reported low general self-efficacy for some of the more specialized nursing activities they performed. Nursing activities are performed with the aim of supporting older people with high health care needs to continue living at home; are tailored to the characteristics of residents and contribute to the provision of timely health examinations and flexible arrangement of services. Findings indicate that it might be advantageous to increase the nursing staffing at these facilities, to provide care guidelines and training opportunities to increase nurse self-efficacy.

Key words: frail elderly, group homes, nursing care, referral and consultation.

SUMMARY STATEMENT

What is already known about this topic?

- Day services, visiting services and short-stay care services are currently provided to enable older people who require care or assistance to continue living in their homes by supporting their activities of daily living and health.

- Such services have previously been provided by individual day, visiting or overnight care service facilities. Multifunctional small group homes are newly established facilities that can comprehensively provide these services.
- Nurses have important roles at these facilities but details of their activities have not yet been clarified.

What this paper adds:

- Nursing activities in multifunctional small group homes include health management, determination of the need for medical consultation and making judgments in emergencies during day, home or overnight care service hours.

Correspondence to: Nobuko Katahira, Department of Nursing, Faculty of Nursing and Rehabilitation, Konan Women's University, 6-2-23 Morikita-machi, Higashinada-ku, Kobe, Hyogo, 658-0001, Japan.
E-mail: katahira-n@konan-wu.ac.jp

- Nursing activities vary in relation to residents' clinical state.
- In 25% of study facilities the content of care services was changed based on nurses' judgments.

The implications of this paper:

- Policy makers and managers should consider increasing the number of nurses in facilities with a large number of residents and examining their work patterns to provide opportunities for care enhancement.
- Policy makers and managers should consider providing training opportunities to increase nurses' self-efficacy and promote their activities, as well as to develop nursing activity guidelines.

INTRODUCTION

Home care for older people has been promoted as a way to improve conventional medical and welfare systems and reduce medical costs in countries with rapidly ageing populations, and many countries are exploring new models and approaches to provide such services. For example, the Program of All-Inclusive Care for the Elderly (PACE) in the United States provides comprehensive medical care and long-term care services to support older individuals living at home. PACE programs have reduced hospitalisation rates among enrollees and have encouraged continuation of home-based care.^{1,2} These programs also improve functional outcomes for older people, and reduces the cost of care.³ Aging in Place (AIP) programs have also contributed to positive health outcomes among group home residents.^{4,5} In the Netherlands, an assessment and management program provided to frail older individuals has successfully improved the detection and management of dementia.^{6,7} Nurses are an integral part of these home-based care programs and have contributed greatly to their findings.

In Japan, the Long-term Care Insurance System was established to support independent community living for older people⁸ and to support the launch of multifunctional small group home services in 2006.⁹ As Japan's population ages, the number of older people requiring long-term care and support is increasing rapidly.¹⁰ To help these individuals remain well and living in their communities, any changes to their situation must be identified, so that community-based flexible care and support can be provided. Because the provision of long-term care in a small and home-like environment has been shown to be an effective way to improve the quality of life and cognitive/functional status of older people,^{11,12} multifunctional small group home services were the chosen form of support. These 'group homes'

are facilities that support up to 25 older community residents to carry on with activities of daily living, by flexible provision of various types of care services, on a 24 h basis. Day care is the main service; overnight and visiting care are provided when necessary. Group homes are required to employ nurses, and group home care is regarded as a new field in nursing. Although a qualitative study has examined the roles of nurses and care workers in group homes,¹³ details of the nursing activities in these facilities have not yet been clarified.

Identifying the types of nursing activities provided in group homes as well as the related characteristics of the facilities, service residents and nurses might contribute to developing measures to promote the provision of effective nursing services. This might also provide a basis for development of services to support older people living at home in other countries with rapidly ageing populations.

Aims

The aim of this study was to explore details of the nursing activities used in group homes to provide a basis for describing effective nursing practices in such facilities. Our research questions were:

1. What types of nursing activities are performed in group homes?
2. What are the characteristics of facilities, service residents and nurses that are linked to particular nursing activities?

METHODS

Design

This study used a descriptive, cross-sectional survey design.

Sample

A random sample was selected of 1000 group homes from the 3192 in Japan, registered with the Welfare and Medical Service Network System (WAMNET).¹⁴ Questionnaires were distributed to administrators and nurses at each selected group home.

Data collection

Two separate versions of the survey were developed by the research team.

One survey, for administrators, contained questions on: (1) facility attributes (business entity, facilities on site, and number of staff members and residents); and (2) resident attributes (age, Long-term Care Grade, family compositions).

The second survey, for nursing staff, included questions on: (1) residents' health condition (disease, identified need for medical care or consultation, medication management, and any emergency management required within the last 3 months); (2) nurse attributes (age, years of experience working at medical institutions, years of experience working at the present facility and working pattern); (3) 17 items based on nursing activities used within the last month, such as monitoring residents' health conditions and medication management,^{13,15} and (4) the self-efficacy of nurses, as measured by the General Self-Efficacy Scale (GSES).^{16,17} Self-efficacy was examined to assess what nurses believed they could do with what was available under a variety of circumstances.¹⁸

Two anonymous self-administered questionnaires with a stamped self-addressed envelope for postal return of the questionnaires were distributed at one time by mail to administrators and nurses in January 2013. We decided that we were unlikely to receive any additional responses after the end of March 2013.

Data analysis

Descriptive statistics including mean, standard deviation and percentages were used to summarize characteristics of facilities, residents and nurses. Facility, resident, and nursing characteristics were dichotomized into higher and lower value subgroups using group mean values. Relationships among facility, resident and nurse attributes and usage ratios of the 17 listed nursing activities were examined using chi-square and Fisher's exact tests. All statistical analyses were performed using IBM SPSS Statistics 19 (IBM Corp., Armonk, NY, USA). All statistical tests were two-sided, and a p -value of <0.05 was considered significant.

Ethical considerations

A written document explaining the study objective, the preservation of anonymity and that participation was voluntary was included with the questionnaire. The receipt of a completed questionnaire was regarded as a respondent's consent to participate in the study. This study was conducted with the approval of the Ethics Committee of Kanazawa University (No.425).

RESULTS

There were 240 facilities from which both administrators and nurses responded (response rate 24.0%).

Facility, nurse and resident attributes

The attributes of facilities and residents are shown in Tables 1 and 2. Table 1 shows that nearly 75% of facilities were non-profit or social welfare corporations; over 90% provided medication management and outpatient medical consultation, and over 60% conducted home visits. Ambulance transportation had been used recently at more than 40% of facilities to deal with changes in residents' pathological conditions or accidents.

On average, facilities had 19.4 residents ($SD = 4.4$) and employed 1.5 nurses ($SD = 0.8$). Table 2 shows that close to 75% of residents were female and more than 75% were 80 years old or more. In total, 9.2% of group home residents in this study were classified as Long-term Care Grade 5, the highest level of care dependency.

The vast majority of nurses (82.8%) were 40 years old or more: 62 (26.1%) were aged 40–49 years; 91 (38.2%) aged 50–59 years and 44 (18.5%) aged 60–69 years. The mean number of years working in group homes was 2.9 years ($SD = 2.0$). Approximately two-thirds of nurses (66.2%, $n = 151$) worked full-time and 33.8% ($n = 77$) worked part-time.

Nursing activities

Table 3 shows the percentage of facilities using each of the 17 listed nursing activities. Four of these activities were carried out in more than 90% of facilities; these were, monitoring residents' health conditions, medication management, determining the necessity of medical consultation and dealing with changes in pathological condition. Nurses' reportage of activity usage was lower than 50% for four specific activities: providing support for residents with more severe medical needs, training of care workers, performing end-of-life care and arranging for overnight care service and changing services. Nurses determined the need for medical consultation at more than 90% of facilities and were engaged in emergency management and care at more than 70%.

Relationships amongst facility attributes, nurse attributes and nursing activities

Statistically significant associations were found between several facility attributes and their nursing activities. For example, a higher percentage of facilities owned by medical corporations had residents with greater medical needs (52.6%, $p = 0.017$) than other entities (31.8%), and a greater percentage of group homes without an attached medical facility reported dealing with changes in

Table 1 Basic characteristics and medical care provided by facilities

		Number of facilities [†]	%
Business entity	For-profit corporation	89	37.2
	Social welfare corporation	88	36.8
	Medical corporation	38	15.9
	Not-for-profit organization	18	7.5
	Incorporated association/foundation	3	1.3
	Others	3	1.3
Attached facilities (Multiple answers were possible)	Visiting care facility	64	26.7
	Group home	54	22.5
	Nursing care and welfare facility	23	9.6
	Hospital/clinic	22	9.2
	Residence with care services	20	8.3
	Nursing home for older people	20	8.3
	Home care support station	13	5.4
	Long-term Care Insurance-covered facility	12	5.0
	Others	38	15.8
	None	82	34.2
Medical care	Skin care	172	71.7
	Enema/faecal removal	126	52.5
	Pressure injury treatment	104	43.3
	Insulin administration	61	25.4
	Blood sugar management	57	23.8
	Bladder catheter placement	36	15.0
	Tube feeding	32	13.3
	Infusion therapy	17	7.1
	None	82	34.2
Medical consultation	Outpatient	232	96.7
	Home visit	152	63.3
Medications	Medication management	219	91.3
Emergency management implemented within the last 3 months	Ambulance transportation	97	40.4
	Unscheduled medical consultation	158	65.8
	Managed within the facility	57	23.8
Nurses' ages	20–29	5	2.0
	30–39	25	10.4
	40–49	62	25.8
	50–59	91	37.9
	60–69	44	18.3
	70–79	11	4.6
	Unclear	2	0.8
		Mean	SD
Number of residents		19.4	4.4
Number of nurses		1.5	0.8
Number of care workers		10.3	4.2
Number of day staff members		5.9	2.8
Number of night staff members		1.3	1.3

[†]N = 240.

Table 2 Demographic and clinical data of facility residents

		Number	%
Sex (<i>N</i> = 4625)	Male	1186	25.6
	Female	3439	74.4
Age (<i>N</i> = 4470)	Under 65	101	2.3
	65–69	115	2.6
	70–74	228	5.1
	75–79	612	13.7
	80–84	1072	24.0
	85–89	1337	29.9
	Over 89	1005	22.5
Long-term care grade [†] (<i>N</i> = 4570)	Assistance Grade 1	140	3.1
	Assistance Grade 2	257	5.6
	Care Grade 1	942	20.6
	Care Grade 2	1152	25.2
	Care Grade 3	1017	22.3
	Care Grade 4	641	14.0
Number of disease diagnosis per facility (<i>N</i> = 4470)	Care Grade 5	421	9.2
		Mean	SD
	Hypertension	8.6	4.4
	Heart disease	4.5	2.9
	Cerebrovascular disease	4.9	3.8
	Diabetes	2.8	1.9
	Dementia	13.0	5.7

[†]The long-term care grade is an index, based on the Long-term Care Insurance System, which indicates the level of need for care services. Those classified as care grades 1, 2 and 3 are permitted to use insurance-covered care services for 32–49, 50–69 and 70–89 min per day, respectively.³¹

pathological conditions (92.7% vs. 72.7%, $p = 0.008$) and advising care workers (88.5% vs. 72.7%, $p = 0.047$). The number of residents per facility was also related to several use ratios. Group homes with more residents conducted more nursing assessments (71.4% vs. 52.3%, $p = 0.005$), managed more emergencies outside service hours (58.7% vs. 42.3%, $p = 0.009$) and relied more on nursing judgment and care in emergencies (77.8% vs. 60.4%, $p = 0.003$).

Working patterns appeared to have a major influence on the use of nursing activities. Nurses who worked full-time made greater use of 12 of the listed activities than those who worked part-time ($p < 0.05$). Length of experience at medical institutions was not associated with nursing activities conducted in the previous month.

Patterns of nursing activities varied with resident and nurse characteristics; for example, associations between nursing activities and residents' diseases (Tables 4–6), and

between nursing activities and nurses' years of experience at the facility (Table 7) revealed different patterns. Nurses at facilities that had more residents with heart disease tended to be engaged in more medication management and care-dependency prevention activities whereas nurses at facilities that had more residents with dementia were more likely to be engaged in determination of medical consultation need and handling changes in residents' pathological condition. More experienced nurses tended to carry out particular activities to a greater extent, such as emergency management outside service hours and end-of-life care.

Relationships among nurses' GSES scores, activities and attributes

The mean GSES score was 58.5 (SD = 11.7). GSES scores were not statistically significantly associated with years of experience or working conditions but were linked with

Table 3 Number and percentage of facilities conducting specific nursing activities

No.	Nursing activity	Number of facilities [†] %	
1	Monitoring users' health conditions (observation, vital sign measurement)	234	97.5
2	Medication management (e.g. dose and schedule adjustments)	221	92.1
3	Determining the necessity of medical consultation	218	90.8
4	Dealing with changes in the pathological condition	218	90.8
5	Advising care workers from a nursing viewpoint	209	87.1
6	Providing care workers with daily guidance on residents' health	203	84.6
7	Providing daily medical care	185	77.1
8	Nursing judgment and care in emergencies (e.g. when accidents or falls occur)	168	70.0
9	Cooperating with other medical professionals (e.g. doctors and care managers)	158	65.8
10	Conducting assessment from a nursing viewpoint	149	62.1
11	Supporting residents' families in relation to medical care	135	56.3
12	Performing care-dependency prevention measures (e.g. rehabilitation and accident prevention)	129	53.8
13	Managing emergencies outside of service hours	122	50.8
14	Providing support for residents with higher medical needs	85	35.4
15	Training care workers	82	34.2
16	Arranging for overnight care service use and changing services	64	26.7
17	Performing end-of-life care	41	17.1

[†]N = 240.

Table 4 Relationships between nursing activities and high/low numbers of residents with heart disease per facility[‡]

Nursing activities		Number of facilities with high/low number of residents with heart disease					
		Low		High		χ^2	$p^{\dagger}$
Medication management (e.g. dosage adjustment and distribution)	Using	120	88.9%	95	100%	11.292	<0.001
	Not using	15	11.1%	0	0%		
Performing care-dependency prevention measures (e.g. rehabilitation and accident prevention)	Using	61	45.2%	62	65.3%	9.035	0.003
	Not using	74	54.8%	33	34.7%		
Providing support for residents with higher medical needs	Using	40	29.6%	43	45.3%	5.909	0.018
	Not using	95	70.4%	52	54.7%		
Training care workers	Using	37	27.4%	41	43.2%	6.172	0.013
	Not using	98	72.6%	54	56.8%		
Arranging for overnight care service use and changing service contents	Using	27	20.1%	35	36.8%	7.845	0.005
	Not using	107	79.9%	60	63.2%		

[‡]Based on chi-square tests; values for each item categorized into two groups based on the mean.

[†] p for results of chi-square tests.

Note: Only those values that were significantly different are included.

several nursing activities. Low self-efficacy scores were more often seen in those who performed (rather than did not perform) care-dependency prevention measures and supported residents with higher medical needs (Table 7).

DISCUSSION

We found that nurses in multifunctional small group homes had an important role in managing residents' health, determining the need for medical consultation and dealing with

Table 5 Relationships between nursing activities and high/low numbers of residents with diabetes per facility[‡]

		Number of facilities with high/low number of residents with diabetes				χ^2	$p^{\dagger}$
		Low		High			
Nursing activities							
Medication management (e.g. dosage adjustment and distribution)	Using	2	10.5%	82	37.6%	5.605	0.022
	Not using	17	89.5%	136	62.4%		

[‡]Based on chi-square tests; values for each item categorized into two groups based on the mean.

[†] p for results of chi-square tests.

Note: Only those values that were significantly different are included.

Table 6 Relationships between nursing activities and high/low numbers of residents with dementia per facility[‡]

		Number of facilities with high/low number of residents with dementia					
		Low		High		χ^2	$p^{\dagger}$
Nursing activities							
Determining the necessity for medical consultation	Using	98	86.7%	120	94.5%	4.327	0.045
	Not using	15	13.3%	7	5.5%		
Dealing with changes in the pathological condition	Using	98	86.7%	120	94.5%	4.327	0.045
	Not using	15	13.3%	7	5.5%		
Providing daily medical care	Using	78	69.0%	107	84.3%	7.847	0.006
	Not using	35	31.0%	20	15.7%		
Cooperating with other medical professionals (e.g. doctors and care managers)	Using	66	58.4%	92	72.4%	5.236	0.029
	Not using	47	41.6%	35	27.6%		
Conducting assessment from a nursing viewpoint	Using	62	54.9%	87	68.5%	4.724	0.033
	Not using	51	45.1%	40	31.5%		
Providing support for residents with higher medical needs	Using	30	26.5%	55	43.3%	7.342	0.007
	Not using	83	73.5%	72	56.7%		

[‡]Based on chi-square tests; values for each item categorized into two groups based on the mean.

[†] p for results of chi-square tests.

Note: Only those values that were significantly different are included.

emergencies as well as making arrangements for flexible care services. We examined the relationships among characteristics of the facilities, residents and nursing activities. As might be expected, we found different patterns of disease associated with different nursing activities.

Nursing activities have a number of functions in multifunctional small group homes. In a large number of facilities, nurses determine the need for medical consultation and manage emergency care. The majority of residents in this study required regular medical consultation and medication management. More than 40% of facilities used

ambulance transportation because of residents' medical needs. These activities suggest that residents have a need for high levels of care. Previous studies have found that the use of at-home or community-based care services does not directly improve residents' functional status,¹⁹ but does prevent hospitalization.²⁰ Another study in the United States reported that the use of home health services delayed nursing home admission.²¹ The large number of nurses in this study who determined need for medical consultations and who were engaged in emergency management and care suggests the potential that these nursing activities might

Table 7 Relationships between nursing activities, nurses' years of experience and nurses' GSES score[‡]

Nursing activities		Nurse attributes							
		Years of experience working in facilities				GSES score			
		Short Nurses	Long Nurses	χ^2	$p^\dagger$	Low Nurses	High Nurses	χ^2	$p^\dagger$
Performing care-dependency prevention measures (e.g. rehabilitation and accident prevention)	Using					65 60.7%	49 45.0%	5.404	0.021
	Not using					42 39.3%	60 55.0%		
Managing emergencies outside of service hours	Using	55 44.4%	67 58.8%	4.941	0.028				
	Not using	69 55.6%	47 41.2%						
Providing support for residents with higher medical needs	Using					47 43.9%	30 27.5%	6.332	0.016
	Not using					60 56.1%	79 72.5%		
Performing end-of-life care	Using	15 12.1%	26 22.8%	4.778	0.039				
	Not using	109 87.9%	88 77.2%						

GSES, General Self-Efficacy Scale.

[‡]Based on chi-square tests with values for each item classified into two groups based on the mean.

[†] p for results of chi-square tests.

Note: Only those values that were significantly different are included.

reduce unnecessary hospitalization and improve older people's ability to continue to live in the community. In 25% of facilities, care services were changed based on the judgment of nurses. Group homes are characterized by flexible care services and the opinions of nurses might contribute to appropriate service use, which confirms the importance of having nurses at such facilities.

Previous research has also shown that care coordination managed by registered nurses can influence utilization and cost outcomes, and can affect the health and functional ability of frail older people.²² Active coordination by nurses in group homes is likely to improve the provision of individually tailored services. For example, in facilities with a higher percentage of heart disease, nursing activities such as medication management and care-dependency prevention were more likely to be used. These services are particularly important for these residents, and our results indicated that nurses tailored their activities to individual resident characteristics. Nurses are often the only qualified healthcare professionals on site. Residents in group homes are often vulnerable older people with multiple diseases, so nurses have an important role in coordinating and managing their care both on a daily basis and during an emergency.

This study showed the importance of nursing activity to enable older people with severe health care needs, such as

the residents of our study facilities, to continue living at home. We also identified that nursing activities can be disease specific. Multifunctional small group homes are unique service providers offering flexible, individual, personalized care to comprehensively support older people so they might continue to live within the community. Our results showed that nurses provided a wide range of care within these services.

We examined relationships between the working conditions of nurses and their activities and found that nurses were more likely to use the listed nursing activities when they worked full-time. This might be explained by the differences in treatment and working hours between full-time and part-time employees in Japan. Similar findings have been reported in Korea's community-based visiting care systems, where the number of visiting nurses at care facilities and their rehire rates have been associated with residents' quality of life.²³ Our study found that emergency management outside service hours was more likely to be used in facilities with more residents. It might therefore be effective to increase the total number of nurses or the proportion working full-time in such facilities. More experienced nurses used certain activities more actively, such as emergency management outside service hours and end-of-life care. It might therefore be necessary to develop

measures to help nurses continue to work in the same facility, to help them develop further experience.

We found that the nurses in this study who more actively adopted care-dependency prevention measures and provided support for residents with severe medical needs had lower GSES scores, indicating lower self-efficacy, or belief in their abilities. In Japan, aspiring nurses generally pursue a 3 year course of basic nursing education following their high school graduation, and they become certified nurses after passing the national certification examination for nursing students.²⁴ Passing this examination is the only necessary qualification for nurses wishing to work in multifunctional small group homes. This perhaps might explain, in part at least, low self-efficacy in carrying out these more specialized nursing activities.

At the time of the study, a number of other factors might also have been influential. First, at the time of this study (2013), group home services had only been in existence for 7 years, and the nurses in this study had only a mean experience of 2.9 years working in multifunctional small group homes. Second, guidelines specifying the roles of nurses working in these facilities had not yet been established, and group home nurses might have been performing particular nursing activities without clear guidelines, resulting in poor self-efficacy. Third, the mean number of nurses per group home was only 1.5, which might mean nurses cannot consult with colleagues about nursing judgments. To improve the quality of nursing activities, nurses' self-efficacy should be enhanced through training opportunities, by developing nursing activity guidelines and enabling nurses to discuss and consult with colleagues on care issues.

In PACE programs, the professionals involved cooperate with each other from the primary to acute phases, to provide flexible and personalized support for older people who meet the criteria for nursing home service use and to help them continue to live in the community.²⁵ PACE is a comprehensive program covering both medical and care services, and enables residents to access appropriate care services even when their pathological condition changes. In contrast, Japanese multifunctional small group homes are social welfare facilities for older people, and nurses are the only medical professionals available, highlighting the importance of their judgments and the heavy burden placed on them during emergencies. It is therefore necessary to strengthen cooperation between staff in group homes and other healthcare professionals and educate care workers to help them provide team-based healthcare

support for residents. It is also important to consider measures to reduce the burden on nurses.

We found that 17.1% of facilities in this study provided end-of-life care. This might be important when considering future directions for nursing practice in multifunctional small group homes. The need for education about pain/symptom control has been found to vary between professionals and facilities, even among group homes for older people with dementia.²⁶ Nurses, however, are present at 60% of deaths.²⁷ Because many Japanese people wish to die at home,²⁸ end-of-life care is likely to become more important in group homes, as the number of older residents increases. Nurses are likely to have a central part in end-of-life care and education about pain/symptom control for care workers. Group homes have a role in confirming the wishes of each resident regarding end-of-life care and helping residents to prepare before their health condition worsens.

Although there have been few studies in Japan on the effects of nursing activities delivered to older individuals living at home, there have been several studies conducted elsewhere. Community-based visiting care systems were found to improve quality of life among older people living at home in South Korea,²³ and a study conducted in Hong Kong confirmed that visits by community nurses (mainly for surveillance) enhanced the self-reported health status of older people with chronic cardiovascular disease after they were discharged from hospital.²⁹ A study in Taiwan found that case management by public health nurses decreased the blood sugar levels of older community residents with diabetes.³⁰ These findings suggest that it might be possible to improve the health of Japanese group home residents by promoting effective nursing activities.

Limitations

This study has a number of limitations. For example, it might be difficult to generalize our results because of the low response rate (24.0%). This was partly because we limited study subjects to those working in facilities from which we received completed questionnaires from both administrators and nurses. We could not determine causal relationships between nursing activities and study factors because we used a cross-sectional design in this study.

CONCLUSION

This is the first study to identify the status of nursing activities provided in multifunctional small group homes of Japan. The results of this study suggest that a number of

nursing activities, including health management, determining the need for medical consultation and managing emergencies, are used with the aim of supporting older people with high health care needs to continue living at home. We also identified nursing activities that were disease specific, providing us with useful suggestions for offering home-based nursing care to older people. In Japan's super-ageing society, these facilities will become increasingly important as service providers to support older people living at home. They might therefore provide a useful model for examining measures to support older people in other countries with rapidly ageing populations. To promote the activities of nurses, who have a central role in this service, it is important to increase the number of nurses in facilities with a large number of residents and to examine their work patterns. Our results also indicate the importance of offering training opportunities to increase nurses' self-efficacy and promote their activities, as well as to develop nursing activity guidelines. Further research is also required: qualitative studies to explore in more detail what determines use of different nursing activities; development of interventions to enhance the self-efficacy of nurses working in group homes; further examination of relationships between nurses' GSES scores and the nursing activities they perform.

ACKNOWLEDGMENTS

The authors would like to thank the group home administrators and nurses who took part in the study. This study was supported by a Grant-in-Aid for Scientific Research (JSPS KAKENHI Grant Number JP23792723, JP.26463496). The research was conducted as part of a doctoral dissertation at Kanazawa University, School of Health Sciences.

REFERENCES

- Segelman M, Szydowski J, Kinosian B *et al.* Hospitalizations in the program of all-inclusive care for the elderly. *Journal of the American Geriatric Society* 2014; **62**: 320–324. DOI: 10.1111/jgs.12637.
- Meret-Hanke LA. Effects of the program of all-inclusive care for the elderly on hospital use. *Gerontologist* 2011; **51**: 774–785. DOI: 10.1093/geront/gnr040.
- Fretwell MD, Old JS, Zwan K, Simhadri K. The elderhaus program of all-inclusive care for the elderly in North Carolina: improving functional outcomes and reducing cost of care: preliminary data. *Journal of the American Geriatric Society* 2015; **63**: 578–583. DOI: 10.1111/jgs.13249.
- Rantz M, Popejoy LL, Galambos C *et al.* The continued success of registered nurse care coordination in a state evaluation of aging in place in senior housing. *Nursing Outlook* 2014; **62**: 237–246. DOI: 10.1016/j.outlook.2014.02.005.
- Rantz MJ, Porter RT, Cheshier D *et al.* TigerPlace, a state-academic-private project to revolutionize traditional long-term care. *Journal of housing for the elderly* 2008; **22**: 66–85.
- Perry M, Melis RJ, Teerenstra S *et al.* An in-home geriatric programme for vulnerable community-dwelling older people improves the detection of dementia in primary care. *International journal of geriatric psychiatry* 2008; **23**: 1312–1319. DOI: 10.1002/gps.2128.
- Stijnen MM, Jansen MW, Duimel-Peeters IG, Vrijhoef HJ. Nurse-led home visitation programme to improve health-related quality of life and reduce disability among potentially frail community-dwelling older people in general practice: a theory-based process evaluation. *BMC family practice* 2014; **15**: 173. DOI: 10.1186/s12875-014-0173-x.
- Campbell JC, Ikegami N. Long-term care insurance comes to Japan. *Health Affairs* 2000; **19**: 26–39. DOI: 10.1377/hlthaff.19.3.26.
- Ministry of Health, Labour and Welfare. Act for partial revision of the Long-Term Care Insurance Act, etc., in order to strengthen long-term care service infrastructure 2011. Available from URL: http://www.mhlw.go.jp/english/policy/care-welfare/care-welfare-elderly/dl/en_tp01.pdf. Accessed 25 August 2016.
- Cabinet Office. Number of households with co-resident persons aged 65 and over, their percentage distribution (by household type) and their share in all households, Section 2 Existing State and Trends of Elderly People and their Environment, Annual Report on the Aging Society: 2013 (Summary). Available from URL: <http://www8.cao.go.jp/kourei/english/annualreport/2013/pdf/1-2-1.pdf>. Accessed 25 August 2016.
- Verbeek H, Zwakhalen SM, van Rossum E, Ambergen T, Kempen GI, Hamers JP. Small-scale, homelike facilities versus regular psychogeriatric nursing home wards: a cross-sectional study into residents' characteristics. *BMC health services research* 2010; **10**: 30. DOI: 10.1186/1472-6963-10-30.
- Kane RA, Lum TY, Cutler LJ, Degenholtz HB, Yu TC. Resident outcomes in small-house nursing homes: a longitudinal evaluation of the initial green house program. *Journal of the American Geriatrics Society* 2007; **55**: 832–9.
- Makino Y. Role of nurses and care workers in small-scale multi-functional care. *Nihon kango fukushi gakkaiishi [Journal of Japanese Society for the Study of Nursing and Social Work]* 2010; **15**: 81–97. (in Japanese)
- Welfare and Medical Service Agency, Welfare and Medical Service Network System(WAMNET). Available from URL: <http://www.wam.go.jp/content/wamnet/pcpub/top/>. (in Japanese) Accessed 25 August 2016.
- Katahira N, Honda A, Ueno M, Kitaoka H, Hirokawa S, Watanabe T, Takeda Y, Hatanaka T, Bessho Y. Current

- nursing systems in group homes for elderly with dementia—focusing on use of visiting nurse services. *Journal of Japan Academy of Community Health Nursing* 2009; **12**: 59–64. (Abstract in English)
- 16 Sherer M, Maddux JE, Mercandante B, Prentice-Dunn S, Jacobs BW, Rogers RW. The self-efficacy scale: construction and validation. *Psychological Reports* 1982; **51**: 663–671. DOI: 10.2466/pr0.1982.51.2.663.
 - 17 Narita K, Shimoyama Y, Nakazato K, Kawaai C, Sato S, Osada YA; Japanese version of the generalized self-efficacy scale: Scale utility from the life-span perspective. *Japanese Journal of Educational Psychology* 1995; **43**: 306–314. (in Japanese)
 - 18 Bandura A. *Self-Efficacy the Exercise of Control*. New York: W. H. Freeman, 1997.
 - 19 Olivares-Tirado P, Tamiya N, Kashiwagi M. Effect of in-home and community-based services on the functional status of elderly in the long-term care insurance system in Japan. *BMC Health Services Research* 2012; **12**: 239. DOI: 10.1186/1472-6963-12-239.
 - 20 Tomita N, Yoshimura K, Ikegami N. Impact of home and community-based services on hospitalisation and institutionalisation among individuals eligible for long-term care insurance in Japan. *BMC Health Services Research* 2010; **10**: 345. doi: 10.1186/1472-6963-10-345.
 - 21 Young Y, Kalamaras J, Kelly L, Hornick D, Yucel R. Is Aging in Place Delaying Nursing Home Admission? *Journal of the American Medical Directors Association* 2015; **16**: 900. e1–6. DOI: 10.1016/j.jamda.2015.07.017.
 - 22 Popejoy LL, Galambos C, Stetzer F *et al.* Comparing Aging in Place to Home Health Care: Impact of Nurse Care Coordination On Utilization and Costs. *Nursing economic\$* 2015; **33**: 306–313.
 - 23 Chow SKY, Wong FK, Chan TM, Chung LY, Chang KK, Lee RP. Community nursing services for postdischarge chronically ill patients. *Journal of Clinical Nursing* 2008; **17**: 260–271. DOI: 10.1111/j.1365-2702.2007.02231.x.
 - 24 Japanese Nursing Association. Basic nursing education. Available from URL: <http://www.nurse.or.jp/jna/english/nursing/education.html#basic>. Accessed 25 August 2016.
 - 25 Mukamel DB, Peterson DR, Temkin-Greener H *et al.* Program characteristics and enrollees' outcomes in the Program of All-Inclusive care for the Elderly (PACE). *The Milbank Quarterly* 2007; **85**: 499–531. DOI: 10.1111/j.1468-0009.2007.00497.x.
 - 26 Hirakawa Y, Kuzuya M, Uemura K. Opinion survey of nursing or caring staff at long-term care facilities about end-of-life care provision and staff education. *Archives of Gerontology and Geriatrics* 2009; **49**: 43–48. DOI: 10.1016/j.
 - 27 Nakanishi M, Honda T. Processes of decision making and end-of-life care for patients with dementia in group homes in Japan. *Archives of Gerontology and Geriatrics* 2009; **48**: 296–9. DOI: 10.1016/j.
 - 28 Fukui S, Yoshiuchi K, Fujita J, Sawai M, Watanabe M. Japanese people's preference for place of end-of-life care and death: a population-based nationwide survey. *Journal of pain and symptom management* 2011; **42**: 882–892. DOI: 10.1016/j.jpainsymman.2011.02.024.
 - 29 Lim JY, Kim GM, Kim EJ, Choi KW, Kim SS. The effects of community-based visiting care on the quality of life. *Western Journal of Nursing Research* 2013; **35**: 1280–1291. DOI: 10.1177/0193945913490237.
 - 30 Lu KY, Lin PL, Tzeng LC, Huang KY, Chang LC. Effectiveness of case management for community elderly with hypertension, diabetes mellitus, and hypercholesterolemia in Taiwan: a record review. *International Journal of Nursing Studies* 2006; **43**: 1001–1010.
 - 31 Labour and Welfare. Youkaigo nintei wa donoyouni okonawareruka.[The Long-term Care Grading System], 2008. Available from URL: <http://www.mhlw.go.jp/topics/kaigo/nintei/gaiyo2.html>. (in Japanese) Accessed 25 August 2016.