Age and sex differences in controlled force exertion measured by a computing bar chart target-pursuit system

著者	Nagasawa Yoshinori, Demura Shinichi
journal or publication title	Measurement in Physical Education and Exercise Science
volume	13
number	3
page range	140-150
year	2009-07-01
URL	http://hdl.handle.net/2297/19136

doi: 10.1080/10913670902812804

Title Page

Original Article

Article title:

Age and sex differences in controlled force exertion measured by a computing bar chart target-pursuit system

Author names:

Yoshinori NAGASAWA¹⁾

Shinichi DEMURA²⁾

Affiliations:

- 1) Kyoto Pharmaceutical University, Department of Health and Sports Sciences,
5 Nakauchi-cho, Misasagi, Yamashina-ku, Kyoto, Kyoto 607-8414 Japan
- 2) Kanazawa University, Graduate School of Natural Science & Technology,
Division of Life Science, Kakuma, Kanazawa, Ishikawa 920-1164 Japan

Running head:

Age and sex differences: controlled force exertion

Address correspondence to

Yoshinori Nagasawa,
Department of Health and Sports Sciences,
Kyoto Pharmaceutical University,
5 Nakauchi-cho, Misasagi, Yamashina-ku, Kyoto, Kyoto 607-8414 Japan.

+81-75-595-4682 (Tel)

+81-75-595-4682 (Fax)

ynaga@mb.kyoto-phu.ac.jp (e-mail)

1 **Title Page**

2

3 Original Article

4

5 **Article title:**

6 Age and sex differences in controlled force exertion measured by a computing bar

7 chart target-pursuit system

8

9

10 **Running head:**

11 Age and sex differences: controlled force exertion

12

13

14

15

16

17

18

19

20

21

22

23

24

1 **Abstract**

2 This study aimed to examine the age and sex differences in controlled force
3 exertion measured by the bar chart display in 207 males (Age 42.1 ± 19.8 yrs) and
4 249 females (Age 41.7 ± 19.1 yrs) aged 15 to 86 years. The subjects matched their
5 submaximal grip strength to changing demand values, which appeared as a moving
6 bar chart on the display of a personal computer. The subjects performed the
7 controlled force exertion test using the dominant hand 3 times with 1-min intervals
8 (one trial was 40 sec.) after one practice trial. A total of the differences between the
9 demand value and the grip exertion value for 25 sec. was used as the evaluation
10 parameter. The errors in controlled force exertion showed a right-skewed
11 distribution in both sexes but showed a normal distribution after logarithmic
12 transformation (males: $W=0.06$, females: $W=0.74$ $p>0.05$). In addition, the errors in
13 controlled force exertion tended to increase constantly with age in both sexes.
14 Significant linear regressions were identified ($r^2_{\text{males}}=0.88$, $r^2_{\text{females}}=0.81$), but there
15 was no significant difference in the increase rate of both sexes. The results of the
16 analysis of variance and multiple comparisons showed insignificant sex differences
17 among means, except for those in individuals older than 70 years, and significant
18 differences between means in age groups older than 40 years and the 20 year old age
19 group were found in both sexes. Individual differences were almost the same in both
20 sexes ($CV_{\text{males}}=20.0 \sim 34.8$, $CV_{\text{females}}=17.7 \sim 36.2$). Errors in controlled force exertion
21 showed a nonsignificant sex difference and increased gradually with age in both
22 sexes but increased remarkably with age after 40 years of age.

23 **Key words:** humans, adult, hand strength, psychomotor performance

24

25

1 **Text**

2

3 **Introduction**

4 Nervous and muscle functions work closely together to control motor
5 performance in humans. Because it is rare to exert maximal ability during daily
6 activities, it will be very important to determine whether submaximal ability is
7 exerted efficiently or continually (Halaney & Carey, 1989). In elderly and
8 developmentally delayed individuals, this is particularly important. In these
9 individuals, it is essential to estimate the voluntary movement functions that
10 primarily contribute to skillful and efficient submaximal movements (Henatsch &
11 Langer, 1985), because the exertion of maximal ability involves risks. Local
12 movements which demand feedback information, such as hand-foot movements,
13 hand-eye coordination and so on, are closely involved in the coordination of the
14 voluntary movement system, i.e. controlled force exertion (Henatsch & Langer, 1985).
15 The controlled force exertion test is one of the useful tests to evaluate motor control
16 function, which acts to coordinate force exertion according to the demands of each
17 task. To smoothly exert motor control function, information from the central and
18 peripheral nervous systems is integrated in the cerebrum, which functions to
19 properly control movements in each motor organ. Motor control function is
20 interpreted to be superior when contraction and relaxation of muscles are smoothly
21 performed according to the movement of a target. In this situation, variability
22 decreases and accuracy increases (Brown & Bennett, 2002). The ability to control
23 this exertion of motor function is postnatally acquired through learning based on
24 motor experiences.

25 Nagasawa and Demura (2002) studied the tracking movement in submaximal

1 strength exertion and developed a new test for rationally and objectively estimating
2 grading, spacing, and timing, which are important elements of controlled force
3 exertion. Their method involves using a grip dynamometer coupled with a personal
4 computer. It was reported that the new test has a high reliability (Nagasawa,
5 Demura, & Nakada, 2003) and that it measures the control of force rather than
6 position tracking than that measured by the pursuit-rotor and pegboard tests
7 (Nagasawa, Demura, & Kitabayashi, 2004). Hence, it is useful as a test to evaluate
8 the neuromuscular function of the elderly (Nagasawa, Demura, Yamaji, Kobayashi,
9 & Matsuzawa, 2000). On the other hand, factors such as fatigue, training, age
10 (growth and development), etc. influence controlled force exertion (Yamamoto, 1983).
11 It is known that physical fitness (neuromuscular function) generally decreases with
12 age, and individual differences are large in the elderly (Bemben, Massey, Bemben,
13 Misner, & Boileau, 1991).

14 Ranganathan, Siemionow, Sabgal, and Yue (2001) examined effects of aging on
15 hand function, and reported that, compared with younger subjects, elderly subjects
16 have weaker handgrip and maximum pinch force, and decreased ability to maintain
17 steady submaximal pinch force. They reported that the decrease in the ability to
18 maintain steady submaximal pinch force is more pronounced in females than males.
19 Voelcker-Rehage and Alberts (2005) reported that younger subjects perform the
20 variable force tracking task at a higher level than elderly subjects. Nagasawa et al.
21 (2000) examined the characteristics of controlled force exertion by the bar chart
22 display in 60 healthy older people (30 males, 30 females) aged 65 to 78 years and
23 compared their performances with those of 60 healthy university students (30 males,
24 30 females). They reported that the elderly had inferior controlled force exertion to
25 the young adults in both sexes. Furthermore, they found that elderly females were

1 inferior to elderly males and that large individual differences were present. However,
2 the above results on age group differences were examined based on a small sample.
3 There are few reports on the change (decrease) of controlled force exertion with age
4 or on the sex and individual differences of these decreases based on a large sample.
5 Nagasawa et al. (2000) indicates that a decrease in the ability to exert controlled
6 force will result in an increase in controlled force exertion errors. We hypothesized
7 that the errors in controlled force exertion would increase gradually with age, and
8 this tendency to increase would differ between the sexes, with sex and individual
9 differences existing in the elderly subjects.

10 This study aimed to examine the age-level and sex differences in controlled force
11 exertion and the above-stated hypotheses.

12

13 **Method**

14 *Subjects*

15 The subjects were 207 males (age 42.1 ± 19.8 yrs, height 168.6 ± 7.2 cm, weight
16 65.8 ± 9.6 kg) and 249 females (age 41.7 ± 19.1 yrs, height 156.3 ± 6.4 cm, weight 53.0
17 ± 6.9 kg) aged 15 to 86 years. Their physical characteristics are summarized by age
18 group in Table 1. All were regarded as right-handed, based on the Oldfield's
19 inventory (1971). Height and weight were similar to Japanese normative values
20 (Laboratory Physical Education in Tokyo Metropolitan University, 1989) for each
21 age-level in both sexes. There were no significant sex differences in the means of age
22 in all age groups. The males had significantly greater mean maximal grip strength
23 and standing height than the females in all age groups. The males had significantly
24 greater mean weight than the females in all age groups except for 70 years or older.
25 Significant correlations were not found among height, weight, age, or controlled force

1 exertion in both sexes, except for the weight of the 15-19yr. old female group
2 ($r^2=0.281$) and the height of the 60-69yr. old female group ($r^2=0.123$). Therefore, we
3 judged that the influences of the above-stated factors on controlled force exertion
4 could be neglected, and we did not control each variable by physique in a comparison
5 of the measurements for these two groups. No subject reported previous wrist
6 injuries or upper limb nerve damage, and all were in good health. Prior to enrollment,
7 the purpose and procedure of this study were explained in detail. This protocol was
8 approved by the Institutional Review Board, and informed written consent was
9 obtained from all subjects. No subject previously experienced a controlled force
10 exertion test. Neuromuscular function generally peaks, with the majority of changes
11 occurring during the period from the late teens to twenties, and it then gradually
12 decreases with age after the age of 30 (Bemben, et al., 1991). The subjects were
13 grouped based on age as follows: 15-19 (27 males, 27 females), 20-24 (29 males, 38
14 females), 25-29 (25 males, 27 females), 30-39 (25 males, 41 females), 40-49 (25 males,
15 27 females), 50-59 (23 males, 26 females), 60-69 (27 males, 36 females), and 70 and
16 older (26 males, 27 females).

17

18 *****Table 1 near here*****

19

20 *Test and Test Procedure*

21 In this study, the subjects performed a grip exertion, attempting to minimize the
22 differences between a demand value and the value of the grip strength as presented
23 on a computer display. This information was transmitted at a sampling rate of 10 Hz
24 to a computer through an RS-232C data output cable after A/D conversion.
25 Measurements of grip strength and controlled force exertion were taken with a

1 Smedley's type handgrip mechanical dynamometer (GRIP-D5101; Takei, Tokyo,
2 Japan), with an accuracy of $\pm 2\%$ in the range of 0 to 979.7 N.

3 Based on a preliminary investigation (Nagasawa & Demura, 2002), a bar chart
4 on the display screen was used. The display showed both the demand value and the
5 actual grip strength simultaneously. The demand value increased and decreased at a
6 constant frequency (range=5 to 25% of maximal grip strength). Changes in the actual
7 grip-exertion value were displayed as vertical changes in that bar, as with the
8 demand value. The demand values varied over a period of 40 sec. at a frequency of
9 0.3 Hz. This rate of change is most easily imitated by the neuromuscular function
10 (Hayashi, 1967; Meshizuka & Nagata, 1972). Figure 1 shows the bar chart displays.
11 Details of the apparatus to measure the controlled force exertion have been described
12 (Nagasawa & Demura, 2002). Sufficient rest time was given to eliminate the
13 influence among the tests and the subjects' fatigue (Nagasawa & Demura, 2002).
14 Subjects wore glasses when required and sat at appropriate distances from the display. They
15 tracked the demand values in the displays, and then measurements were performed.
16 Measurements were not affected by poor vision or fatigue. Subjects in a preliminary
17 experiment were capable of tracking the demand values in the displays.

18

19 *****Figure 1 near here*****

20

21 Relative demand values, not absolute demand values, were utilized, since
22 physical fitness and the muscular strength of each individual are different. The
23 relative demand value varied around 5 to 25% of maximal grip strength. The relative
24 demand value was exactly altered to present the same shape of demand function to
25 all subjects, despite the differences in the scale range (grip strength) observed among

1 subjects. The software program was designed to present the relative demand values
2 within a constant range on the display, regardless of whether maximal grip-strength
3 values were large or small. The demand values in this study used bar chart targets
4 which varied cyclically (see Figure 1).

5 The size of the grip was set so that the subject felt comfortable squeezing the
6 grip. The subject performed maximal grip exertion with the dominant hand twice at
7 1-min. intervals, and the greater value was taken as the value of maximal grip
8 strength (Nagasawa, et al., 2000; Nagasawa & Demura, 2002). The test of controlled
9 force exertion was performed in three trials at 1-min. intervals after one practice
10 trial. The test of controlled force exertion was similar to a commonly used test of grip
11 strength (Walamies & Turjanmaa, 1993; Skelton, Greig, Davies, & Young, 1994),
12 except for the exertion of prolonged submaximal grip. The subject stood upright with
13 the wrist in the neutral position between flexion and extension and the elbow
14 straight and close to the body when exerting the grip. The duration of each trial was
15 40 sec., and the controlled force exertion was estimated using the data from three
16 trials, excluding the first 15 sec. of each trial, according to the previous study of
17 Nagasawa et al. (2000). The total sum of the percent of differences between the
18 demand value and the grip strength was used as an estimate of controlled force
19 exertion (Demura & Nagasawa, 2002). Smaller differences were interpreted to be a
20 better ability to control force exertion. Each subject was free to adopt a standing
21 position most conducive to a clear view of the display (Demura & Nagasawa, 2002).
22 Of three trials, the mean of the second and the third trials was used for analysis
23 (Nagasawa, et al., 2004).

24

1 *Statistical analysis*

2 Data were analyzed using SPSS (Version 11.5 for Windows). The characteristics
3 of the distribution were evaluated for coefficients of skew, kurtosis, and normality
4 (goodness of fit test: Shapiro-Wilk's W -test) in both the sum total and age groups. To
5 examine the variance of the measurements with age, linear regression coefficients
6 were computed for both males and females, and then the difference was examined. To
7 examine significant differences among the means of the age groups (8 x 2 matrix:
8 age-grade x sex group), two-way analysis of variance (ANOVA) was used after
9 logarithmic based 10 transformation. When a significant main effect was found, a
10 multiple-comparison test was done using a Tukey's Honestly Significant Difference
11 (HSD) method for pair-wise comparisons. In addition, the size of mean differences
12 (effect size) between trials of those in their 20-24yr. old group and each other age
13 group were examined. Coefficients of variance were calculated to examine individual
14 differences between age groups. Results are presented as mean and standard
15 deviations unless otherwise specified. An alpha level of 0.05 was taken to be
16 significant for all tests.

17

18 **Results**

19 Table 2 shows the distribution characteristics of each age group for the
20 controlled force exertion values by sex. Skew values of each age group were all
21 positive values, except for the 20-24yr. old group (0.0) in males. The measurements
22 also showed a right-skewed distribution in both sexes. Normality cannot be assumed
23 in both sexes (males: $W=0.87$, females: $W=0.79$, $p<0.05$), but the measurements in
24 both sexes showed a normal distribution after logarithmic transformation (males:
25 $W=0.06$, females: $W=0.74$, $p>0.05$).

1

2

Table 2 near here

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

Table 3 near here

23

Figure 2 near here

24

25

The coefficient of variance was in the same range for all age groups in both sexes

1 (CV_{males}=20.0~34.8, CV_{females}=17.7~36.2), but showed a high value for those older
2 than 40 years of age in males and for those older than 60 years of age in females. The
3 effect size of differences between the mean of 20-24yr. old and the means of age
4 groups older than 40 years of age showed large values over 0.8 in both sexes (Table
5 3).

6

7 **Discussion**

8 The mean value of controlled force exertion increased at an almost constant rate
9 in males and females with age, and their increase rate hardly showed a difference
10 between the sexes. In addition, the remarkable differences in both sexes were not
11 found in all age groups, except for the 70 years and older group. The functional role
12 related to movement performances may differ based on the region of the nervous
13 system controlling it, i.e. the cerebellum is generally associated with skilled motor
14 behavior, and the basal ganglia, in particular the striatonigral system, is associated
15 with actual motor behavior (Kornhuber, 1974). Bemben, Massey, Bemben, Misner, &
16 Boileau (1996) reported that the elderly show a noticeable decrease in peripheral
17 muscle activity compared with young people, based on measurements of muscular
18 endurance using intermittent grip strength. From reports by many researchers
19 (Rikli & Busch, 1986; Rikli & Edwards, 1991; Welford, 1988), including Dustman,
20 Ruhling, Russell, Shearer, Bonekat, Shigeoka, Wood, & Bradford (1984), it has been
21 clarified that the reaction time of muscles decreases with age. Controlled force
22 exertion was confirmed to decrease after 40 years of age in both sexes. The present
23 test was performed by submaximal muscular exertion with a moderate cycle (0.3 Hz)
24 of changing demand values. The achievement of this test requires strong hand-eye
25 coordination in feedback such as 'sense of force exertion', 'matching of target', and so

1 forth (see method). The decrease in muscular strength is based on changes of
2 neuromuscular pathways and muscle fiber composition, spinal motor neuron
3 apoptosis (Galganski, Fuglevand, & Enoka, 1993) and by muscle atrophy with age
4 (Cauley, Petrini, LaPorte, Sandler, Bayles, Robertson, & Slemenda, 1987). Therefore,
5 elderly people are inferior in controlled force exertion due to exercise (i.e., peripheral
6 muscular responses to the changing target and the exertion of neuromuscular
7 function) to young people, and they require more time to specify a movement
8 dimension (Stelmach, Goggin, & Garcia-Colera, 1987). The above-stated functional
9 developmental difference may produce decreases in exertion values or performance
10 with age.

11 According to studies by Aniansson, Rundgren, and Sperling (1980), Sperling
12 (1980), Ruff and Parker (1993), and Speller, Trollinger, Maurer, Nelson, and Bauer
13 (1997), males are superior in manual dexterity to females. Houx and Jolles (1993)
14 examined the sex differences in movement speed to reaction time using a manual
15 function in 20 to 80 year olds and reported that the males were superior in movement
16 speed to females in all age groups. Because females are inferior in manual dexterity
17 and movement speed to males, their controlled force exertion was also considered to
18 be inferior (Nagasawa, et al., 2000). However, a difference between males and
19 females was not found in the majority of the age groups or in the improvement rate.
20 Factors such as the above development difference of neuromuscular function
21 controlling exercise, adaptability to a new task, and the sex difference in learning
22 skill for both sexes may influence very little the rate of decrease in performance with
23 age group, because none of subjects previously experienced a controlled force
24 exertion test. Speller et al. (1997) reported that the assessment of movement
25 performance for a manual dexterity task is more appropriate in males with more

1 experience (manual dexterity). From the above, it is inferred that because manual
2 dexterity and movement speed are closely associated with movement experience in
3 daily activities, the sex difference in movement experience has an effect on controlled
4 force exertion. It is possible that continual exercise prevents the decrease in central
5 nervous system function related to high-level information processing, including
6 judgment, muscle volume, and motor performances, and improves controlled force
7 exertion (Skelton, et al., 1994). In the future, it will be necessary to examine the sex
8 difference of movement experience on controlled force exertion and the relationship
9 between controlled force exertion and exercise frequency in daily life.

10 The differences (effect sizes) between the 20-24yr. old group and groups older
11 than 40 years of age were large in both sexes. Stelmach et al. (1987) examined
12 whether the difference in information prior to the task response affects the elderly's
13 response time and movement time. They reported that, although the elderly persons
14 use pre-information similar to young people to prepare an upcoming movement, the
15 transaction times of information on the movement plan for arms (hands), direction
16 and extension were markedly slower, and the elderly persons required longer
17 movement times. Nakamura, Ide, Sugi, Terada, and Shibasaki (1995) reported that
18 the learning effect of pursuit movements is associated in both the knowledge of a
19 target-locus (declarative memory) and the improvement of procedure to pursue the
20 movement of a target (procedural memory). Although the present controlled force
21 exertion test was the same content (the same locus and speed) in all trials and the
22 information in advance was the same, the controlled force exertion still decreased
23 with age. Of the two memory types, declarative and procedural, the latter is
24 considered to control learning and to make exertion values decrease with age. The
25 present results clarified that the mean in groups older than 40 years of age enlarges

1 in both sexes as compared with that of 20-24yr. old group.

2 Individual differences of the errors in controlled force exertion showed a similar
3 tendency in males and females and tended to increase in individuals older than 60
4 years of age (the elderly) in both sexes. The exertion type of muscular strength
5 examined in this study differs from that of previous studies, but Butki (1994)
6 reported that subjects need 4 trials to gain some familiarity and to show a significant
7 improvement. Experience with a task and the practice effect influence controlled
8 force exertion and may produce the individual difference. Some elderly people may
9 have poorer adaptive functions, perhaps contributing to a floor effect wherein
10 individual differences in performance are small. In contrast, elderly subjects with
11 superior adaptive functions quickly learn the task and individual differences become
12 larger. It appears that such an increase in individual differences in performance
13 occurs in an elderly group. Nagasawa et al. (2004) reported that the ability exerted
14 by a type of displayed demand value is somewhat different in the controlled force
15 exertion test. Hence, when the demand values differ, the above test will need to be
16 examined in terms of age and sex differences. Moreover, it is necessary to establish
17 an evaluation standard value which can present demand values according to sex and
18 age to diagnose arm function and physical fitness of the aged in medical and
19 rehabilitation fields.

20 In conclusion, the errors in controlled force exertion tended to increase
21 constantly with age, but the rate of increase was significant for those greater than 40
22 years of age in both sexes. The change in individual differences was similar for both
23 sexes.

24

1

2 **Acknowledgements**

3 This study was supported in part by a Grant-in-Aid for Scientific Research
4 (project number 13780048 and 17700476) to Y. Nagasawa from the Ministry of
5 Education, Culture, Sports, Science and Technology of Japan, which we gratefully
6 acknowledge.

7

1 **References**

- 2 Aniansson, A., Rundgren, A., & Sperling, L. (1980). Evaluation of functional capacity
3 in activities of daily living in 70-year-old men and women. *Scandinavian Journal*
4 *of Rehabilitation and Medicine*, 12, 145-154.
- 5 Bemben, M. G., Massey, B. H., Bemben, D. A., Misner, J. E., & Boileau, R. A. (1991).
6 Isometric muscle force production as a function of age in healthy 20- to 74-yr.-old
7 men. *Medicine and Science in Sports and Exercise*, 23, 1302-1310.
- 8 Bemben, M. G., Massey, B. H., Bemben, D. A., Misner, J. E., & Boileau, R. A. (1996).
9 Isometric intermittent endurance of four muscle groups in men aged 20-74 yr.
10 *Medicine and Science in Sports and Exercise*, 28, 145-154.
- 11 Brown, S. W., & Bennett, E. D. (2002). The role of practice and automaticity in
12 temporal and nontemporal dual-task performance. *Psychological Research*, 66,
13 80-89.
- 14 Butki, B. D. (1994). Adaptation to effects of an audience during acquisition of rotary
15 pursuit skill. *Perceptual and Motor Skills*, 79, 1151-1159.
- 16 Cauley, J. A., Petrini, A. M., LaPorte, R. E., Sandler, R. B., Bayles, C. M., Robertson,
17 R. J., & Slemenda, C. W. (1987). The decline of grip strength in the menopause:
18 relationship to physical activity, estrogen use and anthropometric factors.
19 *Journal of Chronic Diseases*, 40, 115-120.
- 20 Dustman, R. E., Ruhling, R. O., Russell, E. M., Shearer, D. E., Bonekat, H. W.,
21 Shigeoka, J. W., Wood, J. S., & Bradford, D. C. (1984). Aerobic exercise training
22 and improved neuropsychological function of older individuals. *Neurobiological*
23 *Aging*, 5, 35-42.
- 24 Galganski, M. E., Fuglevand, A. J., & Enoka, R. M. (1993) Reduced control of
25 motor output in a human hand muscle of elderly subjects during submaximal

1 contractions. *Journal of Neurophysiology*, 69, 2108-2115.

2 Halaney, M. E., & Carey, J. R. (1989). Tracking ability of hemiparetic and healthy

3 subjects. *Physical Therapy*, 69, 342-348.

4 Hayashi, Y. (1967). [Human factors on manual control system]. [*Japan Journal of*

5 *Ergonomics*], 3, 265-274. [in Japanese]

6 Henatsch, H. -D., & Langer, H. H. (1985). Basic neurophysiology of motor skills in

7 sport: a review. *International Journal of Sports Medicine*, 6, 2-14.

8 Houx, P. J., & Jolles, J. (1993). Age-related decline of psychomotor speed: effects of

9 age, brain health, sex, and education. *Perceptual and Motor Skills*, 76(1),

10 195-211.

11 Kornhuber, H. H. (1974). Cerebral cortex, cerebellum and basal ganglia, an

12 introduction to their motor functions. In: F. O. Schmitt (Ed.), *The Neurosciences*

13 *III. Study Program* (pp. 267-280). MIT Press: Cambridge- Massachusetts.

14 Laboratory of Physical Education, Tokyo Metropolitan University. (Ed.) (1989).

15 [*Physical Fitness Standards of Japanese People*]. (4th ed.) Tokyo, Japan:

16 Fumaido. [in Japanese]

17 Meshizuka, T., & Nagata, A. (1972). A method for measuring muscular group

18 "control-ability" and its apparatus. *Research Journal of Physical Education*, 16,

19 319-325.

20 Nagasawa, Y., & Demura, S. (2002). Development of an apparatus to estimate

21 coordinated exertion of force. *Perceptual and Motor Skills*, 94, 899-913.

22 Nagasawa, Y., Demura, S., & Kitabayashi, T (2004). Concurrent validity of tests to

23 measure the coordinated exertion of force by computerized target-pursuit.

24 *Perceptual and Motor Skills*, 98:551-560.

25 Nagasawa, Y., Demura, S., & Nakada, M (2003). Trial-to-trial and day-to-day

1 reliability of a computerized target-pursuit system to measure the ability to
2 coordinate exertion of force. *Perceptual and Motor Skills*, 96:1071-1085.

3 Nagasawa, Y., Demura, S., Yamaji, S., Kobayashi, H., & Matsuzawa, J. (2000).
4 Ability to coordinate exertion of force by the dominant hand: comparisons among
5 university students and 65- to 78- year-old men and women. *Perceptual and*
6 *Motor Skills*, 90, 995-1007.

7 Nakamura, M., Ide, J., Sugi, T., Terada, K., & Shibasaki, H. (1995). [Method for
8 studying learning effect on manual tracking of randomly moving visual
9 trajectory and its application to normal subjects]. [*The transactions of the*
10 *Institute of Electronics, Information and Communication Engineers. D-II*],
11 J78-D-II(3), 547-558.

12 Oldfield, R. C. (1971). The assessment and analysis of handedness: the Edinburgh
13 inventory. *Neuropsychologia*, 9, 97-113.

14 Ranganathan, V. K., Siemionow, V., Sabgal, V., & Yue, G. H. (2001) Effects of aging on
15 hand function. *Journal of American Geriatrics Society*, 49, 1478-1484.

16 Rikli, R. E., & Busch, S. (1986). Motor performance of women as a function of age and
17 physical activity level. *Journal of Gerontology*, 41, 645-649.

18 Rikli, R. E., & Edwards, D. J. (1991). Effects of a three-year exercise program on
19 motor function and cognitive processing speed in older women. *Research*
20 *Quarterly*, 62, 61-67.

21 Ruff, R. M., & Parker, S.B. (1993). Gender- and age-specific changes in motor speed
22 and eye-hand coordination in adults: normative values for the Finger Tapping
23 and Grooved Pegboard Tests. *Perceptual and Motor Skills*, 76, 1219-1230.

24 Skelton, D. A., Greig, C. A., Davies, J. M., & Young, A. (1994). Strength, power and
25 related functional ability of healthy people aged 65-89 years. *Age and Ageing*, 23,

1 371-377.

2 Speller, L., Trollinger, J. A., Maurer, P. A., Nelson, C. E. & Bauer, D. F. (1997).

3 Comparison of the test-retest reliability of the Work Box using three

4 administrative methods. *American Journal of Occupational Therapy*, 51,

5 516-522.

6 Sperling, L. (1980). Evaluation of upper extremity function in 70-year-old men and

7 women. *Scandinavian Journal of Rehabilitation and Medicine*, 12, 139-144.

8 Stelmach, G. E., Goggin, N. L., & Garcia-Colera, A. (1987). Movement specification

9 time with age. *Experimental Aging Research*, 13, 39-46.

10 Voelcker-Rehage, C., & Alberts, J. L. (2005) Age-related changes in grasping force

11 modulation. *Experimental Brain Research*, 166, 61-70.

12 Walamies, M., & Turjanmaa, V. (1993). Assessment of the reproducibility of strength

13 and endurance handgrip parameters using a digital analyzer. *European Journal*

14 *of Applied Physiology and Occupational Physiology*, 67, 83-86.

15 Welford, A. T. (1988) Reaction time, speed of performance, and age. *Annals of the*

16 *New York Academy of Science*, 515, 1-17.

17 Yamamoto, T. (1983) [*Control-ability of movements*]. Tokyo: Kyorinshoin. [in

18 Japanese]

Figure. 1. Bar chart display (100 mm X 140 mm) of demand value. [Left bar (A) shows the demand value and right bar (B) is the exertion value of grip strength. The test was to fit line B (exertion value of grip strength) to line A (demand value), which varied in a span of 50 mm on the display. The test time was 40 sec for each trial. The controlled force exertion was calculated using the data from 25 sec of the trial following the initial 15 sec of the 40-sec period. Actual force was shown on the display, left.]

Figure 2. Age-group means of the controlled force-exertion test of the bar chart demand in males (◆) and females (■)
. *p<0.05.

Table 1. Physical characteristics of participants

Age Group (yr.)	<i>n</i>	Age (yr.)		Height (cm)		Weight (kg)		Grip strength (kgf)	
		<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>
Male									
15-19	27	17.2	1.5	171.4	5.4	63.3	9.0	42.0	7.1
20-24	29	21.9	1.4	171.1	4.6	68.2	7.1	51.2	6.3
25-29	25	27.8	1.3	172.9	5.0	69.4	8.2	48.8	8.0
30-39	25	34.4	3.0	173.1	5.7	72.1	10.8	48.0	7.7
40-49	25	44.9	2.8	169.2	7.0	67.4	7.2	46.4	7.7
50-59	23	54.5	2.9	166.2	6.2	65.8	8.4	41.1	7.3
60-69	27	64.3	3.0	165.0	6.2	63.4	9.3	37.0	7.8
70+	26	74.6	4.2	159.8	6.7	57.0	9.9	27.7	7.7
Total	207	42.1	19.8	168.6	7.2	65.8	9.7	42.8	10.3
Female									
15-19	27	17.1	1.4	159.1	5.2	53.4	5.3	29.1	4.9
20-24	38	22.2	1.3	160.1	4.7	52.8	5.3	31.8	4.4
25-29	27	27.0	1.4	159.3	5.8	51.0	6.4	30.8	4.9
30-39	41	35.1	2.6	158.4	4.8	51.8	7.3	29.4	3.9
40-49	27	44.6	2.7	157.0	5.0	52.3	5.9	30.0	3.8
50-59	26	53.2	3.0	154.7	5.2	54.8	7.5	28.9	4.4
60-69	36	63.7	2.8	153.0	6.0	55.6	8.3	25.1	6.5
70+	27	74.9	4.0	147.6	4.4	51.7	7.8	20.4	4.7
Total	249	41.7	19.1	156.3	6.4	53.0	6.9	28.3	5.8

Table 2. Distribution characteristics of controlled force-exertion scores

Age group (yr.)	<i>n</i>	Upper quartile	<i>Mdn</i>	Lower quartile	Skew	Kurtosis	Shapiro-Wilk's W	<i>P</i>
Male								
15-19	27	678.0	602.8	561.3	1.1	1.3	0.90	0.01
20-24	29	712.0	585.6	480.8	0.0	-1.1	0.95	0.24
25-29	25	783.3	581.8	507.5	1.3	1.4	0.89	0.01
30-39	25	771.4	647.5	566.6	0.6	0.0	0.96	0.42
40-49	25	851.0	666.0	572.0	0.9	0.1	0.91	0.02
50-59	23	1107.5	903.6	705.1	1.1	1.1	0.92	0.06
60-69	27	1245.7	958.2	833.9	0.5	-0.8	0.95	0.21
70+	26	1359.2	1084.2	954.3	1.5	1.4	0.83	<0.01
Total	207	956.9	730.4	582.5	1.8	4.6	0.87	<0.01
Female								
15-19	27	798.0	758.9	683.9	1.2	2.7	0.86	<0.01
20-24	38	814.0	648.0	557.7	1.0	0.6	0.92	0.01
25-29	27	808.9	672.7	582.0	1.0	0.9	0.94	0.10
30-39	41	894.5	796.7	688.0	0.8	1.6	0.96	0.22
40-49	27	979.0	900.6	731.3	0.3	-0.5	0.96	0.33
50-59	26	1072.4	900.1	814.8	0.9	0.9	0.93	0.09
60-69	36	1446.7	1110.6	907.2	0.6	-0.5	0.95	0.12
70+	27	1787.1	1506.9	1282.6	1.2	0.7	0.87	<0.01
Total	249	1067.9	829.9	694.8	2.4	8.2	0.79	<0.01

Table3. Means, standard deviations (%), and coefficients of variation and effect size by age group for controlled force-exertion score in the bar chart demand

Age group (yr.)	<i>n</i>	<i>M</i>	<i>SD</i>	<i>CV</i>	<i>ES</i>
Male					
15-19	27	627.8	125.2	19.95	0.31
20-24	29	587.1	137.8	23.47	—
25-29	25	659.8	214.3	32.47	0.40
30-39	25	674.6	135.4	20.07	0.64
40-49	25	730.6	212.5	29.09	0.80
50-59	23	946.1	289.0	30.54	1.59
60-69	27	1024.4	235.8	23.02	2.26
70+	26	1246.5	433.4	34.77	2.05
Female					
15-19	27	751.6	153.0	20.35	0.36
20-24	38	692.3	173.8	25.10	—
25-29	27	702.1	142.3	20.26	0.06
30-39	41	806.2	180.7	22.41	0.64
40-49	27	885.1	157.0	17.74	1.16
50-59	26	943.9	190.5	20.18	1.38
60-69	36	1168.0	336.6	28.82	1.78
70+	27	1675.1	607.1	36.24	2.20

Note. - *ES* shows the effect size of mean differences between trials of those in their 20-24 yr and each age group trial.