Methanotrophic community composition based on pmoA genes in dissolved methane recovery and biological oxidation closed downflow hanging sponge reactors

著者	Matsuura Norihisa, Hatamoto Masashi, Yamaguchi Takashi, Ohashi Akiyoshi
journal or publication title	Biochemical Engineering Journal
volume	124
page range	138-144
year	2017-08-15
URL	http://hdl.handle.net/2297/48598

doi: <https://doi.org/10.1016/j.bej.2017.05.003>

Methanotrophic community composition based on *pmoA* genes in dissolved methane recovery and biological oxidation closed downflow hanging sponge reactors

Norihisa Matsuura ^{a,b*}, Masashi Hatamoto ^b, Takashi Yamaguchi ^c and Akiyoshi Ohashi ^d

^a Faculty of Environmental Design, Institute of Science and Engineering, Kanazawa University, Kakuma, Kanazawa, Ishikawa 920-1192, Japan

^b Department of Environmental System Engineering, Nagaoka University of Technology, 1603-1 Kamitomioka, Nagaoka, Niigata 940-2188, Japan

^c Department of Science of Technology Innovation, Nagaoka University of Technology, Kamitomioka 1603-1, Nagaoka, Niigata 940-2188, Japan.

^d Department of Social and Environmental Engineering, Graduate School of Engineering, Hiroshima University, 1-4-1 Kagamiyama, Higashihiroshima, Hiroshima 739-8527, Japan

*Corresponding author:

Norihisa Matsuura

^a Faculty of Environmental Design, Institute of Science and Engineering, Kanazawa University, Kakuma, Kanazawa, Ishikawa 920-1192, Japan

Tel: +81-76-264-6436

Fax: +81-76-264-6436

E-mail: matsuura@se.kanazawa-u.ac.jp (N. Matsuura)

Fig. S1 Terminal restriction fragment length polymorphism (T-RFLP) analysis of *pmoA* genes amplified from upper, middle, and lower parts of sludge from the two closed downflow hanging sponge (DHS) reactors in summer and winter. The numbers at the terminal restriction fragment (T-RF) peaks indicate the T-RF lengths.