

Some transformation formulas for Lauricella's hypergeometric functions FD

メタデータ	言語: eng 出版者: 公開日: 2017-10-03 キーワード (Ja): キーワード (En): 作成者: メールアドレス: 所属:
URL	http://hdl.handle.net/2297/19390

Some transformation formulas for Lauricella's hypergeometric functions F_D

Keiji MATSUMOTO and Katsuyoshi OHARA

August 7, 2008

Abstract

In this paper, we give some functional equations with a parameter c for Lauricella's hypergeometric functions; they can be regarded as multivariable versions of the Gauss quadratic transformation formula for the hypergeometric function. These functional equations for $c = 1$ are utilized for the study of arithmetic-geometric means of several terms.

MSC2000: 33C65.

Keywords: hypergeometric functions, transformation formulas.

1 Introduction

It is known that the hypergeometric function $F(\alpha, \beta, \gamma; z)$ satisfies the Gauss quadratic transformation formula:

$$(1+z)^{2\alpha} F\left(\alpha, \alpha - \beta + \frac{1}{2}, \beta + \frac{1}{2}; z^2\right) = F\left(\alpha, \beta, 2\beta; \frac{4z}{(1+z)^2}\right).$$

When $\alpha = \beta = \frac{1}{2}$, this equality reduces to

$$\frac{1+z}{2} F\left(\frac{1}{2}, \frac{1}{2}, 1; 1-z^2\right) = F\left(\frac{1}{2}, \frac{1}{2}, 1; 1 - \left(\frac{2\sqrt{z}}{1+z}\right)^2\right),$$

which implies that the reciprocal of the arithmetic-geometric mean of 1 and $x \in (0, 1)$ coincides with $F(\frac{1}{2}, \frac{1}{2}, 1; 1-x^2)$, refer to [HKM].

In this paper, we give some functional equations with a parameter c for Lauricella's hypergeometric functions F_D ; they can be regarded as multivariable versions of the Gauss quadratic transformation formula. Our functional equations for $c = 1$ are given in [KS1],[KS2] and [KM], and they imply the expressions of arithmetic-geometric means of several terms by Lauricella's hypergeometric functions F_D . We also show that each of our functional equations admits no other parameters when we specify the transformations of variables of F_D and an admissible factor to the product of power functions associated with singular locus of F_D .

By considering restrictions of variables for our theorems, we obtain three transformation formulas for the hypergeometric function $F(\alpha, \beta, \gamma; z)$. We remark that they are not listed in [E] and [G], and that one of them was recently found in [BBG].

For proofs of our theorems, we utilize yang [O1], which is a package of computer algebra system Risa/Asir for the ring of differential-difference operators.

2 Lauricella's hypergeometric function F_D

Lauricella's hypergeometric function F_D of m variables z_1, \dots, z_m with parameters $\alpha, \beta_1, \dots, \beta_m, \gamma$ is defined as

$$F_D(\alpha, \beta, \gamma; z) = \sum_{n_1, \dots, n_m \geq 0}^{\infty} \frac{(\alpha, \sum_{j=1}^m n_j) \prod_{j=1}^m (\beta_j, n_j)}{(\gamma, \sum_{j=1}^m n_j) \prod_{j=1}^m (1, n_j)} \prod_{j=1}^m z_j^{n_j},$$

where $z = (z_1, \dots, z_m)$ satisfies $|z_j| < 1$ ($j = 1, \dots, m$), $\beta = (\beta_1, \dots, \beta_m)$, $\gamma \neq 0, -1, -2, \dots$ and $(\alpha, n) = \alpha(\alpha + 1) \cdots (\alpha + n - 1) = \Gamma(\alpha + n)/\Gamma(\alpha)$. This function admits the integral representation of Euler type:

$$F_D(\alpha, \beta, \gamma; z) = \frac{\Gamma(\gamma)}{\Gamma(\alpha)\Gamma(\gamma - \alpha)} \int_0^1 t^\alpha (1-t)^{\gamma-\alpha} \prod_{j=1}^m (1-z_j t)^{-\beta_j} \frac{dt}{t(1-t)}. \quad (1)$$

When $m = 1$, $F_D(\alpha, \beta, \gamma; z)$ coincides with the Gauss hypergeometric function $F(\alpha, \beta, \gamma; z)$, and when $m = 2$, $F_D(\alpha, \beta, \gamma; z)$ is Appell's hypergeometric function $F_1(\alpha, \beta_1, \beta_2, \gamma; z_1, z_2)$.

is $\bigcup_{1 \leq i < j \leq m+2} L_{ij} \subset \mathbb{C}^m$, where

$$L_{ij} = \{\ell_{ij} = z_i - z_j = 0\} \quad (2)$$

and $L_{m+1, m+2}$ is regarded as the hyperplane at infinity.

3 Transformation formulas

Theorem 1 *The hypergeometric function F_D of 2 variables satisfies the transformation formula*

$$\begin{aligned} & \left(\frac{1+z_1+z_2}{3} \right)^c F_D\left(\frac{c}{3}, \frac{c+1}{6}, \frac{c+1}{6}, \frac{c+1}{2}; 1-z_1^3, 1-z_2^3\right) \\ &= F_D\left(\frac{c}{3}, \frac{c+1}{6}, \frac{c+1}{6}, \frac{c+5}{6}; z'_1, z'_2\right), \end{aligned} \quad (3)$$

$$\xi : (z_1, z_2) \mapsto (z'_1, z'_2) = \left(\left(\frac{1+\omega z_1 + \omega^2 z_2}{1+z_1+z_2} \right)^3, \left(\frac{1+\omega^2 z_1 + \omega z_2}{1+z_1+z_2} \right)^3 \right)$$

where $\omega = \frac{-1+\sqrt{-3}}{2}$, $z = (z_1, z_2)$ is in a small neighborhood U of $(1, 1)$, and the value of $\left(\frac{1+z_1+z_2}{3}\right)^c$ at $(z_1, z_2) = (1, 1)$ is 1. This functional equation admits no other parameters when we specify the transformations of variables of the both sides of F_D in (3) and an admissible factor to $\Delta(z) = a \prod_{i=1}^k p_i(z)^{c_i}$, where $p_1(z), \dots, p_k(z)$ are the irreducible factors of the product $\prod_{1 \leq i < j \leq m+2} \xi^* \ell_{ij}$ of the pull-back of ℓ_{ij} in (2) for $m = 2$ under the map ξ .

Theorem 2 *The hypergeometric function F_D of 2 variables satisfies the transformation formula*

$$\begin{aligned} & \left(\frac{1+\sqrt{z_1 z_2}}{2} \right)^{2c-1} F_D\left(\frac{c}{2}, \frac{2c-1}{4}, \frac{2c-1}{4}, c; 1-z_1^2, 1-z_2^2\right) \\ &= F_D\left(\frac{c}{2}, \frac{2c-1}{4}, \frac{2c-1}{4}, \frac{c+1}{2}; z'_1, z'_2\right), \end{aligned} \quad (4)$$

$$\begin{aligned} \xi : (z_1, z_2) &\mapsto (z'_1, z'_2), \\ z'_1 &= \frac{(\sqrt{(1-z_1^2)(1-z_2^2)} - \sqrt{-1}(z_1 - z_2))^2}{(1 + \sqrt{z_1 z_2})^4}, \\ z'_2 &= \frac{(\sqrt{(1-z_1^2)(1-z_2^2)} + \sqrt{-1}(z_1 - z_2))^2}{(1 + \sqrt{z_1 z_2})^4}, \end{aligned}$$

where $z = (z_1, z_2)$ is in a small neighborhood U of $(1, 1)$, the values of $\sqrt{z_1 z_2}$ and $(\frac{1+\sqrt{z_1 z_2}}{2})^{2c-1}$ at $(z_1, z_2) = (1, 1)$ are 1, and the value of $\sqrt{(1-z_1^2)(1-z_2^2)}$ in the expression of z'_2 is same as that of z'_1 . Though z'_1 and z'_2 are exchanged by the choice of branches of $\sqrt{(1-z_1^2)(1-z_2^2)}$, the right hand side of (4) is single-valued by the coincidence of the parameters β_1 and β_2 of F_D . This functional equation admits no other parameters when we specify the transformations of variables of the both sides of F_D in (3) and an admissible factor to $\Delta(z) = a \prod_{i=1}^k p_i(z)^{c_i}$, where $p_1(z), \dots, p_k(z)$ are the irreducible factors of the product $\prod_{1 \leq i < j \leq m+2} \xi^* \ell_{ij}$ of the pull-back of ℓ_{ij} in (2) for $m = 2$ under the map ξ .

Theorem 3 *The hypergeometric function F_D of 3 variables satisfies the transformation formula*

$$\begin{aligned} & \left(\frac{1+z_1+z_2+z_3}{4} \right)^{\frac{c}{2}} F_D\left(\frac{c}{4}, \frac{c+2}{12}, \frac{c+2}{12}, \frac{c+2}{12}, \frac{c+2}{3}; 1-z_1^2, 1-z_2^2, 1-z_3^2\right) \\ &= F_D\left(\frac{c}{4}, \frac{c+2}{12}, \frac{c+2}{12}, \frac{c+2}{12}, \frac{c+5}{6}; z'_1, z'_2, z'_3\right), \end{aligned} \quad (5)$$

$$\xi : (z_1, z_2, z_3) \mapsto (z'_1, z'_2, z'_3),$$

$$z'_1 = \left(\frac{1-z_1-z_2+z_3}{1+z_1+z_2+z_3} \right)^2,$$

$$z'_2 = \left(\frac{1-z_1+z_2-z_3}{1+z_1+z_2+z_3} \right)^2,$$

$$z'_3 = \left(\frac{1+z_1-z_2-z_3}{1+z_1+z_2+z_3} \right)^2,$$

where $z = (z_1, z_2, z_3)$ is in a small neighborhood U of $(1, 1, 1)$, and the value of $(\frac{1+z_1+z_2+z_3}{4})^{c/2}$ at $(z_1, z_2, z_3) = (1, 1, 1)$ is 1. This functional equation admits no other parameters when we specify the transformations of variables of the both sides of F_D in (3) and an admissible factor to $\Delta(z) = a \prod_{i=1}^k p_i(z)^{c_i}$, where $p_1(z), \dots, p_k(z)$ are the irreducible factors of the product $\prod_{1 \leq i < j \leq m+2} \xi^* \ell_{ij}$

of the pull-back of ℓ_{ij} in (2) for $m = 3$ under the map ξ .

Remark 1 The transformation formulas (3) and (4) for $c = 1$ are utilized for the study of arithmetic-geometric means of three terms in [KS1] and [KS2], respectively. The transformation formula (5) for $c = 1$ appears in [KM] as Proposition 1, which is a key to express the common limit of a quadruple sequence by Lauricella's hypergeometric function F_D of 3 variables.

4 Proof

In this section, we prove Theorem 1. Since we can show the others similarly, we omit their proofs.

Let $\Omega_{\hat{f}}(z)$ and $\Omega_{\hat{g}}(z)$ be the connection 1-forms in Fact 1 for

$$\begin{aligned}\hat{f}(z) &= {}^t(f_0(z_1, z_2), z_1 \frac{\partial f_0}{\partial z_1}(z_1, z_2), z_2 \frac{\partial f_0}{\partial z_2}(z_1, z_2)), \\ f_0(z_1, z_2) &= F_D(\alpha, \beta_1, \beta_2, \gamma; z_1, z_2),\end{aligned}$$

and

$$\begin{aligned}\hat{g}(z) &= {}^t(g_0(z_1, z_2), z_1 \frac{\partial g_0}{\partial z_1}(z_1, z_2), z_2 \frac{\partial g_0}{\partial z_2}(z_1, z_2)), \\ g_0(z_1, z_2) &= F_D(\alpha', \beta'_1, \beta'_2, \gamma'; z_1, z_2),\end{aligned}$$

respectively. It is easy to see that the vector-valued functions

$$f(z) = {}^t(f_0, \frac{\partial f_0}{\partial z_1}, \frac{\partial f_0}{\partial z_2}) \text{ and } g(z) = {}^t(g_0, \frac{\partial g_0}{\partial z_1}, \frac{\partial g_0}{\partial z_2})$$

satisfy the Pfaffian systems

$$df = \Omega_f(z)f, \quad dg = \Omega_g(z)g,$$

respectively, where

$$\Omega_f(z) = P\Omega_{\hat{f}}(z)P^{-1} + dPP^{-1}, \quad \Omega_g(z) = P\Omega_{\hat{g}}(z)P^{-1} + dPP^{-1},$$

$$P = \text{diag}\left(1, \frac{1}{z_1}, \frac{1}{z_2}\right) = \begin{pmatrix} 1 & & \\ & \frac{1}{z_1} & \\ & & \frac{1}{z_2} \end{pmatrix}.$$

Consider the vector-valued function

$$G(x) = {}^t(G_0, \frac{\partial G_0}{\partial x_1}, \frac{\partial G_0}{\partial x_2})$$

for the pull-back $G_0(x_1, x_2)$ of $g_0(z_1, z_2)$ under the map

$$\xi : (x_1, x_2) \mapsto (z_1, z_2) = \left(\left(\frac{1 + \omega x_1 + \omega^2 x_2}{1 + x_1 + x_2} \right)_3, \left(\frac{1 + \omega^2 x_1 + \omega x_2}{1 + x_1 + x_2} \right)_3 \right).$$

It satisfies

$$G(1, 1) = {}^t(1, 0, 0) \quad (6)$$

and the Pfaffian system $dG = \Omega_G(x)G$, where

$$\Omega_G(x) = J_2 \Omega_g(x) J_2^{-1} + dJ_2 J_2^{-1}, \quad J_2 = \begin{pmatrix} 1 & \\ & {}^t J \end{pmatrix},$$

$\Omega_g(x)$ is the pull-back of $\Omega_g(z)$ under the map ξ , and J is the Jacobi matrix of the map ξ . The singular locus of $\Omega_G(x)$ consists of 12 lines $p_i(x) = 0$ ($i = 1, \dots, 12$), where

$$\begin{aligned} p_1(x) &= 1 + x_1 + x_2, & p_2(x) &= x_1 - \omega, & p_3(x) &= x_1 - \omega^2, \\ p_4(x) &= x_2 - \omega, & p_5(x) &= x_2 - \omega^2, & p_6(x) &= x_1 - \omega x_2, \\ p_7(x) &= x_1 - \omega^2 x_2, & p_8(x) &= x_1 - x_2, & p_9(x) &= x_1 - 1, \\ p_{10}(x) &= x_2 - 1, & p_{11}(x) &= 1 + \omega x_1 + \omega^2 x_2, & p_{12}(x) &= 1 + \omega^2 x_1 + \omega x_2. \end{aligned}$$

Put $a_i = p_i(1, 1)$; note that $a_i \neq 0$ for $i = 1, \dots, 7$, $a_i = 0$ for $i = 8, \dots, 12$. Since $f_0(1, 1) = g_0(1, 1) = 1$, we have $\Delta(1, 1) = 1$. Thus $\Delta(x)$ should be

$$\Delta(x) = \prod_{i=1}^7 \left(\frac{p_i(x)}{a_i} \right)^{c_i}.$$

Consider the vector-valued function

$$F(x) = {}^t(F_0, \frac{\partial F_0}{\partial x_1}, \frac{\partial F_0}{\partial x_2})$$

for

$$F_0(x_1, x_2) = \Delta(x) f_0(1 - x_1^3, 1 - x_2^3).$$

It satisfies

$$F(1, 1) = {}^t(1, \sum_{i=1}^7 \frac{c_i}{a_i} \frac{\partial p_i}{\partial x_1} - \frac{3\alpha\beta_1}{\gamma}, \sum_{i=1}^7 \frac{c_i}{a_i} \frac{\partial p_i}{\partial x_2} - \frac{3\alpha\beta_2}{\gamma}) \quad (7)$$

and the Pfaffian system $dF = \Omega_F(x)F$, where

$$\Omega_F(x) = Q[J_1 \Omega_f(x) J_1^{-1} + dJ_1 J_1^{-1}] Q^{-1} + dQQ^{-1}, \quad J_1 = \text{diag}(1, -3x_1^2, -3x_2^2),$$

$$Q = \begin{pmatrix} \frac{\Delta(x)}{\frac{\partial \Delta(x)}{\partial x_1}} & & & \\ \frac{\partial \Delta(x)}{\partial x_1} & \Delta(x) & & \\ \frac{\partial \Delta(x)}{\partial x_2} & & \Delta(x) & \\ & & & \Delta(x) \end{pmatrix} = \Delta(x) \begin{pmatrix} 1 & & & \\ \frac{\partial \log \Delta(x)}{\partial x_1} & 1 & & \\ \frac{\partial \log \Delta(x)}{\partial x_1} & & & \\ \frac{\partial \log \Delta(x)}{\partial x_1} & & & 1 \end{pmatrix},$$

and $\Omega_f(x)$ is the pull-back of $\Omega_f(z)$ under the map

$$(x_1, x_2) \mapsto (z_1, z_2) = (1 - x_1^3, 1 - x_2^3).$$

The singular locus of $\Omega_F(x)$ consists of 12 lines $x_1 = 0$, $x_2 = 0$ and $p_i(x) = 0$ ($i = 1, \dots, 10$).

Note that $F_0(x) = G_0(x)$ on U if and only if $F(1, 1) = G(1, 1)$ and $\Omega_F(x) = \Omega_G(x)$. By (6) and (7), we have

$$\frac{c_1}{3} + \frac{c_2 + c_6}{1 - \omega} + \frac{c_3 + c_7}{1 - \omega^2} - \frac{3\alpha\beta_1}{\gamma} = \frac{c_1}{3} + \frac{c_4 - c_6\omega}{1 - \omega} + \frac{c_5 - c_7\omega^2}{1 - \omega^2} - \frac{3\alpha\beta_2}{\gamma} = 0. \quad (8)$$

We compare the entries of $\Omega_F(x)$ with those of $\Omega_G(x)$ by utilizing yang [O1], which is a package of computer algebra system Risa/Asir for the ring of differential-difference operators. We have a necessary and sufficient condition for the identity $\Omega_F(x) = \Omega_G(x)$ expressed as a system of 802 linear and 399 quadratic equations of 15 variables ${}^t v = (c_1, \dots, c_7, \alpha, \beta_1, \beta_2, \gamma, \alpha', \beta'_1, \beta'_2, \gamma')$. The 802 linear equations include the followings 14 linear equations:

$$\begin{aligned} -c_1 - c_4 - c_5 - c_6 - c_7 + 3\beta_2 + \alpha' + \beta'_1 + \beta'_2 - \gamma' &= 0, \\ -2c_4 - 2c_5 + 3\alpha + 3\beta_2 - 3\gamma - 2\alpha' + \beta'_1 + \beta'_2 + 2\gamma' &= 1, \\ -2c_6 - 2c_7 + 3\alpha + 3\beta_1 + 3\beta_2 - 3\gamma - \alpha' - \beta'_1 - \beta'_2 + \gamma' &= 0, \\ -c_6 - c_7 + 3\beta_1 - \alpha' - \beta'_1 - \beta'_2 + \gamma' &= 1, \\ c_6 + c_7 + \alpha' + \omega\beta'_1 - (\omega + 1)\beta'_2 - \gamma' &= -1, \\ 3\alpha - 3\beta_1 + 3\beta_2 - 3\beta'_1 - 3\beta'_2 &= -1, \\ \alpha' - 2\beta'_1 - 2\beta'_2 + 2\gamma' &= 1, \\ 3\alpha + 3\beta_1 - 3\gamma &= -1, \\ 3\alpha + 3\beta_2 - 3\gamma &= -1, \\ (\omega + 1)c_2 - \omega c_3 &= 0, \\ c_2 + c_3 &= 0, \\ (\omega + 1)c_4 - \omega c_5 &= 0, \\ c_4 + c_5 &= 0, \\ \omega c_6 - (\omega + 1)c_7 &= 0. \end{aligned}$$

Thus we have a 1-dimensional solution space of these 14 linear equations, which can be expressed as

$$\begin{aligned} \alpha = \alpha' = \frac{c_1}{3}, \quad \beta_1 = \beta_2 = \beta'_1 = \beta'_2 = \frac{c_1 + 1}{6}, \\ \gamma = \frac{c_1 + 1}{2}, \quad \gamma' = \frac{c_1 + 5}{6}, \quad c_2 = c_3 = c_4 = c_5 = c_6 = c_7 = 0, \end{aligned} \quad (9)$$

where we regard c_1 as a free parameter in \mathbb{C} .

We can see that the solution (9) satisfies the 802 linear and 399 quadratic equations and (8) by Risa/Asir. Hence $F_0(x) = G_0(x)$ on U if and only if the condition (9) holds. Refer to [O2] for our computation by yang and Risa/Asir. \square

5 Restrictions

In this section, we derive some corollaries by considering restrictions of variables for our transformation formulas.

Corollary 1 (Theorem 2.3 in [BBG]) *We have*

$$\begin{aligned} & \left(\frac{1+2z}{3} \right)^c F\left(\frac{c}{3}, \frac{c+1}{3}, \frac{c+1}{2}; 1-z^3 \right) \\ &= F\left(\frac{c}{3}, \frac{c+1}{3}, \frac{c+5}{6}; \left(\frac{1-z}{1+2z} \right)^3 \right) \end{aligned}$$

for z sufficiently near to 1, where the value of $\left(\frac{1+2z}{3} \right)^c$ at $z = 1$ is 1.

Proof. Put $z = z_1 = z_2$ for the transformation formula (3) and use the integral representation (1). \square

Corollary 2 *We have*

$$\begin{aligned} & \left(\frac{1+z}{2} \right)^{2c-1} F\left(\frac{c}{2}, \frac{2c-1}{4}, c; 1-z^4 \right) \\ &= F\left(\frac{c}{2}, \frac{2c-1}{2}, \frac{c+1}{2}; -\left(\frac{1-z}{1+z} \right)^2 \right). \end{aligned}$$

for z sufficiently near to 1, where the value of $\left(\frac{1+z}{2} \right)^{2c-1}$ at $z = 1$ is 1.

Proof. Put $z = \sqrt{z_1}$, $z_2 = 1$ for the transformation formula (4) and use the integral representation (1). \square

Corollary 3 *We have*

$$\begin{aligned} & \left(\frac{1+3z}{4}\right)^{\frac{c}{2}} F\left(\frac{c}{4}, \frac{c+2}{4}, \frac{c+2}{3}; 1-z^2\right) \\ &= F\left(\frac{c}{4}, \frac{c+2}{4}, \frac{c+5}{6}; \left(\frac{1-z}{1+3z}\right)^2\right). \end{aligned}$$

for z sufficiently near to 1, where the value of $\left(\frac{1+3z}{4}\right)^{c/2}$ at $z = 1$ is 1.

Proof. Put $z = z_1 = z_2 = z_3$ for the transformation formula (3) and use the integral representation (1). \square

Remark 2 The equalities in Corollaries 1 and 3 for $c = 1$ are used in [BB] to study modified arithmetic-geometric means.

Remark 3 It is written in [BBG] that Corollary 1 can not be deduced from the cubic transformation formulas in [G]. The authors remark that our corollaries are not listed in [E] and [G], and think that Corollaries 2 and 3 can not be obtained by classical results.

References

- [BB] J.M. Borwein and P.B. Borwein, A cubic counterpart of Jacobi's identity and the AGM, *Trans. Amer. Math. Soc.* **323**(2) (1991), 691–701.
- [BBG] B.C. Berndt, S. Bhargava and F.G. Garvan, Ramanujan's theories of elliptic functions to alternative bases, *Trans. Amer. Math. Soc.*, **347** (1995), 4163–4244.
- [E] A. Erdélyi, *Higher Transcendental Functions*, volume I, Robert E. Krieger Publishing Co., Inc., Melbourne, Florida, 1981.
- [G] E.M. Goursat, Sur l'Équation Différentielle Linéaire qui Admet pour Intégrale la Série Hypergéométrique, *Ann. Sci. l'Ecole Normale Sup.* (2) **10** (1881), 3–142.
- [HKM] R. Hattori, T. Kato and K. Matsumoto, Mean iterations derived from transformation formulas for the hypergeometric function, *preprint* 2008.

- [IKSY] K. Iwasaki, H. Kimura, S. Shimomura and M. Yoshida, *From Gauss to Painlevé*, Vieweg, Braunschweig, Wiesbaden, 1991.
- [KM] T. Kato and K. Matsumoto, The common limit of a quadruple sequence and the hypergeometric function F_D of three variables, *preprint* 2007.
- [KS1] K. Koike and H. Shiga, Isogeny formulas for the Picard modular form and a three terms arithmetic geometric mean, *J. Number Theory* **124** (2007), 123–141.
- [KS2] K. Koike and H. Shiga, Extended Gauss AGM and corresponding Picard modular forms, to appear in *J. Number Theory*.
- [O1] K. Ohara, yang — a package for computation in the ring of differential-difference operators, <http://www.openxm.org>, 2007.
- [O2] K. Ohara, Calculations of some transformation for hypergeometric functions, <http://www.math.kobe-u.ac.jp/OpenXM/Math/AGM>, 2008.

Keiji MATSUMOTO
(matsu@math.sci.hokudai.ac.jp)
Department of Mathematics
Hokkaido University
Sapporo 060-0810, Japan

Katsuyoshi OHARA
(ohara@kanazawa-u.ac.jp)
Faculty of Mathematics and Physics
Kanazawa University
Kanazawa 920-1192, Japan