

Systematic studies on the Conducting Tissue of the Gametophyte in Musci (14) Anatomy of the Stems of Rhizogonium, Mnium and Fissidens

メタデータ	言語: eng 出版者: 公開日: 2017-10-03 キーワード (Ja): キーワード (En): 作成者: 河合, 功 メールアドレス: 所属:
URL	https://doi.org/10.24517/00011194

This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 International License.



Systematic Studies on the Conducting Tissue of the Gametophyte in Musci

(14) Anatomy of the Stems of *Rhizogonium*, *Mnium* and *Fissidens*

Isawo KAWAI, Sadako YOSHITAKA and Emi YAMAMOTO

Department of Biology, Faculty of Science, Kanazawa University

(Received October 31, 1986)

Abstract Experiments on the absorption of pigments by the gametophytes of *Rhizogonium*, *Mnium* and *Fissidens* in which each tissue within the stems was stained with different dyes, led to the conclusion that the stems of *Rhizogonium*, *Mnium* and *Fissidens* appear to be differentiated into an epidermis, cortex, leptome and hydrome.

In future studies, it is hoped that the relationships existing among gametophytes of Musci will be investigated according to the color adopted by each tissue, those showing similar dyeing properties being considered to be homologous tissues.

Introduction

Previous investigations on the absorption of pigments by the gametophytes of *Polytrichum*, in which each tissue within the stem was stained with different dyes, led to the following results. (1) The cell walls of the epidermis were dyed red with eosin solution. (2) The cell walls of the cortex were dyed blue-green with methyl green solution. (3) The cytoplasm of the leptome was dyed red with eosin solution. (4) The cell walls of the hydrome were dyed violet-brown with a respective of solutions of aniline blue and eosin, Janus green and eosin, and Congo red and gentian violet.

Using the Congo red-gentian violet-eosin-methyl green-method it was shown that in *Lepidium virginicum* L. the cytoplasm of phloem was dyed red with eosin, and that the cell walls of xylem vessels were dyed violet-brown with the solution of Congo red and gentian violet (Figs. 1 and 2).

In conformity with these results, the inner structures of the stems of certain members of the Musci were investigated, and the findings are reported here.

Materials and Methods

The materials used in this study were samples of the mosses *Rhizogonium*, *Mnium* and *Fissidens*. Observations were conducted after subjecting the mosses to the following staining procedures.




Fig. 1 (x 60) and Fig. 2 (x 200) : Cross sections of the stem of *Lepidim Virginicum* L. The cell walls of the xylem vessels were dyed violet with the solution of Congo red and gentian violet. The cytoplasm of the phloem were dyed red with eosin solution.

(1) *Rhizogonium*

- a) A solution of aniline blue and eosin was absorbed into the gametophyte for seventy-two hours.
- b) After washing, a solution of eosin was absorbed into the gametophyte for seventy-two hours.
- c) After further washing, a solution of methyl green was absorbed into the gametophyte, again for seventy-two hours. After washing, cross-sections of the moss about $15\mu\text{m}$ in thickness were cut with a cryo-microtome and mounted in gum arabic.

(2) *Mnium*

- a) A solution of aniline blue and eosin was absorbed into the gametophyte for seventy-two hours.
- b) After washing, a solution of eosin was absorbed into the gametophyte for seventy-two hours.
- c) After further washing, a solution of methyl green was absorbed into the gametophyte, again for seventy-two hours. After washing, cross-sections of the moss about $15\mu\text{m}$ in thickness were cut with a cryo-microtome and mounted in gum arabic.

(3) *Fissidens*

- a) A solution of Janus green and eosin was absorbed into the gametophyte for thirty-six hours.
- b) After washing, a solution of eosin was absorbed for a further thirty-six hours.
- c) After further washing, a solution of methyl green was absorbed into the moss, again for thirty-six hours. After washing cross-sections of the moss about $15\mu\text{m}$ in thickness were cut with a cryo-microtome and mounted in gum arabic.

Results and Discussion

In the vascular plant *Lepidium*, the cell walls of the xylem vessels were dyed violet with the Congo red and gentian violet, and the cytoplasm of the phloem was dyed red with eosin.

In *Polytrichum* of the Musci, the cell walls of the hydrome were dyed violet with the combination of Congo red and gentian violet, anilin blue and eosin, and Janus green and eosin, and the chloroplasts of the cortex and the cytoplasm of the leptome were dyed red with eosin. (Plate I)

From these results, it may be considered that the xylem vessels of *Lepidium* are homologous to the hydrome of *Polytrichum*, and that the phloem of *Lepidium* is homologous to the leptome of *Polytrichum*.

In *Rhizogonium*, the cell walls of the epidermis were dyed red with eosin, and the cell walls of the cortex were dyed blue-green with methyl green. The chloroplasts of the cortex and the cytoplasm of the several layers inside the cortex were dyed red with eosin. These results suggest that in *Rhizogonium*, the several layers inside the cortex may be homologous to the leptome of *Polytrichum*. Similarly, the central part of the stem in *Rhizogonium* may be homologous to the hydrome of *Polytrichum*, because the cell walls of the latter were dyed violet-blue with the solution of aniline blue and eosin. (Plate I)

In *Mnium*, the cell walls of the epidermis were dyed red with eosin, and the cell walls of the cortex were dyed blue-green with methyl green. The chloroplasts of the cortex and the cytoplasm of several layers inside the cortex were dyed red with eosin. These results suggest that in *Mnium*, the several layers inside the cortex may be homologous to the leptome of *Polytrichum*. Similarly, the central part of the stem in *Mnium* may be homologous to the hydrome of *Polytrichum*, because the cell walls of the latter were dyed violet-blue with the solution of aniline blue and eosin. (Plate II)

In *Fissidens*, the cell walls of the epidermis were dyed red with eosin, and the cell walls of the cortex were dyed blue-green with methyl green. The chloroplasts of the cortex and the cytoplasm of several layers inside the cortex were dyed red with eosin. These results suggest that in *Fissidens*, the several layers inside the cortex may be considered to be homologous to the leptome of *Polytrichum*. Similarly, the central part of the stem in *Fissidens* may be homologous to the hydrome of *Polytrichum*, because the cell walls of the central part of the stem were dyed violet-blue with the solution of aniline blue and eosin. (Plate II)




Plate I. Cross sections of the stem

Fig. 1-Fig. 4: *Polytrichum*, Fig. 5-Fig. 8: *Rhizogonium*.

Fig. 1: The cell walls of the hydrome were dyed violet-blue with the solution of aniline blue and eosin.

Fig. 2: The cytoplasm of the leptome were dyed red with the eosin solution. Fig. 3: The cell walls of the cortex were dyed blue with the methyl green solution. Fig. 4: The cell walls of the epidermis were dyed red with the eosin solution and the cell walls of the cortex were dyed blue-green with the methyl green solution (distilled water). Fig. 5: The cell walls of the hydrome were dyed blue with the solution of aniline blue and eosin. Figs. 6, 7: The cytoplasm of the leptome were dyed red with the eosin solution. Fig. 8: The cell walls of the cortex were dyed green with the methyl green solution.




Plate II. Cross sections of the stem

Fig. 1-Fig. 4: *Mnium*, Fig. 5-Fig. 8: *Fissidens*

Figs. 1, 2: The cell walls of the hydrome were dyed violet-blue with the solution of aniline blue and eosin. Fig. 3: The cytoplasm of the leptome were dyed red with the eosin solution. Fig. 4: The cell walls of the cortex were dyed blue with the methyl green solution. Fig. 5: The cell walls of the hydrome were dyed violet-brown with the solution of aniline blue and eosin. Fig. 6: The cytoplasm of the leptome were dyed red with the eosin solution. Fig. 7: The cell walls of the cortex were dyed blue with the methyl green solution. Fig. 8: The cell walls the epidermis were dyed red with the eosin solution, and the cell walls of the cortex were dyed blue with the methyl green solution (distilled water).

From these experimental results, it appears that the stems of *Rhizogonium*, *Mnium* and *Fissidens* are differentiated into an epidermis, cortex, leptome and hydrome.

References

- FREY, W. (1974) Bau und Entstehung der Zentralstrangverzweigung bei Laubmoosen. *Nova Hedwigia* 25 : 211-233.
- FRYE, T. C. (1946) *Pogonatum convolutum* (HEDW.) P. BEAUV. *Bryologist* 49 : 36-40.
- (1948) *Pogonatum semipellucidum*. *Bryologist* 51 : 248-250.
- HÉBANT, C. (1966) Differentiation des tissus conducteurs dans la tige feuillée. De *Polytrichum juniperinum* WILLD. C. R. Acad. Sc. Paris 262 : 2585-2588.
- (1967) Structure et différenciation des tissus conducteurs dans le gaméophyte des *Polytrichum*. *Nat. Monsspelienica, Bot.* 18 : 293-298.
- (1969) Histologie végétale-nouvelles observations sur le leptome de la tige feuillée des *Polytrichum*. C. R. Aca. Sc. Paris 269 : 2530-2533.
- (1970) A new look at the conducting tissues of mosses (bryopsida). Their structure, distribution and significance. *Phytomorphology* 20 : 390-410.
- (1973) Acid phosphomonoesterase activities (β -glycerophosphatase and naphthol AS-MX phosphatase) in conducting tissues. *Protoplasma* 77 : 231-241.
- (1974) Studies on the development of the conducting tissue-system in the gametophytes of some Polytrichales. II. Development and structure at maturity of the hydroids of the central strand. *Journ. Hattori Bot. Lab.* 38 : 565-607.
- (1975) The phloem (leptome) of Bryophytes. *Phloem transport* : New York 211-215.
- (1976) Studies on the development of the conducting tissue-system in the gametophyte of some Polytrichales. III. Further observations on leptoids, with particular reference to their endoplasmic reticulum. *Protoplasma* 87 : 79-90.
- (1977) The conducting tissues of Bryophytes. *Cramer. Vaduz* 22-77.
- HORMANN, H. (1963) Beitrag zur Anatomie von *Pogonatum baldwini* (MUELL.) PAR. *Nova Hedwigia* 5 : 279-282.
- KAWAI, I. and K. Ikeda (1970) Systematic studies on the conducting tissue of the gametophyte in Musci (1) On the affinity regarding the conducting tissue of some species of the Polytrichaceae. *Sci. Rep. Kanazawa Univ.* 15 : 71-98.
- KAWAI, I. (1971) ——— (2) On the affinity regarding the inner structure of the stem in some species of Dicranaceae, Bartramiaceae, Entodontaceae and Fissidentaceae. *Ann. Rep. Bot. Garden, Fac. Sci. Univ. Kanazawa* 4 : 18-39.
- KOPONEN, T. (1980) A synopsis of Mniaceae (Bryophyta). IV Taxa in Europe, Macaronesia, NW Africa and the near East. *Ann. Bot. Fennici* 17 : 125-162.
- LORENTZ, P. G. (1867) Studien zur vergleichenden Anatomie der Laubmoose II. *Flora* 50 : 529-540, 545-558.
- MESSMER, L. W. and T. C. Frye (1947) The *Polytrichum juniperinum* group between America and the United States. *Bryologist* 50 : 259-268.
- OSADA, T. A. Noguchi (1962) *Pogonatum inflexum* (LINDB.) PAR. and its allies. *Journ. Bot.* 37 : 361-365.
- (1965) Japanese Polytrichaceae I. Introduction and the genus *Pogonatum*. *Journ. Hattori Bot. Lab.* 28 : 171-201.

- (1966) Japanese Polytrichaceae II. The genera *Polytrichum*, *Oligotrichum*, *Bartramiopsis* and *Atrichum*, and phytogeography. Journ. Hattori Bot. Lab. 29 : 1-52.
- SCHEIRER, D. C. (1972) Anatomical studies in the Polytrichaceae. I. The gametophor of *Dendroligotrichum dendroides* (HEDW.) BROTH. Bryologist 75 : 305-314.
- (1973) Hydrolysed walls in the water-conducting cells of *Dendroligotrichum* (Bryophyta). Histochemistry and ultrastructure. Planta 115 : 37-46.
- (1975) Anatomical studies in the Polytrichaceae II. Histochemical observations on thickened lateral walls of Hydroides of *Dendroligotrichum*. Bryologist 78 : 113-123.
- (1976) Some fine structural observations on the rhizome of *Dendroligotrichum* (Bryophyta). Protoplasm 89 : 323-337.
- TANSLEY, A. G. (1901) Notes on the conducting tissue-system in Bryophyta. Annales Bot. 1155 : 1-38.
- TRACHTENBERG, S and E. ZAMSKI (1978) Conduction of ionic solutes and assimilates in the leptom of *Polytrichum juniperinum* WILLD. Journ. Exp. Bot. 29 : 719-727.
- ZAMSKI, E. and S. TRACHTENBERG (1976) Water movement through hydroids of a moss gametophyte. Israel Journ. Bot. 25 : 168-173.