

Water in amine-mediated single electron transfer reaction of N-allylic trichloroacetamides

メタデータ	言語: eng 出版者: 公開日: 2017-10-04 キーワード (Ja): キーワード (En): 作成者: メールアドレス: 所属:
URL	https://doi.org/10.24517/00015336

This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 International License.



WATER IN AMINE-MEDIATED SINGLE ELECTRON TRANSFER REACTION OF *N*-ALLYLIC TRICHLOROACETAMIDES


Tsuyoshi Taniguchi, Masamichi Sasaki, and Hiroyuki Ishibashi*

School of Pharmaceutical Sciences, Institute of Medical, Pharmaceutical and Health Sciences, Kanazawa University, Kakuma-machi, Kanazawa 920-1192, Japan. E-mail: isibasi@p.kanazawa-u.ac.jp

Abstract – Water contaminating 1,4-dimethylpiperazine was thought to play an important role in effecting a single electron transfer reaction (radical cyclization) of *N*-allylic trichloroacetamides.

INTRODUCTION

We have recently reported that *N*-allylic trichloroacetamides (**1**), upon heating in 1,4-dimethylpiperazine, gave γ -lactams (**2**) in good yields (Scheme 1).¹⁻³ These reactions might involve a single electron transfer reaction of 1,4-dimethylpiperazine to **1**. Compounds (**1**) gave dichloro-substituted carbamoylmethyl radicals after removal of a mono-chlorine atom. Cyclization of these radicals to an olefinic bond and successive addition reaction of a H-atom at the resulting terminal radical intermediates gave γ -lactams (**2**). Our attention was next turned to the application of this method to synthesis of (-)-kainic acid (**3**).^{4,5} We report herein that water contaminating 1,4-dimethylpiperazine plays an important role in effecting a single electron transfer reaction (radical cyclization).




Scheme 1

RESULTS AND DISCUSSION


We initiated our investigation by examining the cyclization of compound (**8**) giving (+)-kainic acid in boiling 1,4-dimethylpiperazine in place of that giving the desired (-)-kainic acid. Synthesis of radical

precursor (**8**) was begun by conversion of L-serine (**4**) into (-)-Garner's aldehyde (**5**).⁶ Horner-Wadsworth-Emmons reaction of compound (**5**) gave the known *E*-ester (**6**).⁷ After transformation of compound (**6**) into **7**, the hydroxy group of compound (**7**) was protected with a *tert*-butyldiphenylsilyl (TBDPS) group and the nitrogen atom was trichloroacetylated to give radical precursor (**8**) (Scheme 2).



Scheme 2

When compound (**8**) was heated in 1,4-dimethylpiperazine (1,4-DMP), which is commercially available and was used without further purification, the expected radical cyclization product (**9**) (stereoisomers' ratio = 9:1) was obtained in 28-38% yield together with *tert*-butyldiphenylsilanol (**10**)⁸ after 10 min of heating (Scheme 3).




Scheme 3

The stereochemistry between C4 and C5 of the major isomer of compound (**9**) was probably *trans*-configuration.⁹ We next carried out a similar reaction in pure 1,4-DMP, since reproducibility was not observed, and the yield of product (**9**) was relatively low. However, the yield of **9** from compound (**8**) in distilled (pure) 1,4-DMP was found to be lower (7%) than that using non-distilled

1,4-dimethylpiperazine (Table 1, Entry 1). In this case, silanol (**10**) was obtained as a major product in 67% yield.

We soon found that the yield of compound (**9**) was improved to 57% by addition of 5 equiv of water to distilled 1,4-DMP (Table 1, Entry 2). The addition of 0.2 equiv of water gave a similar result (56% yield of **9**) (Table 1, Entry 3). However, when 100 equiv of water was added to 1,4-DMP, compound (**9**) was obtained in only 19% yield with an increase in the yield of silanol (**10**) (80%) (Table 1, Entry 4). Therefore, an appropriate quantity of water was found to play an important role in effecting a single electron transfer reaction (radical cyclization).

Table 1. Effect of water in amine-mediated radical cyclization of **8**.



Entry	H ₂ O (equiv)	Yield (%)	
		9	10
1	none	7	67
2	5	57	11
3	0.2	56	11
4	100	19	80

It was thought that the formation of silanol (**10**) might involve elimination of **8** to give diene (**11**), which was decomposed under the present reaction conditions (Fig. 1). The presence of a small amount of water might accelerate the radical cyclization to give compound (**9**) in good yield. The exact reason, however, is not clear at the moment.

CONCLUSION

We found that water contaminating 1,4-dimethylpiperazine played an important role in effecting a single electron transfer reaction (radical cyclization).




Figure 1

EXPERIMENTAL

Ethyl (R)-4-(N-benzylamino)-5-hydroxy-2-pentenoate (7) Acetyl chloride (2.3 g, 29.4 mmol) was added dropwise to EtOH (20 mL) at 0 °C over 5 min, and the mixture was stirred at the same temperature for 10 min. To the resultant solution was added dropwise a solution of **6** (4.4 g, 14.7 mmol) in EtOH (10 mL) at 0 °C over 5 min. After the mixture was stirred at room temperature for 20 h, the solution was concentrated to give brown oil. To a solution of the residue in CH₂Cl₂ (70 mL) were added successively Et₃N (1.5 g, 14.7 mmol), MgSO₄ (4.4 g, 36.7 mmol) and benzaldehyde (2.3 g, 22.0 mmol) at 0 °C and the mixture was stirred at room temperature for 3 h. The reaction mixture was filtered through celite[®] and the filtrate was concentrated. The residual yellow oil was diluted with MeOH (50 mL) and to the resultant solution was added NaBH₄ (4.5 g, 36.7 mmol) in portion at 0 °C. The reaction was quenched by addition of water and the reaction mixture was extracted with AcOEt. The organic phase was washed with brine, dried (Na₂SO₄), and concentrated. The residue was chromatographed on silica gel (hexane/AcOEt, 3:1 to 1:1 to 1:2) to give **7** (3.1 g, 85%) as a yellow oil: $[\alpha]_D^{24}$ -63.8 (*c* 2.4, CHCl₃); IR (CHCl₃) ν 1715 cm⁻¹; ¹H NMR (270 MHz, CDCl₃) δ 1.30 (3H, t, *J* = 7.1 Hz), 2.39 (2H, br s), 3.33-3.41 (1H, m), 3.44 (1H, dd, *J* = 10.1, 7.4 Hz), 3.64 (1H, dd, *J* = 10.1, 3.6 Hz), 3.64 (1H, d, *J* = 13.0 Hz), 3.84 (1H, d, *J* = 13.0 Hz), 4.20 (2H, q, *J* = 7.1), 6.00 (1H, dd, *J* = 15.8, 0.8 Hz), 6.79 (1H, dd, *J* = 15.8, 7.3 Hz), 7.25-7.35 (5H, m); ¹³C NMR (67.8 MHz, CDCl₃) δ 14.1, 51.0, 60.1, 60.5, 63.8, 123.4, 127.2, 128.1, 128.4, 139.4, 146.5, 166.0; HRMS calcd for C₁₄H₁₉NO₃: 249.1365. Found: 249.1363.

Ethyl (R)-4-(N-benzyl-N-trichloroacetyl-amino)-5-(*t*-butyldiphenylsilyloxy)-2-pentenoate (8) To a solution of **7** (500 mg, 2.01 mmol) in DMF (2 mL) were added Et₃N (507 mg, 5.02 mmol), DMAP (25 mg, 0.201 mmol) and TBDPSCl (1.10 g, 4.01 mmol) and the mixture was stirred at room temperature for 2 h. The resultant suspension was diluted with a saturated aqueous solution of NaHCO₃ and the mixture was extracted with Et₂O. The organic phase was washed with brine, dried (Na₂SO₄), and

concentrated. The residue was chromatographed on silica gel (hexane/AcOEt, 1:0 to 40:1 to 20:1 to 8:1) to give ethyl (*R*)-4-(*N*-benzylamino)-5-(*t*-butyldiphenylsilyloxy)-2-pentenoate (910 mg, 93%) as a colorless oil: $[\alpha]_D^{25}$ -32.4 (*c* 1.5, CHCl₃); IR (CHCl₃) ν 1715 cm⁻¹; ¹H NMR (270 MHz, CDCl₃) δ 1.04 (9H, s), 1.28 (3H, t, *J* = 7.1), 2.10 (1H, s), 3.36-3.43 (1H, m), 3.60 (1H, dd, *J* = 10.1, 3.1), 3.63 (1H, d, *J* = 13.5), 3.68 (1H, dd, *J* = 10.1, 4.8), 3.85 (1H, d, *J* = 13.5), 4.18 (2H, q, *J* = 7.1), 5.99 (1H, dd, *J* = 15.8, 1.0), 6.78 (1H, dd, *J* = 15.8, 7.4), 7.23-7.45 (11H, m), 7.58-7.62 (4H, m); ¹³C NMR (67.8 MHz, CDCl₃) δ 14.2, 19.2, 26.8, 51.2, 60.1, 60.3, 65.8, 123.3, 126.9, 127.7, 128.0, 128.4, 129.8, 132.9, 133.0, 135.50, 135.51, 140.1, 147.4, 166.2; HRMS calcd for C₃₀H₃₇NO₃Si: 487.2543. Found: 487.2545.

To a solution of thus obtained amine (910 mg, 1.87 mmol) in CH₂Cl₂ (20 mL) were added Et₃N (378 mg, 3.73 mmol) and trichloroacetyl chloride (441 mg, 2.43 mmol) at 0 °C and the mixture was stirred at room temperature for 30min. The reaction was quenched with a saturated aqueous solution of NaHCO₃ and the reaction mixture was extracted with CH₂Cl₂. The organic phase was dried (Na₂SO₄) and concentrated. The residue was chromatographed on silica gel (hexane/AcOEt, 20:1) to give **8** (1.11 g, 94%) as a colorless oil: $[\alpha]_D^{25}$ -38.8 (*c* 0.45, CHCl₃); IR (CHCl₃) ν 1680, 1715 cm⁻¹; ¹H and ¹³C NMR spectra of **8** showed it to contain two rotamers. ¹H NMR (270 MHz, CDCl₃) δ 1.01 (s), 1.06 (total 9H, both s), 1.20-1.35 (3H, m), 3.75-3.85, 4.00-4.40 (total 5H, both m), 4.88 (br d, *J* = 16.0), 5.38 (d, *J* = 16.2), 5.52 (d, *J* = 16.2), 6.03 (br d, *J* = 15.5), 6.95 (br dd, *J* = 16.0, 6.3), 7.07-7.60 (15H, m); HRMS calcd for C₃₂H₃₆³⁵Cl₃NO₄Si: 631.1480. Found: 631.1481.

(4*S*,5*R*)-1-Benzyl-5-(*t*-butyldiphenylsilyloxymethyl)-3,3-dichloro-4-(ethoxycarbonylmethyl)pyrrolidin-2-one (9) A mixture of **8** (30 mg, 0.0474 mmol), 1,4-dimethylpiperazine (541 mg, 4.74 mmol) and H₂O (4.3 mg, 0.237 mmol) was heated at reflux for 10 min. After cooling, the reaction mixture was diluted with a saturated aqueous solution of NH₄Cl and the mixture was extracted with CH₂Cl₂. The organic phase was dried (Na₂SO₄) and concentrated. The residue was purified by preparative thin layer chromatography (hexane/AcOEt, 3:1) to give **9** (16.2 mg, 57%) as a colorless oil: IR (CHCl₃) ν 1725, 1730 cm⁻¹; ¹H NMR (500 MHz, CDCl₃, for a major isomer) δ 1.09 (9H, s), 1.26 (3H, t, *J* = 7.1), 2.45 (1H, dd, *J* = 15.9, 6.1), 2.89 (1H, dd, *J* = 15.9, 6.8), 3.08 (1H, dt, *J* = 8.1, 2.4), 3.57-3.64 (3H, m), 3.73 (1H, dd, *J* = 12.0, 2.4), 4.14-4.22 (2H, m), 5.10 (1H, d, *J* = 15.1), 6.81-6.83 (2H, m), 7.19-7.21 (3H, m), 7.38-7.42 (2H, m), 7.44-7.47 (3H, m), 7.50-7.53 (1H, m), 7.61-7.63 (2H, m), 7.68-7.69 (2H, m); ¹³C NMR (125 MHz, CDCl₃, for a major isomer) δ 14.0, 19.1, 26.7, 32.7, 44.3, 46.7, 58.2, 59.8, 61.1, 85.6, 127.6, 127.7, 127.8, 128.0, 128.7, 130.0, 130.2, 132.3, 132.3, 134.6, 135.6, 135.8, 167.0, 170.3; HRMS calcd for C₃₂H₃₇³⁵Cl₂NO₄Si: 597.1869. Found: 597.1860, calcd for C₃₂H₃₇³⁷Cl₂NO₄Si: 601.1810. Found: 601.1812.

ACKNOWLEDGEMENT

This work was supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

REFERENCES

1. H. Ishibashi, S. Haruki, M. Uchiyama, O. Tamura, and J. Matsuo, *Tetrahedron Lett.*, 2006, **47**, 6263.
2. T. Taniguchi, R. Kawajiri, and H. Ishibashi, *Arkivoc*, 2008, xiv, 7.
3. H. Ishibashi, M. Sasaki, and T. Taniguchi, *Tetrahedron*, 2008, **64**, 7771.
4. For reviews on (–)-kainic acid, see: (a) M. G. Moloney, *Nat. Prod. Rep.*, 1998, **15**, 205. (b) M. G. Moloney, *Nat. Prod. Rep.*, 1999, **16**, 485. (c) M. G. Moloney, *Nat. Prod. Rep.*, 2002, **19**, 597.
5. For synthesis of (–)-kainic acid, see: H. Sakaguchi, H. Tokuyama, and T. Fukuyama, *Org. Lett.*, 2008, **10**, 1711 and references cited therein.
6. (a) A. Dondoni and D. Perrone, *Org. Synth.*, 2000, **77**, 64. (b) P. Garner and J. M. Park, *Org. Synth.*, 1992, **70**, 18.
7. K. L. Debendra, *J. Chem. Soc., Perkin Trans. 1*, 2001, 1851.
8. S. McN. Sieburth and W. Mu, *J. Org. Chem.*, 1993, **58**, 7584.
9. A. L. J. Beckwith and C. H. Schiesser, *Tetrahedron*, 1985, **41**, 3925.