

Zn speciation in two Fe-Mn banded systems

メタデータ	言語: eng
	出版者:
	公開日: 2017-10-05
	キーワード (Ja):
	キーワード (En):
	作成者: 田崎, 和江
	メールアドレス:
	所属:
URL	https://doi.org/10.24517/00035483

This work is licensed under a Creative Commons
Attribution-NonCommercial-ShareAlike 3.0
International License.



Zn speciation in two Fe-Mn banded system

Matthew Marcus¹, Alain Manceau², Micheal Kersten³ and Kazue Tazaki⁴

¹*Advanced Light Source, Lawrence Berkeley National Laboratory*

²*Enironmental Geochemistry Group, LGIT, University J. Fourier and CNRS*

³*Geoscience Institute, Gutenberg Institute*

⁴*Department of Earth Sciences, Faculty of Science, Kanazawa University*

What we did: Micro-XRF and micro-EXAFS at Zn and Mn on two Fe-Mn banded systems: looking at Fe/Mn/Zn distribution and Mn and Zn speciation.

Why we care: Fe-Mn banded systems are ubiquitous, occurring in soils, shallow oceans, deep oceans, hotspots,...




They have similarities despite differences in formation. Some are biogenic, some not.

Question: Are there common themes? How transferrable is the structural information? Does it matter if bacteria were involved?

Acknowledgements: This work was done at the Advanced Light Source, Beamline 10.3.2, and at the ESRF FAME beamline. The authors thank Robert Sublett for help with data collection. M. Kersten acknowledges support by Jan Harff from the Baltic Sea Research Institute.



Samples

Bacterial mat from Japan




Ref: Tazaki, K., *Clays and Clay Minerals* 48, 511-520 (2002)

Shallow-ocean nodule from Baltic




Ref: S. Hlawatsch, T. Neumann, C. M. G. van den Berg, M. Kersten, J. Harff, E. Suess, *Marine Geology* 182, 373-387 (2002)

Elemental maps



Bacterial mat from hot springs at seashore
Zn is at natural abundance.

Tricolor elemental maps



Zn at rim is anthropogenic and was deposited only when outer layers were growing.

Similarities: Fe/Mn alternation due to oscillating chemistry. Zn prefers Mn to Fe.

Differences: Japanese sample has Zn everywhere; Baltic one mostly in rim, due to natural/anthropogenic Zn source.

Zn less segregated in Japan sample than Baltic.



Zn XANES: Japan

Look at different points and analyzed by XANES.

Use PCA to characterize spectra

Colored points mark where spectra were taken. Colors match those on XANES and PCA plots


Result principal Components Analysis
Points colored according to PCA component strength



Mn map (high Mn=light)

Zn extended XANES from spots shown above

Isosbestic points (arrowed in insert) show that data approximates linear combinations of two components (end-members)




Results: XANES shows a clear variation with location on sample. Data are approximately described as linear combinations of two end-members, with fractions (related to PCA 2nd component strength) correlated with Fe/Mn ratio.


This is consistent with EXAFS picture of competing sorption on hydrous ferric oxides and phyllo-manganates.

Zn EXAFS: Baltic

Zn and Mn EXAFS and XANES is uniform over area tested, unlike Japan sample.




Sample




*Zn sorbed on turbostratic birnessite:
Tetrahedral Zn and some Mn³⁺*

Lab-made reference
Birnessite, [Zn]/[Mn]=0.008




*Tetrahedral and some octahedral
Zn, with Mn³⁺*

Zn EXAFS: Japan



*EXAFS differs from place to place,
unlike in Baltic nodule.*

*Fit of Mn-rich spot consistent with
mixture of Zn-sorbed ferrihydrite
(Note: XRD finds 6-line Fh) and
Zn-sorbed birnessite (similar to
Baltic).*





*Fit of Fe-rich area not yet clear
Possible phosphate component.*

*Zn in Japanese sample is
coordinated to ferrihydrite,
birnessite, and at least one
other species, in proportions
depending on Fe/Mn ratios.*

Mn EXAFS

Comparison of Baltic and Japan



Mn EXAFS from the two samples look mostly alike, but with differences. Analysis shows:

Baltic: Turbostratic (from XRD) birnessite with some Mn^{3+}

Japan: Consistent with mixture of birnessite and lithiophorite. Slight Mn-Mn distance expansion suggesting more Mn^{3+} in layer than in Baltic.

Both: Uniform over several spots of varying composition.