

HDL3 Cholesterol Levels in an Elderly Population

メタデータ	言語: eng 出版者: 公開日: 2018-05-10 キーワード (Ja): キーワード (En): 作成者: メールアドレス: 所属:
URL	https://doi.org/10.24517/00042065

This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 International License.





HDL₃ Cholesterol Levels in an Elderly Population

Akihiro Inazu^{1,2}

¹Department of Clinical Laboratory Science, School of Health Sciences, Kanazawa University, Kanazawa, Japan

²Wellness Promotion Science Center, Institute of Medical, Pharmaceutical and Health Sciences, Kanazawa University, Kanazawa, Japan

See article vol. 24: 000-000

The HDL hypothesis was almost negated by three negative results of three low-molecular-weight chemical CETP inhibitors in the phase 3 trials, although low CETP activity, high levels of large HDL (i.e., HDL₂), and decrease in coronary artery disease prevalence have been demonstrated in genetic epidemiological studies of several SNPs¹). However, there are still hopes for the revival of the HDL hypothesis. One has not been well tested whether or not enhanced lipoprotein lipase activity, TG lipolysis, and enhanced HDL synthesis, i.e., VLDL-HDL pathway hypothesis is anti-atherogenic²). The other is cellular cholesterol efflux and small HDL (i.e., HDL₃) hypothesis³).

HDL consists of heterogeneous particles in terms of density (1.063–1.21 g/mL) and has an apolipoprotein composition of apoA-I, apoA-II, apoCs, and apoE and lipid composition of FC, CE, TG, and PL. Recent studies have suggested that specific lipids, such as sphingomyelin, ceramide, and sphingosine-1-phosphate (S-1-P), may play a key role in the HDL's functions⁴). S-1-P receptors are coupled with G proteins, thereby stimulating cellular migration and proliferation and modulating vascular tonus, inflammation, and a barrier function between cells. S-1-P is rich in dense HDL⁵). Therefore, HDL₃ levels may act as S-1-P carriers, playing a specific role in vascular protection.

The current study by Dr. Notsu *et al.* investigated the association between HDL subclasses and carotid intima-media thickness in the Shimane CoHRE Study conducted in the Oki Island in the Japan Sea⁶). HDL₃ cholesterol is negatively associated

with age both in men and women. Simple correlation showed that max-IMT has an inverse relation with HDL₃ cholesterol levels, but multivariate analyses negated the association when age was included. One explanation for this finding is age acting as a confounding factor associated with both low HDL₃ cholesterol and increased carotid atherosclerosis (**Fig. 1**).

In terms of the future direction of HDL research, it would be interesting to examine whether or not low HDL₃ phenotype related with sedentary life could be corrected by enhancing ABCA1 or LPL activity through dietary and physical activity instruction or drug therapy.

Conflict of Interest

None.

References

- 1) Mabuchi H, Nohara A, Inazu A: Cholesteryl ester transfer protein (CETP) deficiency and CETP inhibitors. *Mol Cells*, 2014; 37: 777-784. doi: 10.14348/molcells.2014.0265
- 2) Brinton EA, Eisenberg S, Breslow JL: Increased apo A-I and apo A-II fractional catabolic rate in patients with low high density lipoprotein-cholesterol levels with or without hypertriglyceridemia. *J Clin Invest*, 1991; 87: 536-544
- 3) Mehta NN, Li R, Krishnamoorthy P, Yu Y, Farver W, Rodrigues A, Raper A, Wilcox M, Baer A, DerOhanesian S, Wolfe M, Reilly MP, Rader DJ, VanVoorhees A, Gelfand JM: Abnormal lipoprotein particles and cholesterol efflux capacity in patients with psoriasis. *Atherosclerosis*, 2012; 224: 218-221
- 4) Ståhlman M, Fagerberg B, Adiels M, Ekroos K, Chapman JM, Kontush A, Borén J: Dyslipidemia, but not hyperglycemia and insulin resistance, is associated with marked alterations in the HDL lipidome in type 2 diabetic subjects in the DIWA cohort: impact on small HDL particles. *Biochim Biophys Acta*, 2013; 1831: 1609-1617
- 5) Kontush A, Therond P, Zerrad A, Couturier M, Nègre-Salvayre A, de Souza JA, Chantepie S, Chapman MJ: Preferential sphingosine-1-phosphate enrichment and sphingomyelin depletion are key features of small dense

Address for correspondence: Akihiro Inazu, Department of Clinical Laboratory Science, School of Health Sciences, Kanazawa University, Kanazawa, Ishikawa, Japan
E-mail:

Received: March 22, 2017

Accepted for publication: March 23, 2017




Fig. 1. Clinical significance of HDL₃ cholesterol levels

Low HDL₃ cholesterol is correlated with carotid atherosclerosis, but it is not causal. Aging may be a confounding factor between HDL₃ cholesterol and carotid atherosclerosis.

HDL₃ particles: relevance to antiapoptotic and antioxidative activities. *Arterioscler Thromb Vasc Biol*, 2007; 27: 1843-1849

6) Notsu Y, Yano S, Takeda M, Yamasaki M, Isomura M,

Nabika T, Nagai A: Association of high-density lipoprotein subclasses with carotid intima-media thickness: Shimane CoHRE Study. *JAT*, 2017; in press