

Chromosome numbers of five species of Lamium (Labiatae) in Japan

メタデータ	言語: eng 出版者: 公開日: 2017-10-04 キーワード (Ja): キーワード (En): 作成者: メールアドレス: 所属:
URL	http://hdl.handle.net/2297/00049786

Norihito Miura¹, Yoshikane Iwatsubo² and Naohiro Naruhashi²: **Chromosome numbers of five species of *Lamium* (Labiatae) in Japan**

¹Nikko Museum of Natural History, Chuguji 2480-1, Nikko 321-1661, Japan ; ²Department of Biology, Faculty of Science, University of Toyama, Gofuku 3190, Toyama 930-8555, Japan

The genus *Lamium* L. (Labiatae), distributed in the north of Africa and Eurasia, comprises about 40 species (Mabberley 1997). In Japan the following six species including two naturalized species occur : *L. album* var. *barbatum*, *L. amplexicaule*, *L. humile*, *L. hybridum*, *L. purpureum* and *L. tuberiferum* (Murata and Yamazaki 1993 ; Murata 2003). As shown in Table 2, chromosome number of the *Lamium* taxa distributed in Japan is known for the following four species : *L. album* var. *barbatum* : $n=9$ and $2n=18$; *L. amplexicaule* : $n=9$ and $2n=18$; *L. hybridum* : $n=18$ and $2n=36$; and *L. purpureum* : $n=9$, $2n=14$ and 18 . Except for a report of $2n=14$ chromosomes in *L. purpureum* from SW Finland (Aronhson 1982, sec. Goldblatt 1988), the counts reported for these species correspond to the basic chromosome number of $x=9$ proposed for the genus by Darlington and Wylie (1955). In the course of our cytological studies for Japanese Labiatae, the authors found $2n=34$ chromosomes, along with $2n=18$ and 36 chromosomes in *Lamium*. This paper reports on the somatic chromosome numbers of *L. album* var. *barbatum*, *L. amplexicaule*, *L. humile*, *L. hybridum* and *L. purpureum* in Japan.

Materials and methods

A total of 28 individuals of *Lamium album* L. var. *barbatum* (Siebold et Zucc.) Franch. et Sav., *L. amplexicaule* L., *L. humile* (Miq.) Maxim., *L. hybridum* Vill. and *L. purpureum* L. collected from eight localities were used for the study (Table 1). These plants were cultivated in plastic pots at the experimental garden of University of Toyama, Japan. Their root tips were pretreated

in 2.0 or 2.1 mM 8-hydroxyquinoline solution at approximately 25°C for 1 h and then kept at 6°C for 15 h. They were fixed in a mixture of glacial acetic acid and ethyl alcohol (1 : 3) for 1 h, soaked in 1 N HCl at room temperature for 1 h, macerated in 1 N HCl at 60°C for 10 minutes, and then washed in tap water. The root tips were stained in a drop of 1.5% lacto-propionic orcein on the slide glass, and ordinary squash technique was applied for the examination of somatic chromosomes. Voucher specimens of these plants are deposited in the Toyama Science Museum (TOYA).

Results and discussion

Chromosome counts in the five taxa were as follows :

Lamium album var. *barbatum*


Two individuals collected from two localities (Table 1) were examined. They had $2n=18$ chromosomes, (Fig. 1, Table 2) which verified the counts reported previously for this taxon (Tanaka 1974 ; Terasaka and Tanaka 1974 ; Starodubtsev 1985, as *L. barbatum* ; Sokolovskaya et al. 1986, as *L. barbatum* ; Nishikawa 1981 ; Rudyka 1995, as *L. barbatum*).

Lamium amplexicaule

Six individuals collected from two localities listed in Table 1 were investigated. All showed $2n=18$ chromosomes (Fig. 2). As shown in Table 2, the count found in the present study agreed with almost all the counts reported previously for this taxon.

Lamium humile

Four individuals collected from one locality were investigated (Table 1). A chromosome count



Figs. 1–5. Somatic chromosomes of the five species in *Lamium* of Japan. 1: *L. album* var. *barbatum* ($2n=18$). 2: *L. amplexicaule* ($2n=18$). 3: *L. humile* ($2n=34$). 4: *L. hybridum* ($2n=36$). 5: *L. purpureum* ($2n=18$). Bar indicates 5 μ m.

Table 1. Collection localities and number of individuals examined (in parenthesis) of *Lamium* in Japan

Taxon	Collection locality
<i>L. album</i> var. <i>barbatum</i>	Toyama Pref. : Gofuku, Toyama City, (1). Ishikawa Pref. : Fukami, Wajia City, (1).
<i>L. amplexicaule</i>	Gunma Pref. : Kamishiroi, Komochi-mura, Kitagunma-gun, (1). Toyama Pref. : Gofuku, Toyama City, (5).
<i>L. humile</i>	Nara Pref. : Akadani, Ootou-mura, Yoshino-gun, (4).
<i>L. hybridum</i>	Gifu Pref. : Kamigiri, Takayama City, (1).
<i>L. purpureum</i>	Akita Pref. : Nigorikawa, Akita City, (1) ; Detomachigomon, Honjo City, (1) ; Komeyama, Yuri-machi, Yuri-gun, (1). Toyama Pref. : Gofuku, Toyama City, (12).

of $2n=34$ was obtained from all samples (Fig. 3, Table 2). This is the first report of chromosome number for this species.

Lamium hybridum

One individual was investigated (Table 1). This plant had $2n=36$ chromosomes (Fig. 4), agreeing with all the previous reports (Bernström 1944, 1955, as $n=18$; Löve and Löve 1956; Morton 1973; Fernandes and Leitão 1984; Elena-Rosselló et al. 1988; Ortega-Olivencia and Ruíz-Tellez 1990, as $n=18$) (Table 2).

Lamium purpureum

In this species, $n=9$ and $2n=14$, 18 chromosomes are known (Table 2). Fifteen individuals collected from four localities listed in Table 1 were investigated. These showed a somatic chromosome number of $2n=18$ (Fig. 5, Table 2).

In *Lamium*, two somatic chromosome numbers, $2n=18$ and 36, are known to be by far the most

common (Fedorov 1969). On the basis of the chromosome numbers, the basic number of *Lamium* was considered as $x=9$ by Darlington and Wylie (1955). In this genus, a unique chromosome number of $2n=34$ had been reported in *L. iranicum* Parsa (Aryavand 1977, as $n=17$). Present study disclosed that *L. humile* also has $2n=34$ chromosomes. Their somatic chromosome numbers show that they have a basic chromosome number of $x=17$.

The Labiatae, which is composed of approximately 6,700 species of 252 genera (Mabberley 1997), has basic chromosome numbers of $x=5-19$, within the basic numbers both $x=8$ and 9 are the most common primary basic numbers, and $x=17$ is the most common secondary basic number (Singh 1995). In Labiatae the secondary basic number of $x=17$ is considered by Singh (1995) to arise as the result of combination between the

Table 2. Chromosome numbers of *Lamium* taxa used in the study

Taxon	Present count (2n)	Previous count		Reference
		(n)	(2n)	
<i>L. album</i> var. <i>barbatum</i>	18	9		Terasaka and Tanaka (1974)
			18	Tanaka (1974), Starodubtsev (1985, as <i>L. barbatum</i> Siebold et Zucc.), Sokolovskaya et al. (1986), Nishikawa (1981), Rudyka (1995, as <i>L. barbatum</i> Siebold et Zucc.)
<i>L. amplexicaule</i>	18	9		Jørgensen (1927), Tischler (1934), Arohonka (1982, sec. Goldblatt 1988), Gill (1983), Saggoo and Bir (1983 a, 1983 b, 1986), Ortega-Olivencia and Ruíz-Tellez (1990)
			18	Bernström (1944, 1952, 1953a, 1955), Heiser and Whitaker (1948), Löve and Löve (1956), Strid (1965), Gadella and Kliphuis (1966), Skalinska et al. (1968), Májovsky et al. (1970), Dahlgren et al. (1971), Fernandes and Queirós (1971), Feráková (1972), Morton (1973), Löve and Kjellqvist (1974), Markova and Thu (1974), Aryavand (1977), Van den Brand et al. (1979), Strid and Franzén (1981), Gill (1983), Fernandes and Leitão (1984), Elena-Rosselló et al. (1988), Dobes et al. (1997)
<i>L. humile</i>	34			
<i>L. hybridum</i>	36	18		Bernström (1944, 1955), Ortega-Olivencia and Ruíz-Tellez (1990)
			36	Löve and Löve (1956), Morton (1973), Elena-Rosselló et al. (1988), Fernandes and Leitão (1984)
<i>L. purpureum</i>	18	9		Jørgensen (1927), Tischler (1934, 1937), Griesinger (1937), Gill (1983)
			14	Arohonka (1982, sec. Goldblatt 1988)
			18	Heitz (1926), Bernström (1944, 1953a, 1953b, 1955), Pólya (1949), Löve and Löve (1956), Sorsa (1963, as 2n=ca.18), Skalinska et al. (1968), Gadella and Kliphuis (1971), Morton (1973), Löve and Kjellqvist (1974), Markova and Thu (1974), Májovsky et al. (1976), Van den Brand et al. (1979), Fernandes and Leitão (1984), Elena-Rosselló et al. (1988), Hill (1989), Ortega-Olivencia and Ruíz-Tellez (1990)

two dissimilar primary numbers of 8 and 9. In *Lamium*, however, the primary basic number reported is merely 9, and 8 has not been found in any species in the genus. Thus, the basic chromosome number of $x=17$ found in both *L. humile* and *L. iranicum* seems to be arisen by a dysploidal change from $2n=36$ chromosomes having a primary basic chromosome number of $x=9$ to $2n=34$ chromosomes with the secondary basic chromosome number because a primary basic number of $x=8$, which is found in many genera in La-

biatae, has not been known in any species in *Lamium*.

The authors are indebted to two anonymous reviewers for their helpful comments on the manuscript. We thank Dr. M. Hakki for his kindness in correcting the manuscript.

References

- Aryavand, A. 1977. Löve, Á. (ed.). IOPB chromosome number reports LVII. Taxon **26**: 443-444.

- Bernström, P. 1944. Two new hybrids in *Lamium*. *Hereditas* **30**: 257-260.
- Bernström, P. 1952. Cytogenetic intraspecific studies in *Lamium* I. *Hereditas* **38**: 163-220.
- Bernström, P. 1953 a. Increased crossability in *Lamium* after chromosome doubling. *Hereditas* **39**: 241-256.
- Bernström, P. 1953 b. Cytogenetic intraspecific studies in *Lamium* II. *Hereditas* **39**: 381-437.
- Bernström, P. 1955. Cytogenetic studies on relationships between annual species of *Lamium*. *Hereditas* **41**: 1-122.
- Dahlgren, R., Karlsson, Th. and Lassen, P. 1971. Studies on the flora of the Balearic Islands I. Chromosome numbers in Balearic angiosperms. *Bot. Notiser* **124**: 249-269.
- Darlington, C.D. and Wylie, A.P. 1955. Chromosome atlas of flowering plants. 2nd ed. p.329. George Allen and Unwin, London.
- Dobes, C., Hahn, B. and Morawetz, W. 1997. Chromosomenzahlen zur Gefäßpflanzen-Flora Österreichs. *Linzer biol. Beitr.* **29**: 5-43.
- Elena-Rosselló, J. A., González Zapatero, M. A. and Navarro Andrés, F. 1988. Consideraciones cariológicas sobre algunas especies nitrófilas de *Lamium* L. *Acta Bot. Barcinon.* **37**: 149-154.
- Fedorov, A. A. 1969. Chromosome numbers of flowering plants. 926 pp. Komar. Bot. Inst., Acad. Sci. U.S.S.R., Leningrad. (in Russian)
- Feráková, V. 1972. Bemerkungen zur Flora der Umgebung der Stadt Hlohovec in der Slowakei III. Ergänzungen. *Acta Fac. Rerum Nat. Univ. Comenianae Bot.* **20**: 149-158.
- Fernandes, A. and Leitão, M. T. 1984. Contribution á l'étude cytotaxinomique des spermatophyta du Portugal XVIII-Lamiaceae. *Mem. Soc. Broteriana* **27**: 27-75.
- Fernandes, A. and Queirós, M. 1971. Sur la caryologie de quelques plantes récoltées pendant la III^{ème} Réunion de Botanique Péninsulaire. *Mem. Soc. Broteriana* **21**: 343-385.
- Gadella, Th. W. J. and Kliphuis, E. 1966. Chromosome numbers of flowering plants in the Netherlands II. *Proc. Roy. Neth. Acad. Sci. Ser. C.* **69**: 541-556.
- Gadella, Th. W. J. and Kliphuis, E. 1971. Chromosome numbers of flowering plants in the Netherlands V. *Proc. Kon. Ned. Acad. Wetensch. Ser. C* **74**: 335-343.
- Gill, L. S. 1983. Cytotaxonomic studies of the tribe Stachydeae (Labiatae) in Canada. *Willdenowia* **13**: 175-181.
- Goldblatt, P. 1988. Index to plant chromosomes numbers 1984-1985. *Monogr. Syst. Bot. Missouri Bot. Gard.* **23**: 126.
- Griesinger, R. 1937. Über hypo-und hyperdiploide Formen von *Petunia*, *Hyoscyamus*, *Lamium* und einige andere Chromosomenzählungen. *Ber. Deutsch. Bot. Ges.* **55**: 556-571.
- Heiser, C. B. and Whitaker, T. W. 1948. Chromosome number, polyploidy, and growth habit in California weeds. *Amer. Jour. Bot.* **35**: 179-186.
- Heitz, E. 1926. Der nachweis der chromosomen. Vergleichende studien über ihre zahl, grösse und form im Pflanzenreich I. *Zeitsch. Bot.* **18**: 625-681.
- Hill, L. M. 1989. IOPB chromosome data 1. *Int. Organ. Pl. Biosyst. Newslett. (Zurich)* **13**: 17-19.
- Jørgensen, C. A. 1927. Cytological and experimental studies in the genus *Lamium*. *Hereditas* **9**: 126-136.
- Löve, Á. and Kjellqvist, E. 1974. Cytotaxonomy of Spanish plants. IV. Dicotyledons: Caesalpiniaceae-Asteraceae. *Lagascalia* **4**: 153-211.
- Löve, Á. and Löve, D. 1956. Cytotaxonomical conspectus of the Icelandic flora. *Acta Horti Gotoburgensis* **20**: 65-291.
- Mabberley, D. J. 1997. The plant book, 2 nd. ed. p. 611. Cambridge University Press, Cambridge.
- Májovsky, J. et al. 1970. Index of chromosome numbers of Slovakian flora part 2. *Acta Fac. Rerum Nat. Univ. Comenianae Bot.* **18**: 45-60.
- Májovsky, J. et al. 1976. Index of chromosome numbers of Slovakian flora (part 5). *Acta Fac. Rerum Nat. Univ. Comenianae Bot.* **25**: 1-18.
- Markova, M. L. and Thu, N. T. 1974. Löve, Á. (ed.). IOPB chromosome number reports XLIII. *Taxon* **23**: 193-194.
- Morton, J. K. 1973. A cytological study of the British Labiatae(excluding *Mentha*). *Watsonia* **9**: 239-246.
- Murata, G. 2003. Labiatae. In Shimizu, T. (ed.) *Naturalized plants of Japan*. pp.170-175. pl.

- 80-83. Heibonsha, Tokyo. (in Japanese)
- Murata, G. and Yamazaki, 1993. *Lamium* L. In Iwatsuki, K., Yamazaki, T., Boufford, D. E. and Ohba, H. (eds.). *Flora of Japan IIIa*. pp. 294-296. Kodansha, Tokyo.
- Nishikawa, T. 1981. Chromosome counts of flowering plants of Hokkaido (5). Rep. Taisetsuzan Inst. Sci. **16**: 45-53. (in Japanese with English summary)
- Ortega-Olivencia, A. and Ruíz-Tellez, T. 1990. Números cromosómicos de plantas occidentales, 551-555. *Anales Jard. Bot. Madrid* **47**: 175-177.
- Pólya, L. 1949. Fedorov, A.A. 1969. Chromosome numbers of flowering plants. p. 364. Komar. Bot. Inst. Acad. Sci. U.S.S.R., Leningrad. (in Russian)
- Rudyka, E. G. 1995. Chromosome numbers in vascular plants from the southern part of the Russian Far East. *Bot. Zhurn.* **80**: 87-90.
- Saggo, M. I. S. and Bir, S. S. 1983 a. Cytological studies on Indian members of Acanthaceae and Labiatae. *J. Palynol.* **19**: 243-277.
- Saggo, M. I. S. and Bir, S. S. 1983 b. Bir, S. S. (ed.). SOCGI plant chromosome number reports-I. *J. Cytol. Genet.* **18**: 61-63.
- Saggo, M. I. S. and Bir, S. S. 1986. Meiotic studies on some East Himalayan members of family Labiatae. *J. Indian Bot. Soc.* **65**: 304-309.
- Singh T. P. 1995. Alterations in the basic chromosome numbers as a means of speciation in Labiatae. *Feddes Repert. Berlin* **106**: 39-47.
- Skalinska, M., Pogan, E. and Jankun, A. et al. 1968. Further studies in chromosome numbers of Polish angiosperms seventh contribution. *Acta Biol. Cracov. Ser. Bot.* **11**: 199-224. pl. 27-29.
- Sokolovskaya, A. P., Probatova, N. S. and Rudyka, E. G. 1986. A contribution to the study of chromosome numbers and geographical distribution of some species of the family Lamiaceae in the Soviet Far East. *Bot. Zhurn.* **71**: 195-200.
- Sorsa, V. 1963. Chromosomenzahlen finnischer Kormophyten. II. *Ann. Acad. Sci. Fennicae, Ser. A, IV. Biol.* **68**: 1-14.
- Starodubtsev, V. N. 1985. Chromosome numbers in the representatives of some families from the Soviet Far East. *Bot. Zhurn.* **70**: 275-277.
- Strid, A. 1965. Studies in the Aegean flora. VI. Notes on some genera of Labiatae. *Bot. Notiser* **118**: 104-122.
- Strid, A. and Franzén R. 1981. Löve, Á. (ed.). Chromosome number reports LXXIII. *Taxon* **30**: 829-842.
- Tanaka, R. 1974. Organizational system of meiotic division and the development of reproductive cells in higher plants. *The Cell* **6**: 22-25. (in Japanese)
- Terasaka O. and Tanaka, R. 1974. Cytological studies on the nuclear differentiation in microspore division of some angiosperms. *Bot. Mag. Tokyo* **87**: 209-207.
- Tischler, G. 1934. Die Bedeutung der Polyploidie für die Verbreitung der Angiospermen, erläutert an den Arten Schleswig-Holsteins, mit Ausblicken auf andere Florengebiete. *Bot. Jahrb.* **67**: 1-36.
- Tischler, G. 1937. Die Halligenflora der Nordsee im Lichte cytologischer Forschung. *Cytologia Fujii Jub.* pp. 162-170.
- Van Den Brand, C. Van Meel, F. C. M. and Wieffering J. H. 1979. Löve, Á. (ed.). IOPB chromosome number reports LXIV. *Taxon* **28**: 395-397.
- (Received June 7, 2005; accepted December 6, 2005)
- 三浦憲人¹・岩坪美兼²・鳴橋直弘²: 日本産オドリコソウ属 5 種の染色体数
- オドリコソウ属のオドリコソウ (*L. album* var. *barbatum*), ホトケノザ (*L. amplexicaule*), ヤマジオウ (*L. humile*), モミジバヒメオドリコソウ (*L. hybridum*), それにヒメオドリコソウ (*L. purpureum*) の染色体数を調べた。オドリコソウは $2n=18$, ホトケノザは $2n=18$, ヤマジオウは $2n=34$, モミジバヒメオドリコソウは $2n=36$, そしてヒメオドリコソウは $2n=18$ であった。この属の染色体基本数は $x=9$ が知られていることから, オドリコソウとホトケノザは二倍体, モミジバヒメオドリコソウは四倍体であり, それらはいずれも過去の報告と一致した。ヤマジオウの染色体数は本研究により $2n=34$ であることが初めて明らかにされ, 染色体基本数は $x=17$ と判断された。
- (¹〒321-1661 日光市中宮祠 2480-1; ²〒930-8555 富山市五福 3190 富山大学理学部生物学科)