

A Fundamental study on analysis of ECT imaging using model animals of psycho-neuro disorders

メタデータ	言語: jpn 出版者: 公開日: 2022-04-14 キーワード (Ja): キーワード (En): 作成者: Mori, Hirofumi メールアドレス: 所属:
URL	https://doi.org/10.24517/00056880

This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 International License.



1992 Fiscal Year Final Research Report Summary

A Fundamental study on analysis of ECT imaging using model animals of psycho-neuro disorders

Research Project

Project/Area Number

03670542

Research Category

Grant-in-Aid for General Scientific Research (C)

Allocation Type

Single-year Grants

Research Field

Radiation science

Research Institution

Kanazawa University

Principal Investigator

MORI Hirofumi Kanazawa Univ., Radioisotope Center, Professor, アイソトープ総合センター, 教授 (90019604)

Co-Investigator(Kenkyū-buntansha)

KIDO Hideki Kanazawa Univ., School of Medicine, Assist. Prof., 医学部付属病院, 講師 (50161518)

MATSUDA Hiroshi Kanazawa Univ., School of Medicine, Assist. Prof., 医学部付属病院, 講師 (90173848)

SHIBA Kazuhiro Kanazawa Univ., Radioisotope Center, Assistant, アイソトープ総合センター, 助手 (40143929)

Project Period (FY)

1991 – 1992

Keywords

Emission CT / receptor / autoradiography / animal model / schizophrenia / antipsychotic drugs / radiopharmaceuticals

Research Abstract

The effects of acute intravenous or chronic subcutaneous haloperidol (HPD) administration on the local cerebral blood flow (LCBF) in 41 regions of the rat brain were studied by means of the quantitative autoradiographic N-isopropyl-p-[125-I] iodoamphetamine technique. After acute HPD administrations, the LCBFs decreased in the anterior cingulate cortex, primary motor cortex and primary auditory cortex, and increased the n. habenula. Chronic HPD administration reduced LCBF in the substantia nigra, and increased LCBFs in the n.caudatus-putamen, n.accumbens and n.habenula. No change in LCBF

was observed in any cortical region after chronic HPD administratin.

An in vivo receptor binding techniques was applied to evaluate the affinities of and clozapine, risperidone, RMI-81582 and haloperidol for dopamine D1, D2, 5- HT2 receptors in the rat brain with [3-H] -SCH23390, [3-H] -YM-09151-2 and [3-H] -Ketanserine as selective ligands. Time course study of receptor occupancy at 25 to 250 min after i ... More

Research Products (6 results)

All Other

All Publications (6 results)

[Publications] Shiba K.: "Synthesis of tecnetium-99m labeled diaminodithiol for bifunctionl chelating agents" Int. J. Radiat. Appl. Instrum. part A.42. 1159-1164 (1991) ▾

[Publications] Shiba K.: "Synthesis and characterization of Tc-99m-p-halophenethyl diaminodithiol analogs" Nucl. Med. Communications. 19. 303-310 (1992) ▾

[Publications] Sumiyosi T.: "Time Course of Dopamin-1,2 and Serotonin-2 Peceptor Occupancy Rates by Haloperidol and Clozapin in Vivo" Japan. J. Psychiat. Neurol.(1993) ▾

[Publications] Shiba K.: "Shnthesis of technetium-99m labeled diaminoditiol for bifunctional chelating agents" Appl. Radiat. Isot..42(12). 1159-1164 (1991) ▾

[Publications] Shiba K.: "Technetium-99m p-iodophenethyl(DADT-IPE); Potential brain perfusion imaging agent for SPECT" Nucl. Med. Biol.19(3). 303-310 (1992) ▾

[Publications] Sumiyosi T.: "Time course of dopamine-2 and serotonine-2 receptor occupancy rates by haloperidol and clozapine in vivo" Jap. J. Psychiat. Neurol.

URL: https://kaken.nii.ac.jp/report/KAKENHI-PROJECT-03670542/036705421992kenkyu_seika_hokoku

Published: 1994-03-23