

Study on Nursing care for Dementia

メタデータ	言語: jpn 出版者: 公開日: 2022-11-11 キーワード (Ja): キーワード (En): 作成者: Kanagawa, Katsuko メールアドレス: 所属:
URL	https://doi.org/10.24517/00067910

This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 International License.



1987 Fiscal Year Final Research Report Summary

Study on Nursing care for Dementia

Research Project

Project/Area Number

61571121

Research Category

Grant-in-Aid for General Scientific Research (C)

Allocation Type

Single-year Grants

Research Field

Nursing

Research Institution

Kanazawa University

Principal Investigator

KANAGAWA Katsuko School of Allied Medical Professions,Kanazawa University, 医療技術短期大学部, 教授 (10019565)

Co-Investigator(Kenkyū-buntansha)

KAWASHIMA Kazuyo School of Allied Medical Professions,Kanazawa University, 医療技術短期大学部, 助手 (40157855)

IZUMI Kiyoko School of Allied Medical Professions,Kanazawa University, 医療技術短期大学部, 講師 (20115207)

ONO Turuko School of Allied Medical Professions,Kanazawa University, 医療技術短期大学部, 助教授 (10020025)

HAGINO Taeko School of Allied Medical Professions,Kanazawa University, 医療技術短期大学部, 助教授 (90115262)

Project Period (FY)

1986 – 1987

Keywords

dementia / nursing home / dementia in home / abnormal behavior / communication disorder / 看護ケア

Research Abstract

We investigated the abnormal behavior and communication disorders of dementia in institution elderly and home elderly.

Subjects were a total of 248 cases (61 males and 187 females) in Y nursing home, and a total of 30 cases (8 males and 22 females) in homes.

1. The results were as follows in institution elderly.

1) 32% of institutional elderly were classified into Dementia(0-10 points in scales) by Hasegawa Dementia Scale, 26% in males and 33% in females.

2) Incidence of dementia became higher with increasing of ages.

3) There were relation between degree of Dementia and ADL of elderly.

2. The results were as follows for 30 dementia (8 males and 22 females) in the home.

1) Over 80% of families with dementia complained the abnormal behavior and communication disorders of them.

2) Half of dementia had the effective method of relieving anxiety and strain.

We felt that we must offer not only nursing care to the each disorder, but also total care to dementia.

Also,we must search dementia how into throw ourselves into dementias' feelings.

Research Products (8 results)

All Other

All Publications (8 results)

[Publications] 川島和代: 金沢大学医療技術短期大学部紀要. 11. 63-68 (1987)



[Publications] 金川克子: 老年社会科学. 10. (1988)



[Publications] 川島和代: 老年社会科学. 10. (1988)



[Publications] 金川克子: 老年社会科学. 11. (1989)



[Publications] KAWASIMA,Kazuyo: "The Study on Factors Relating to Health Behavior of the Health Aged." Memoirs of the school of allied medical Professions,Kanazawa university.11. 63-68 (1987)



[Publications] KANAGAWA,Katsuko: "The Problems of Health Care in A Nursing Home." Japanese journal of gerontology. 10. (1988)



[Publications] KAWASIMA,Kazuyo: "Relationship between Degree of Demential and ADL." Japanese journal of gerontology. 10. (1988)



[Publications] KANAGAWA,KATSUKO: "Study on ADL of Dementia and Correspondence of Families" Japanese journal of gerontology. 11. (1989)



URL: https://kaken.nii.ac.jp/report/KAKENHI-PROJECT-61571121/615711211987kenkyu_seika_hokoku_

Published: 1989-03-29