

Effective Dissolution of Biomass in Ionic Liquids by Irradiation of Non-thermal Atmospheric Pressure Plasma

著者	Kuroda Kosuke, Shimomura Kai, Ishijima Tatsuo, Takada Kenji, Ninomiya Kazuaki, Takahashi Kenji
journal or publication title	Australian Journal of Chemistry
volume	70
number	6
page range	731-734
year	2016-01-01
URL	http://hdl.handle.net/2297/46338

doi: <https://doi.org/10.1071/ch16554>

Supplementary Materials

Effective Dissolution of Biomass in Ionic Liquids by Irradiation of Non-thermal Atmospheric Pressure Plasma

Kosuke Kuroda^A, Kai Shimomura^A, Tatsuo Ishijima^B, Kenji Takada^A, Kazuaki Ninomiya^C, and Kenji Takahashi^A

^A*Division of Natural System, Graduate School of Natural Science and Technology, Kanazawa University, Kakuma-machi, Kanazawa 920-1192, Japan*

^B*Research Center for Sustainable Energy & Technology, Institute of Science and Engineering, Kanazawa University, Kakuma-machi, Kanazawa 920-1192, Japan*

^C*Institute for Frontier Science Initiative, Kanazawa University, Kakuma-machi, Kanazawa 920-1192, Japan*

Fig. S1 shows the molecular weight distribution of lignin after heating at 50 °C for 4h in an oil bath. There was no change of molecular weight distribution before and after heating. It supports that plasma irradiation decomposed the lignin structure.




Fig. S1. Molecular weight distribution of lignin after heating at 50 °C for 4h in an oil bath.

Fig. S2 shows that ^1H NMR spectra of [Emim]OAc before and after plasma irradiation for 4h, and Table S1 shows the integral values of the peaks. There is no change in both spectra and the integral values after plasma irradiation for 4h. The results show that [Emim]OAc was not decomposed during plasma irradiation.




Fig. S2. ^1H NMR spectra of [Emim]OAc before and after plasma irradiation for 4h.

Table S1. Integral values of the peaks of the protons in ^1H NMR spectra of [Emim]OAc before and after plasma irradiation for 4h. The numbers of protons correspond to the numbers shown in Figure S2.

	Integral value					
	H_1 (3H)	H_2 (1H)	H_3 (2H)	H_4 (3H)	H_5 (2H)	H_6 (3H)
Before plasma irradiation	3.00	0.96	1.98	2.86	1.81	2.88
After plasma irradiation	3.00	0.96	2.01	2.92	1.79	2.95