

Differential expression of CD45RO (UCHL1) and its functional relevance in two subpopulations of circulating TCR- / + lymphocytes

| | |
|------------------------------|--|
| 著者 | Miyawaki Toshio, Kasahara Yoshihito, Taga Kazuyuki, Yachie Akihiro, Taniguchi Noboru |
| journal or publication title | Journal of Experimental Medicine |
| volume | 171 |
| number | 5 |
| page range | 1833-1838 |
| year | 1990-01-01 |
| URL | http://hdl.handle.net/2297/7446 |

doi: 10.1084/jem.171.5.1833

DIFFERENTIAL EXPRESSION OF CD45RO (UCHL1) AND
ITS FUNCTIONAL RELEVANCE IN TWO SUBPOPULATIONS
OF CIRCULATING TCR- γ/δ^+ LYMPHOCYTES

By TOSHIO MIYAWAKI, YOSHIHITO KASAHARA, KAZUYUKI TAGA,
AKIHIRO YACHIE, AND NOBORU TANIGUCHI

*From the Department of Pediatrics, School of Medicine, Kanazawa University,
Kanazawa, Ishikawa 920, Japan*

Human T cells express various isoforms of CD45 (leukocyte common antigen), which arise from alternate mRNA splicing (1). The mAb UCHL1 recognizes a 180 kD molecular mass isoform, CD45RO (2). Recent studies on major two ($CD4^+$ and $CD8^+$) subpopulations of circulating T cells have suggested that UCHL1 identifies a primed population of T cells that includes memory cells (3). Within $CD4^+$ cells, the CD45RO population mediates proliferative responses to recall antigens and is able to support B cell differentiation. A reciprocal population of $CD4^+$ cells, identified by mAbs to the 205- and 220-kD molecular mass isoforms termed CD45RA, do not show these capabilities. It has also been shown that the CD45RO population of $CD8^+$ cells contains precursors exerting memory-dependent cytotoxicity (4).

While the majority of mature T cells express the TCR- α/β -CD3 complex, a minor population of blood T cells lacking both CD4 and CD8 bear a distinct TCR composed of γ and δ subunits. Although the function and the repertoire of TCR- γ/δ remain to be fully defined, it has been shown that blood TCR- γ/δ^+ T cells can be subdivided largely into two nonoverlapping subsets defined by BB3 and δ TCS1 mAb (5, 6). In this paper we examined the expression of CD45RO by TCR- γ/δ^+ in healthy children and adults to elucidate some of their biological roles. We will show here that CD45RO is preferentially expressed on BB3 $^+$ subsets among circulating TCR- γ/δ^+ cells. BB3 $^+$ cells appear to represent a population of T γ/δ^+ cells capable of responding to some antigens.

Materials and Methods

Blood Samples. Cord blood was collected from the umbilical vein immediately after the uneventful delivery of full-term newborn babies. Venous blood from healthy infants and children of different ages was obtained after the informed parental consent was given. Adult subjects were healthy volunteers between the ages of 25 and 35 yr.

Monoclonal Antibodies. The mAb UCHL1 (IgG2a) against CD45RO (2) was purchased from Dakopatts a/s (Copenhagen, Denmark). The BB3 mAb (IgG1) specific for human TCR- γ

This work was supported by Grant-in-Aid for Scientific Research from the Ministry of Education, Science and Culture in Japan and by a grant from the Ministry of Welfare in Japan.

Address correspondence to Dr. Toshio Miyawaki, Department of Pediatrics, School of Medicine, Kanazawa University, 13-1 Takara-Machi, Kanazawa, Ishikawa 920, Japan.

chain (5) was the kind gift of Dr. L. Moretta (University of Geneva, Italy). Anti-Ti γ A mAb (IgG2a), generously provided by Dr. T. Hercend (Institut Gustave-Roussy, Villejuif, France), recognizes the human TCR- γ molecule (6). FITC-conjugated TCR δ 1 mAb (IgG1) (7) and FITC-conjugated δ TCS1 mAb (IgG1) (8), both of which have been found to be directed to human TCR- δ chains, were obtained from T Cell Sciences (Cambridge, MA). FITC-conjugated anti-Leu-3a mAb (IgG1) specific for CD4 was from Becton Dickinson & Co. (Mountain View, CA).

Immunofluorescence Analysis. Viable mononuclear cells from heparinized blood were isolated on Ficoll/Hypaque gradients (Lymphoprep, Nycomed AS, Oslo, Norway) and stained with mAb. Two-color staining was performed to evaluate expression of CD45RO by TCR- γ/δ^+ cells. The cells were first incubated with UCHL1 or isotype-matched control mouse IgG, and counterstained with biotinylated rabbit anti-mouse IgG2a antibody (Zymed Laboratories, San Francisco, CA), and further with phycoerythrin (PE)-conjugated streptavidin (Becton Dickinson & Co.). Next, the cells were stained for determination of TCR- γ/δ^+ cells with FITC-conjugated TCR δ 1, FITC-conjugated δ TCS1, or unconjugated BB3 mAb. For BB3 staining, the cells were finally treated with FITC-conjugated horse anti-mouse IgG1 antibody (Zymed Laboratories). The labeled cells were analyzed by an Epics C flow cytometer (Coulter Electronics, Inc., Hialeah, FL).

Proliferative Responses. TCR- γ/δ^+ cell (Ti γ A $^+$ and δ TCS1 $^+$) subsets were isolated using an Epics C flow cytometer from mononuclear cells stained with the corresponding mAbs as above. These cells were seeded on flat-bottomed microtiter plates (Corning Glass Works, Corning, NY) at 5×10^4 cells/well in 0.2 ml of RPMI 1640 supplemented with 10% AB serum in the presence of 5×10^3 irradiated adherent monocytes from the same donor. Cultures were stimulated with 1 μ g/ml of purified protein derivative (PPD; Japan BCG Co., Ltd., Tokyo, Japan) or 10 ng/ml of anti-CD3 mAb (OKT3; Ortho Pharmaceutical Co., Raritan, NJ) for 120 h and were pulsed with 0.5 μ Ci of [3 H]TdR 12 h before harvesting. As controls, the similar cultures were carried out for CD45RO $^+$ and CD45RO $^-$ CD4 $^+$ cells separated by the cell sorting.

Results and Discussion

We used TCR δ 1 mAb to examine the frequency of TCR- γ/δ^+ cells and their expression of CD45RO, since TCR δ 1 has been shown to detect the whole population of TCR- γ/δ^+ cells (6). The appearance of blood TCR- γ/δ^+ cells was scarce (<0.5%) at birth, gradually increased with age, and reached the adult values in older children (Table I). On the other hand, the proportions of TCR- α/β^+ cells with CD4 or CD8 in the neonatal blood are similar to those observed in the adult blood (9). Consistent

TABLE I
Frequency of TCR δ 1 $^+$ Cells and Their Expression of CD45RO in Peripheral Blood of Different Ages

| Age | n | Percentage of TCR δ 1 $^+$ cells | Percent CD45RO expression by TCR δ 1 $^+$ cells |
|---------|----|---|--|
| Newborn | 6 | <0.5 | ND |
| 0-1 mo | 7 | 1.5 \pm 0.2* | 72.5 \pm 4.6 |
| 2-3 mo | 8 | 1.9 \pm 0.3 | 63.8 \pm 3.3 |
| 4-11 mo | 6 | 2.3 \pm 0.4 | 61.0 \pm 5.1 |
| 1-2 yr | 12 | 2.7 \pm 0.5 | 69.6 \pm 4.0 |
| 3-6 yr | 14 | 3.9 \pm 0.4 | 72.4 \pm 3.5 |
| 7-15 yr | 16 | 4.4 \pm 0.6 | 69.3 \pm 3.1 |
| Adult | 12 | 5.8 \pm 1.1 | 72.4 \pm 3.7 |

* Means \pm SEM.

with the current concept of naive and memory T cells, we observed that the circulating pool of CD4⁺ and CD8⁺ cells expressing CD45RO increased as a function of age (unpublished observations). In adult subjects, the mean percentages of CD45RO⁺ cells within CD4⁺ and CD8⁺ cells are 35 and 25%, respectively. In marked contrast to CD4⁺ and CD8⁺ cells, expression of CD45RO was observed on a sizable portion (~70%) of blood TCR- γ/δ ⁺ cells in all donors examined (Table I). The remarkable finding was that most TCR- γ/δ ⁺ cells possessed CD45RO on their surfaces even around 1 mo of life shortly after birth.

We often observed that some of donors exhibited a clear bimodal distribution of CD45RO expression on TCR- γ/δ ⁺ cells (Fig. 1 A). This suggests the presence of a small CD45RO⁻ population of TCR- γ/δ ⁺ cells in blood. The recent availability of other mAb specific for human TCR- γ/δ molecules have allowed to identify two mutually exclusive subsets defined by BB3 and δ TCS1 mAbs (5, 6). In blood TCR- γ/δ ⁺ cells, most cells reacts with BB3, whereas a few cells are positive for δ TCS1. Based on our observations, we next asked whether expression of CD45RO by these two TCR- γ/δ ⁺ cells might be different from each other. Fig. 1 shows a representative of two-color immunofluorescence analysis for CD45RO expression by T γ/δ ⁺ subsets. Interestingly, a large percentage of BB3⁺ cells were considered CD45RO⁺, whereas CD45RO expression was absent from δ TCS1⁺ cells. As presented in Table II, this was the rule in all of seven subjects studied.




FIGURE 1. Two-color immunofluorescence analysis of CD45RO expression by TCR δ 1⁺, BB3⁺ and δ TCS1⁺ cells. Contour maps (A, C, and E) show two-color staining patterns of lymphocytes gated by forward and 90° light scatters. A second gate was set to include the whole populations of TCR δ 1⁺ (B), BB3⁺ (D) and δ TCS1⁺ cells (F) and 5,000 cells accumulating within the gate were analyzed for CD45RO expression, which are expressed as the histogram patterns. A vertical dashed line in the histograms indicates the negative control.

TABLE II
Expression of CD45RO by TCR- γ/δ^+ Subsets

| Subjects | Percent CD45RO Expression by | | |
|----------|--|--|---|
| | TCR $\delta 1^+$ Cells | BB3 $^+$ Cells | δ TCS1 $^+$ Cells |
| NT | 50.1 (4.6)* | 92.1 (3.5) | 1.4 (1.6) |
| NY | 84.5 (6.6) | 94.0 (5.8) | ND (<0.5) |
| TT | 82.6 (6.6) | 93.4 (5.1) | 0.9 (0.9) |
| NK | 62.3 (6.2) | 97.2 (4.3) | 0.8 (0.9) |
| AY | 86.0 (12.0) | 91.6 (11.2) | ND (<0.5) |
| YH | 92.0 (4.5) | 92.0 (1.2) | 2.6 (1.4) |
| YK | 38.2 (11.0) | 92.0 (7.8) | 1.7 (3.8) |
| | 61.8 \pm 8.9 \ddagger (7.4 \pm 2.8) | 93.2 \pm 0.7 \ddagger (5.6 \pm 1.2) | 1.5 \pm 0.3 \ddagger (1.7 \pm 0.5) |

CD45RO expression by each TCR- γ/δ^+ subset was examined for the blood from adult subjects.

* Parentheses indicate the proportions of each subset in lymphocytes.

\ddagger Mean \pm SEM.

Although studies for established clones suggest that all TCR- γ/δ^+ cell clones mediate non-MHC-restricted cytotoxicity (10), the antigen-specificity and immunological functions of these cells remain speculative. On the basis of the correlation between CD45RO expression and memory T cells, it seemed feasible to suppose that BB3 $^+$ cells, but not δ TCS1 $^+$ cells lacking CD45RO, might contain a population responding to certain antigens in a similar manner to memory CD4 $^+$ cells. Preliminary experiments suggested that staining with BB3 mAb for the cell sorting resulted in somewhat enhanced background responses of isolated cells, indicating the activation of corresponding subsets by this antibody (11). It has been demonstrated that most BB3 $^+$ cells are identified by anti-Ti γ A mAbs as well (12). In addition, resting Ti γ A cells do not proliferate in response to anti-Ti γ A mAb in the absence of exogenous IL-2 (13). Thus, we substituted anti-Ti γ A for BB3 to isolate CD45RO $^+$ TCR- γ/δ^+ subsets. As shown in Table III, both Ti γ A $^+$ and δ TCS1 $^+$ cells proliferated well in response to anti-CD3 mAb. In PPD-reactive individuals, appreciable proliferative responses to PPD, though less than those seen in memory (CD45RO $^+$) CD4 $^+$ cells, were obtained only from Ti γ A $^+$ cells, but not from δ TCS1 $^+$ cells. These results strongly suggest that a fraction of Ti γ A $^+$ cells may contribute the immune response to a nominal antigen. Since 95% of BB3 cells, though not completely identical, coexpress Ti γ A antigen (13), we speculate that functional profiles of Ti γ A $^+$ cells may represent those of BB3 $^+$ cells. Regarding the antigen-specific response of TCR- γ/δ^+ cells, it has recently been reported that TCR- γ/δ^+ clones established from blood lymphocytes respond to some antigens such as tetanus toxoid or PPD in a MHC-restricted fashion (14, 15).

Moretta et al. (16) have indicated that BB3 $^+$ cells express C γ 1-encoded TCR- γ/δ heterodimers, whereas δ TCS1 $^+$ cells express C γ 2-encoded molecules. BB3 $^+$ cells predominate in the peripheral blood, but are very scantily found in the thymic tissues. In contrast, δ TCS1 $^+$ cells are enriched in the thymus. Immunohistochemical analysis have shown that δ TCS1 $^+$ cells are predominant in intraepithelial lymphocytes of the jejunum as compared with the blood (17). However, the reason for such different tissue distribution between BB3 $^+$ and δ TCS1 $^+$ cells is unknown.

TABLE III
*Proliferative Responses of TCR- γ/δ^+ and CD4 $^+$ Subpopulations
 by PPD or Anti-CD3 mAb*

| Donors | Stimuli | TCR- γ/δ^+ Cells | | CD4 $^+$ Cells | |
|--------|----------|------------------------------|---------------------|----------------|-------------|
| | | Ti γ A $^+$ | δ -TCS1 $^+$ | CD45RO $^+$ | CD45RO $^-$ |
| YH | None | 266 | 501 | 562 | 722 |
| | PPD | 2,305 | 488 | 24,968 | 902 |
| | Anti-CD3 | 31,600 | 28,098 | 53,338 | 48,221 |
| YK | None | 320 | 341 | 982 | 1,102 |
| | PPD | 14,614 | 256 | 34,682 | 1,490 |
| | Anti-CD3 | 29,646 | 12,660 | 63,052 | 38,962 |
| NT | None | 664 | 125 | 902 | 562 |
| | PPD | 8,420 | 102 | 23,315 | 875 |
| | Anti-CD3 | 23,528 | 32,628 | 100,192 | 48,802 |

Each subpopulation was isolated from PPD-reactive adult donors. The cells (5×10^4 /well) were cultured with PPD (1 μ g/ml) or anti-CD3 mAb (10 ng/ml) in the presence of autologous irradiated monocytes for 5 d. The results indicate the mean [3 H]TdR incorporation of triplicate cultures.

In conclusion, the present study showed that the BB3 $^+$ subset in the peripheral blood selectively expressed CD45RO, corresponding with their responding capability to recall antigens. These observations implicate a special role of this subset in the immune response to many pathogens and infectious agents.

Summary

We examined the developmental profile of TCR- γ/δ^+ cells with respect to CD45RO expression. Although total TCR- γ/δ^+ cells were negligible in the neonatal blood and increased with advancing age, most blood TCR- γ/δ^+ cells markedly expressed CD45RO without a distinction of age, probably reflecting a different CD45RO expression of two subsets defined by BB3 and δ TCS1 mAbs. The vast majority of BB3 $^+$ cells expressed CD45RO, whereas expression of CD45RO was virtually absent in the δ TCS1 $^+$ population. Functional studies revealed that, while both TCR- γ/δ^+ cell subsets showed CD3-mediated activation, only BB3 $^+$ (or Ti γ A $^+$) cells, but not δ TCS1 $^+$ cells, appeared to proliferate in response to PPD in PPD-reactive individuals. The results suggested that the CD45RO $^+$ (BB3 $^+$ or Ti γ A $^+$) subset among blood TCR- γ/δ^+ cells may be mainly involved in the memory or primed component of the immune system responding to some foreign antigens.

We thank Dr. L. Moretta for providing BB3 mAb and for Dr. T. Hercend for kind supply of anti-Ti γ A mAb.

Received for publication 29 November 1989 and in revised form 29 January 1990.

References

1. Thomas, M. L., and L. Lefrançois. 1988. Differential expression of the leukocyte-common antigen family. *Immunol. Today*. 9:320.

2. Terry, L. A., M. H. Brown, and P. C. L. Beverley. 1988. The monoclonal antibody, UCHL1, recognizes a 180,000 MW component of the human leukocyte-common antigen, CD45. *Immunology*. 64:331.
3. Sanders, M. E., M. W. Makgoba, and S. Shaw. 1988. Human naive and memory T cells: reinterpretation of helper-inducer and suppressor-inducer subsets. *Immunol. Today*. 9:195.
4. Merckenschlager, M., and P. C. L. Beverley. 1989. Evidence for differential expression of CD45 isoforms by precursors for memory-dependent and independent cytotoxic responses: human CD8 memory CTLp selectively express CD45RO (UCHL1). *Int. Immunol.* 1:450.
5. Bottino, C., G. Tambussi, S. Ferrini, E. Ciccone, P. Varese, M. C. Mingarri, L. Moretta, and A. Moretta. 1988. Two subsets of human T lymphocytes expressing γ/δ antigen receptor are identifiable by monoclonal antibodies directed to two distinct molecular forms of the receptor. *J. Exp. Med.* 168:491.
6. Faure, F., S. Jitsukawa, F. Triebel, and T. Hercend. 1988. Characterization of human peripheral blood lymphocytes expressing the CD3- γ/δ complex with anti-receptor monoclonal antibodies. *J. Immunol.* 141:3357.
7. Band, H. F. Hochstenbach, J. Mclean, S. Hata, M. S. Krangel, M. B. Brenner. 1987. Immunochemical proof that a novel rearranging gene encodes the T cell receptor δ subunit. *Science (Wash. DC)*. 238:682.
8. Wu, Y.-J., W.-T. Tian, R. M. Snider, C. Rittershaus, P. Rogers, L. LaManna, and S. H. Ip. 1988. Signal transduction of γ/δ T cell antigen receptor with a novel mitogenic anti- δ antibody. *J. Immunol.* 141:1476.
9. Yachie, A., T. Miyawaki, T. Nagaoki, T. Yokoi, M. Mukai, N. Uwadana, and N. Taniguchi. 1981. Regulation of B cell differentiation by T cell subsets defined with monoclonal OKT4 and OKT8 antibodies in human cord blood. *J. Immunol.* 127:1314.
10. Hercend, T., and F. Triebel. 1989. Subpopulations of human peripheral T gamma delta lymphocytes. *Immunol. Today*. 10:186.
11. Ciccone, E., S. Ferrini, C. Bottino, O. Viale, I. Prigione, G. Pantaleo, G. Tambussi, A. Moretta, and L. Moretta. 1988. A monoclonal antibody specific for a common determinant of the human T cell receptor γ/δ directly activates CD3⁺WT31⁻ lymphocytes to express their functional program(s). *J. Exp. Med.* 168:1.
12. Triebel, F., F. Faure, F. Mami-Chouaib, S. Jitsukawa, A. Griscelli, C. Genevée, S. Roman-Roman, and T. Hercend. 1988. A novel human V δ gene expressed predominantly in the T γ A fraction of γ/δ ⁺ peripheral lymphocytes. *Eur. J. Immunol.* 18:2021.
13. Faure, F., S. Jitsukawa, F. Tribel, and T. Hercend. 1988. CD3/T γ A: a functional γ -receptor complex expressed on human peripheral lymphocytes. *J. Immunol.* 140:1372.
14. Kozbor, D., G. Trinchieri, D. S. Monos, M. Isobe, G. Russo, J. A. Haney, C. Zmijewski, and C. M. Croce. 1989. Human TCR- γ/δ ⁺, CD8⁺ T lymphocytes recognize tetanus toxoid in an MHC-restricted fashion. *J. Exp. Med.* 169:1847.
15. Haregewoin, A., G. Soman, R. C. Hom, and R. W. Finberg. 1989. Human $\gamma\delta$ ⁺ T cells respond to mycobacterial heat-shock protein. *Nature (Lond.)*. 340:309.
16. Moretta, L., E. Ciccone, M. C. Mingarri, C. Bottino, S. Ferrini, G. Tambussi, G. Melioli, C. E. Grossi, and A. Moretta. 1989. Human T lymphocytes expressing γ/δ T cell antigen receptor. *Clin. Immunol. Immunopathol.* 50:S117.
17. Kyes, S., E. Carew, S. R. Carding, C. A. Janeway, Jr., and A. Hayday. 1989. Diversity in T-cell receptor γ gene usage in intestinal epithelium. *Proc. Natl. Acad. Sci. USA*. 86:5527.