

Fat-free mass and calf circumference as body composition indices to determine non-exercise activity thermogenesis in patients with diabetes

著者	Isobe Yuki, Sakurai Masaru, Kita Yuki, Takeshita Yumie, Misu Hirofumi, Kaneko Shuichi, Takamura Toshinari
journal or publication title	Journal of Diabetes Investigation
volume	7
number	3
page range	352-358
year	2016-05-01
URL	http://hdl.handle.net/2297/43921

doi: 10.1111/jdi.12421

Supplementary Table 1. Clinical anthropometry and biochemical characteristics of the study subjects

	All	Males	Females	<i>p</i>
Age (year)	52.5±15.6	53.5±13.9	51.3±17.7	0.669
Bodyweight (kg)	70.8±24.2	67.9±17.4	74.0±30.2	0.435
Body mass index (kg/m ²)	26.81±9.14	23.48±4.89	30.49±11.28	0.013
Fat mass (kg)	25.60±19.82	16.27±9.09	35.91±23.36	0.001
Fat-free mass (kg)	45.24±10.89	51.66±9.07	38.15±8.02	0.000
Waist circumference (cm)	93.58±18.66	87.39±12.73	100.42±21.92	0.025
Hip circumference (cm)	96.98±14.58	92.43±7.65	101.99±18.57	0.037
Thigh circumference (cm)	52.07±10.53	48.05±6.25	56.52±12.54	0.009
Calf circumference (cm)	36.37±6.40	35.14±5.37	37.48±7.28	0.416
Arm muscle area (cm ²)	50.82±18.64	49.16±11.69	52.66±24.38	0.560
Basal energy expenditure (kcal/day)	1508±349	1513±249	1502±442	0.921
Basal energy expenditure / body weight	22.16±3.51	22.86±3.15	21.39±3.81	0.193
Diet-induced thermogenesis (kcal/day)	220±140	217±109	223±170	0.884
Diet-induced thermogenesis / body weight	3.15±2.09	3.36±2.04	2.91±2.17	0.498
Total protein (g/dL)	7.0±0.7	7.0±0.7	6.8±0.6	0.343
Total cholesterol (mg/dL)	196±56	185±40	209±69	0.181
Triglyceride (mg/dL)	164 (54-1545)	160 (54-371)	170 (83-1545)	0.707
HDL-cholesterol (mg/dL)	43±14	44±12	42±16	0.550
Aspartate aminotransferase (U/L)	31±21	28±14	34±27	0.397
Alanine aminotransferase (U/L)	46±38	41±25	51±48	0.425
Lactate dehydrogenase (U/L)	176 (139-326)	179 (139-317)	166 (140-326)	0.648
γ-Glutamyl transpeptidase (IU/L)	34 (11-287)	27 (16-287)	39 (11-254)	0.940
Creatinine (mg/dL)	0.75±0.71	0.98±0.93	0.49±0.08	0.028
Immunoreactive insulin (mU/mL)	7.6 (0.9-41.6)	10.9 (0.9-41.6)	7.4 (0.9-25.3)	0.158
Fasting plasma glucose (mg/dL)	173±64	152±57	197±64	0.025
Quantitative insulin sensitivity check index	0.33±0.48	0.34±0.04	0.32±0.06	0.221
HbA1c (%)	9.7±2.2	9.0±2.2	10.6±2.0	0.030
Thyroid-stimulating hormone (mU/mL)	2.00±1.10	2.23±1.08	1.73±1.10	0.159
Free triiodothyronine (pg/mL)	2.45±0.80	1.98±0.78	2.87±0.57	0.016
Free thyroxine (ng/dL)	1.31 (0.90-8.90)	1.24 (0.90-3.29)	1.35 (0.96-8.90)	0.440
Type 1/type 2 diabetes	3/37	1/20	2/17	
Treated with/without antidiabetic agents	23/17	11/10	12/7	
Antidiabetic agents#	5/8/2/9/2/6	1/5/2/2/1/4	4/3/0/7/1/2	

Data are mean±standard deviation, median (range). P-value are Student's t-test for normal distribution, and Mann–Whitney U test for irregular distribution.

insulin/DPP-4 inhibitors/glinide/metformin/a-glucosidase inhibitors/sulfonylureas

Supplementary Table 2. The independent explanatory variables for diet-induced thermogenesis.

	Independent variables	Diet-induced thermogenesis								
		All			Males			Females		
		β	t	<i>p</i>	β	t	<i>p</i>	β	t	<i>p</i>
Model 1	Gender	0.122	0.860	0.396						
	Age	-0.442	-3.296	0.002	-0.213	-0.931	0.365	-0.510	-2.674	0.017
	HbA1c	-0.425	-2.987	0.005	-0.352	-1.539	0.142	-0.374	-1.961	0.069
Model 2	Gender	0.208	1.049	0.302						
	Age	-0.377	-2.208	0.034	-0.240	-0.876	0.394	-0.273	-1.114	0.284
	HbA1c	-0.415	-2.873	0.007	-0.354	-1.502	0.153	-0.298	-1.559	0.141
	Fat-free mass	0.134	0.626	0.536	-0.052	-0.192	0.850	0.367	1.469	0.164
Model 3	Gender	-0.044	-0.308	0.762						
	Age	-0.482	-2.760	0.015	-0.728	-2.566	0.043	-0.345	-1.944	0.100
	HbA1c	-0.473	-3.234	0.006	-0.831	-3.376	0.015	-0.237	-1.478	0.190
	CC	0.207	1.153	0.267	-0.398	-1.464	0.193	0.537	2.858	0.029

Model 1, adjusted for gender, age and HbA1c; Model 2, adjusted for gender, age, HbA1c and fat-free mass; Model 3, adjusted for gender, age, HbA1c and CC.

The multiple regression was used for the analysis.