

Tumor necrosis factor- modifies the effects of Shiga toxin on glial cells

著者	Leu Hue, Sugimoto Naotoshi, Shimizu Masaki, Toma Tomoko, Wada Taizo, Ohta Kunio, Yachie Akihiro
journal or publication title	International Immunopharmacology
volume	38
page range	139-143
year	2016-09-01
URL	http://hdl.handle.net/2297/45911

doi: 10.1016/j.intimp.2016.05.022

Tumor necrosis factor- α modifies the effects of Shiga toxin on glial cells

Hue Leu^{a, b}, Naotoshi Sugimoto^{a, c, *}, Masaki Shimizu^a, Tomoko Toma^a, Taizo Wada^a, Kunio Ohta^a,

Akihiro Yachie^a

^aDepartment of Pediatrics, Graduate School of Medical Science, Kanazawa University, Kanazawa,

Japan

^bDan Phuong General Hospital, Hanoi, Vietnam

^cDepartment of Physiology, Graduate School of Medical Science, Kanazawa University, Kanazawa,

Japan

E-mail addresses

HL: nevergiveup.hue@gmail.com

NS: ns@med.kanazawa-u.ac.jp

MS: shimizum@staff.kanazawa-u.ac.jp

TT: tomoko-t@staff.kanazawa-u.ac.jp

TW: taizo@staff.kanazawa-u.ac.jp

KO: kohta@staff.kanazawa-u.ac.jp

AY: yachie@staff.kanazawa-u.ac.jp

*Corresponding author:

N. Sugimoto, MD, PhD

Department of Pediatrics, Graduate School of Medical Science, Kanazawa University, 13-1

Takara-machi, Kanazawa 920-8640, Japan

E-mail: ns@med.kanazawa-u.ac.jp

Tel: +81-76-265-2313

Fax: +81-76-262-1866

Abstract

Shiga toxin (STX) is one of the main factors inducing hemorrhagic colitis and hemolytic-uremic syndrome (HUS) in infections with STX-producing *Escherichia coli* (STEC). Approximately 62% of patients with HUS showed symptoms of encephalopathy in the 2011 Japanese outbreak of STEC infections. At that time, we reported elevated serum concentrations of tumor necrosis factor (TNF)- α in patients with acute encephalopathy during the HUS phase. In the current study, we investigated whether TNF- α augments the effects of STX in glial cell lines and primary glial cells. We found that TNF- α alone or STX in combination with TNF- α activates nuclear factor- κ B (NF- κ B) signaling and inhibits growth of glial cells. The magnitude of the NF- κ B activation and the inhibition of cell growth by the STX and TNF- α combination was greater than that obtained with TNF- α alone or STX alone. Thus, this in vitro study reveals the role of TNF- α in glial cells during STEC infections.

Keywords

nuclear factor- κ B, Shiga toxin, encephalopathy, outbreak

Abbreviations

BBB: blood-brain barrier

CNS: central nervous system

JNK: c-Jun N-terminal kinase

HUS: hemolytic-uremic syndrome

NF- κ B: nuclear factor- κ B

STX: Shiga toxin

STEC: STX-producing *Escherichia coli*

TNF- α : tumor necrosis factor- α

1. Introduction

Shiga toxin (STX), which is produced by some strains of *Escherichia coli*, causes hemorrhagic colitis and hemolytic-uremic syndrome (HUS) [1]. Twenty to fifty percent of HUS cases involve central nervous system (CNS) complications that are the most common cause of death in STEC infections [1, 2]. However, approximately 62% of patients with HUS showed symptoms of encephalopathy in the 2011 Japanese outbreak of infection with STX-producing *E. coli* (STEC) [3, 4]. Encephalopathy is a syndrome involving cerebral dysfunction with organic and inorganic causes, including cytokine storm and toxic reaction [5, 6].

We previously reported that the serum concentration of tumor necrosis factor (TNF)- α was higher in patients with acute encephalopathy than in patients without encephalopathy during the HUS phase in the 2011 Japanese outbreak of STEC infections [7]. This study suggested that TNF- α is a critical factor leading to encephalopathy in STEC infections.

The brain contains mainly neuronal and glial cells. Glial cells are the most abundant cells in the brain and play key roles in cerebral homeostasis, plasticity, and inflammation. Glial cells are functional components of the blood-brain barrier (BBB) [8]. Therefore, glial cell hypofunction and death impair cerebral homeostasis [9] and are thought to cause CNS complications, including encephalopathy [8].

In this study, we investigated the effect of TNF- α with or without STX on glial cells in vitro. Initially, we studied the effects of TNF- α alone on death of and inflammatory responses in glial cells. The subsequent analysis demonstrated that the combination of STX and TNF- α led to greater cell death and inflammatory responses than those noted with TNF- α alone in glial cells. Possibly, the role of TNF- α in sensitizing glial cells to STX contributes to the development of CNS complications in STEC infections.

2. Materials and methods

2.1. Materials

Dulbecco's modified Eagle's medium (DMEM), TNF- α , and Cell Counting Kit 8 were obtained from Wako Pure Chemical Industries, Ltd (Osaka, Japan). Shiga toxin-2 (STX) was obtained from Nacalai Tesque (Kyoto, Japan). Fetal bovine serum (FBS) was obtained from Invitrogen Corporation (Carlsbad, CA, USA). Anti-phospho-specific nuclear factor- κ B (NF- κ B) p65 (Ser536), anti-phospho-specific c-Jun N-terminal kinase (JNK; Thr183/Tyr185), anti-phospho-specific p38 mitogen-activated protein kinase (MAPK; Thr180/Tyr182), anti- β -actin, and horseradish-peroxidase-conjugated anti-rabbit IgG antibodies were purchased from Cell Signaling Technology (Danvers, MA, USA).

2.2. Cell culture

B92 rat cells [10] and primary rat glial cells [11] were gifted by Dr. Ohno-Shosaku (Kanazawa University, Japan). The glial fibrillary acidic protein, a marker of astrocyte, was expressed in primary cells that Dr. Ohno-shosaku provided, indicating astrocytes [9]. The cells were maintained in DMEM containing 10% FBS at 37°C in a 5% CO₂ incubator. Subcultures were required to maintain that both cells were in the logarithmic phase of growth during culture.

2.3. Cell proliferation assay

Cell proliferation was analyzed using Cell Counting Kit 8. B92 rat glial cells and primary glial cells were seeded in 96 well-plates at a density of 3×10^3 cells/well. After 24-h incubation, the cells were then treated with TNF- α and/or STX2 for 48 h. Next, the cells were incubated with 10 μ l WST-8 for 2 h. The absorbance of the colored formazan product produced by mitochondrial dehydrogenases in metabolically active cells was recorded at 450 nm as the background value. Cell proliferation was expressed as a percentage of absorbance obtained in treated wells relative to that in untreated (control) wells.

2.4. Western blotting

Western blotting was performed as described previously [12, 13].

2.5. Statistical analysis

Data are presented as the mean \pm SEM from at least three independent experiments. Statistical analysis was conducted using a Kruskal Wallis non-parametric ANOVA followed by a Bonferroni Inequality. Results were considered statistically significant when p -values were less than 0.05 or 0.01.

3. Results

3.1. TNF- α -induces downregulation of cell growth and upregulation of NF- κ B activity in B92 glial cells

First, we examined the effects of TNF- α on growth of cultured glial cells. TNF- α (50–500 ng/mL) considerably inhibited cell growth (Fig. 1A) and cleaved caspase-3 (Suppl. Fig. 1) in a concentration-dependent manner in B92 glial cells.

Next, we examined the effects of TNF- α on inflammation in cultured glial cells. TNF- α (5–500 ng/mL) significantly phosphorylated NF- κ B in a concentration-dependent manner in B92 glial cells (Fig.1B), indicating TNF- α induced activation of NF- κ B signaling.

3.2. STX facilitates TNF- α -induced downregulation of cell growth and TNF- α -induced upregulation of NF- κ B activity in B92 glial cells

Our previous study showed that STX (0.03–3 ng/mL) did not inhibit growth of B92 glial cells [14]. Likewise, in the current study, STX (3 ng/mL) alone failed to alter growth of B92 glial cells; however, it significantly facilitated TNF- α -induced downregulation of growth of these cells (Fig.2A).

Although STX alone failed to phosphorylate NF- κ B, it facilitated TNF- α -induced NF- κ B phosphorylation in B92 glial cells (Fig.2B).

3.3. STX facilitates TNF- α -induced activation of JNK and p38 MAPK in B92 glial cells

STX alone phosphorylated JNK but not p38 MAPK in B92 glial cells, indicating that it activates JNK but not p38 MAPK (Fig.3). In contrast, TNF- α alone phosphorylated both JNK and p38 MAPK in B92 glial cells, indicating that it activates both JNK and p38 MAPK (Fig.3). The combination of STX and TNF- α strongly phosphorylated both JNK and p38 MAPK in glial cells. STX significantly facilitated TNF- α -mediated phosphorylation of JNK (Fig.3).

3.4. STX facilitates TNF- α -induced upregulation of NF- κ B activity in primary glial cells

TNF- α , but not STX, phosphorylated NF- κ B in primary B92glial cells (Fig.4). Although STX alone failed to activate NF- κ B, it significantly facilitated TNF- α -induced NF- κ B phosphorylation, indicating the activation of NF- κ B (Fig.4).

3.5. STX facilitates TNF- α -induced activation of JNK and p38 MAPK in primary glial cells

TNF- α , but not STX, phosphorylated both JNK and p38 MAPK in primary glial cells (Fig.5). STX alone failed to activate JNK or p38 MAPK, similar to its effect on NF- κ B activity. However, STX significantly facilitated TNF- α -induced JNK and p38 MAPK phosphorylation in primary cells (Fig.5).

4. Discussion

In the current study, we demonstrated that TNF- α inhibits cell growth and activates NF- κ B, JNK, and p38 MAPK in a concentration-dependent manner in glial cells. STX combined with TNF- α decreased cell growth and activated NF- κ B, JNK, and p38 MAPK to a greater extent than STX alone or TNF- α alone. These results suggest that TNF- α modifies disease progression in STEC infections.

STEC and STX cause hemorrhagic diarrhea and the potentially lethal HUS in severe cases [1]. HUS in STEC infections often involves neurological symptoms and signs of encephalopathy [1, 2]. The neurological symptoms and signs of encephalopathy include seizures, alteration of consciousness, focal neurological deficits, visual disturbances, and brainstem symptoms. Neurological manifestations are associated with increased mortality.

STEC are gram-negative bacilli that have lipopolysaccharide (LPS) in their cell walls [15]. LPS acts as an endotoxin, stimulates macrophages, and causes release of cytokines from

macrophages, including TNF- α [16]. TNF- α is a master cytokine that mediates the inflammatory response and innate immunity [17]. TNF- α causes cell death and elicits stress responses, including JNK and p38 MAPK signaling [17]. In our previous study, we showed that the serum concentration of TNF- α in patients with acute encephalopathy was higher than that in patients without encephalopathy in the 2011 Japanese outbreak of STEC infections, suggesting that TNF- α is a critical cytokine in relation to the development of encephalopathy [7]. In this study, TNF- α induced death (Fig.1) of and NF- κ B, JNK, and p38 MAPK activation (Fig.3, 5) in glial cells, indicating that TNF- α might modify glial cell conditions.

However, the role of STX in the mechanism underlying encephalopathy is still unclear. Glial cells are the most abundant cells in the brain and are functional components of the BBB [8]. Thus, glial cells play key roles in cerebral homeostasis, plasticity, and inflammation. When the glial cells are dead or show abnormally low function, cerebral homeostasis might be disrupted. In our present in vitro study, the combination of STX and TNF- α caused much greater damage to glial cells than TNF- α alone or STX alone (Fig.2-5). STX is multifunctional protein inducing protein biosynthesis inhibition and endoplasmic reticulum (ER) stress response [18]. ER stress is widely reported to induce apoptotic cell death [19]. Thus, STX may enhance TNF- α induced cell death. These results indicate that STX, together with TNF- α , likely causes glial cell death and disrupts cerebral homeostasis, resulting in brain damage (including encephalopathy).

Indeed, STX and TNF- α have been found in the blood of STEC patients with encephalopathy [7]. When STX and TNF- α reach the brain, STX, TNF- α , or both attack the vascular endothelial cell layer that forms the anatomical BBB and disrupt the endothelial layer [20, 21]. In turn, TNF- α or the combination of STX and TNF- α affect glial cells that play roles in the functional BBB and homeostasis, and induce their death. Encephalopathy is recognized as a major determinant of mortality during the acute phase of STEC infections. Although the pathogenesis of CNS complications is unclear, BBB disruption and neuronal disturbances are observed during the acute phase of STEC infections [22-24]. The overall result of these events is likely to be encephalopathy in STEC patients.

5. Conclusions

The current study showed that TNF- α potentiated glial cell death and inflammation induced by STX treatment. Glial cells play key roles in cerebral homeostasis, plasticity, and inflammation; thus, these results suggest that TNF- α is a critical factor in relation to the development of encephalopathy in STEC infections.

Potential conflicts of interest

The authors do not have any conflicts of interest to declare.

Authors' contributions

HL conducted experiments. NS designed the study and prepared the manuscript. HL, MS, TT, and TW analyzed the data. NS and KO obtained funding. KO and AY edited the manuscript. All authors read and approved the final version of this manuscript.

Acknowledgments

We thank Dr. Takako Ohno-Shosaku for providing the B92 rat glial cells and primary rat glial cells.

This work was supported by Grants-in-Aid for Science and Culture (25460614 to KO, and 25282021, 26650173 and 16K13013 to NS) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

References

1. T. Magnus, J. Röther, O. Simova, M. Meier-Cillien, J. Repenthin, F. Möller, J. Gbadamosi, U. Panzer, M. Wengenroth, C. Hagel, S. Kluge, R.K. Stahl, K. Wegscheider, P. Urban, B. Eckert, M. Glatzel, J. Fiehler, C. Gerloff, The neurological syndrome in adults during the 2011 northern German *E. coli* serotype O104:H4 outbreak, *Brain* 135 (2012) 1850–1859. doi: 10.1093/brain/aws090.
2. F. Obata, Influence of *Escherichia coli* shiga toxin on the mammalian central nervous system, *Adv. Appl. Microbiol.* 71 (2010) 1–19. doi: 10.1016/S0065-2164(10)71001-7.
3. M. Watahiki, J. Isobe, K. Kimata, T. Shima, J. Kanatani, M. Shimizu, A. Nagata, K. Kawakami, M. Yamada, H. Izumiya, S. Iyoda, T. Morita-Ishihara, J. Mitobe, J. Terajima, M. Ohnishi, T. Sata, Characterization of enterohemorrhagic *Escherichia coli* O111 and O157 strains isolated from outbreak patients in Japan, *J. Clin. Microbiol.* 52 (2014) 2757-2763. doi: 10.1128/JCM.00420-14.
4. Y. Yahata, T. Misaki, Y. Ishida, M. Nagira, M. Watahiki, J. Isobe, J. Terajima, S. Iyoda, J. Mitobe, M. Ohnishi, T. Sata, K. Taniguchi, Y. Tada, N. Okabe, N. Epidemiological analysis of a large enterohaemorrhagic *Escherichia coli* O111 outbreak in Japan associated with haemolytic uraemic syndrome and acute encephalopathy, *Epidemiol. Infect.* 143 (2015) 2721-2732. doi: 10.1017/S0950268814003641.

5. M.R. Dobbs, Toxic encephalopathy, *Semin. Neurol.* 31(2011)184-193. doi: 10.1055/s-0031-1277989.
6. T.E. Gofton, G.B. Young, Sepsis-associated encephalopathy, *Nat. Rev. Neurol.* 8 (2012) 557-566. doi: 10.1038/nrneurol.2012.183.
7. M. Shimizu, M. Kuroda, N. Sakashita, M. Konishi, H. Kaneda, N. Igarashi, J. Yamahana, H. Taneichi, H. Kanegane, M. Ito, S. Saito, K. Ohta, T. Taniguchi, K. Furuichi, T. Wada, M. Nakagawa, H. Yokoyama, A. Yachie, A. Cytokine profiles of patients with enterohemorrhagic *Escherichia coli* O111-induced hemolytic-uremic syndrome, *Cytokine* 60 (2012) 694–700. doi: 10.1016/j.cyto.2012.07.038.
8. C. Iadecola, M. Nedergaard, Glial regulation of the cerebral microvasculature, *Nat. Neurosci.* 10 (2007) 1369–1376. doi:10.1038/mn2003.
9. N. Sugimoto, H. Leu, N. Inoue, M. Shimizu, T. Toma, M. Kuroda, T. Saito, T. Wada, A. Yachie, The critical role of lipopolysaccharide in the upregulation of aquaporin 4 in glial cells treated with Shiga toxin, *J. Biomed. Sci.* 22 (2015) 78. doi: 10.1186/s12929-015-0184-5.
10. D. Schubert, S. Heinemann, W. Carlisle, H. Tarikas, B. Kimes, J. Patrick, J.H. Steinbach, W. Culp, B.L. Brandt, Clonal cell lines from the rat central nervous system, *Nature* 249 (1974) 224-227.
11. T. Ohno-Shosaku, T. Maejima, M. Kano, Endogenous cannabinoids mediate retrograde signals from depolarized postsynaptic neurons to presynaptic terminals, *Neuron* 2001 (29) 729-38.

12. T. Saito, N. Sugimoto, K. Ohta, T. Shimizu, K. Ohtani, Y. Nakayama, T. Nakamura, Y. Hitomi, H. Nakamura, S. Koizumi, A. Yachie, Phosphodiesterase inhibitors suppress *Lactobacillus casei* cell-wall-induced NF- κ B and MAPK activations and cell proliferation through protein kinase A–or exchange protein activated by cAMPdependent signal pathway, *Sci. World J.* 2012 (2012) 748572. doi: 10.1100/2012/748572.
13. N. Sugimoto, K. Ohta, T. Saito, Y. Nakayama, T. Nakamura, A. Maeda, A. Yachie, *Lactobacillus casei* cell wall extract directly stimulates the expression of COX2 independent of Toll-like receptor 2 in rat glial cells, *J. Microbiol. Immunol. Infect.* 46 (2013) 389-392. doi: 10.1016/j.jmii.2012.03.004.
14. N. Sugimoto, T. Toma, M. Shimizu, M. Kuroda, T. Wada, A. Yachie, Shiga toxin-2 enhances heat-shock-induced apoptotic cell death in cultured and primary glial cells, *Cell Biol. Toxicol.* 30 (2014) 289-299. doi: 10.1007/s10565-014-9286-1.
15. H. Chart, C. Jenkins, The serodiagnosis of infections caused by Verocytotoxin-producing *Escherichia coli*, *J. Appl. Microbiol.* 86 (1999) 731-740.
16. C.A. Dinarello, Infection, fever, and exogenous and endogenous pyrogens: some concepts have changed, *J. Endotoxin Res.* 10 (2004) 201-222. doi: 10.1177/09680519040100040301.
17. C.L. Tsai, W.C. Chen, H.L. Hsieh, P.L. Chi, L.D. Hsiao, C.M. Yang, TNF- α induces matrix metalloproteinase-9-dependent soluble intercellular adhesion molecule-1 release via

- TRAF2-mediated MAPKs and NF- κ B activation in osteoblast-like MC3T3-E1 cells, *J. Biomed. Sci.* 21 (2014) 12. doi: 10.1186/1423-0127-21-12.
18. M.-S. Lee, S. Koo, D.G. Jeong, V.L. Tesh, Shiga toxins as multi-functional proteins: induction of host cellular stress responses, role in pathogenesis and therapeutic applications, *Toxins* 8 (2016) 77. Doi:10.3390/toxins8030077.
19. R.K. Gara, V.K. Srivastava, S. Duggal, J.K. Bagga, M. Bhatt, S. Sanyal, D.P. Mishra, Shikonin selectively induces apoptosis in human prostate cancer cells through the endoplasmic reticulum stress and mitochondrial apoptotic pathway, *J Biomed Sci.* 22 (2015) 26. doi: 10.1186/s12929-015-0127-1.
20. V.I. Landoni, P. Schierloh, M. de Campos Nebel, G.C. Fernández, C. Calatayud, M.J. Lapponi, M.A. Isturiz, Shiga toxin 1 induces on lipopolysaccharide-treated astrocytes the release of tumor necrosis factor-alpha that alter brain-like endothelium integrity, *PLoS Pathog.* 8 (2012) e1002632. doi: 10.1371/journal.ppat.1002632.
21. V.L. Tesh, Induction of apoptosis by Shiga toxins, *Future Microbiol.* 5 (2010) 431–453. doi: 10.2217/fmb.10.4.
22. M. Kuroda, M. Shimizu, N. Inoue, I. Ikeno, H. Nakagawa, A. Yokoi, Y. Niida, M. Konishi, H. Kaneda, N. Igarashi, J. Yamahana, H. Taneichi, H. Kanegane, M. Ito, S. Saito, K. Furuichi, T. Wada, M. Nakagawa, H. Yokoyama, A. Yachie, Serum tau protein as a marker of disease

activity in enterohemorrhagic *Escherichia coli* O111-induced hemolytic uremic syndrome,

Neurochem. Int. 85-86 (2015) 24-30. doi: 10.1016/j.neuint.2015.04.003.

23. T. Skripuletz, U. Wurster, H. Worthmann, M. Heeren, R. Schuppner, C. Trebst, J.T. Kielstein, K.

Weissenborn, M. Stangel, Blood-cerebrospinal fluid barrier dysfunction in patients with

neurological symptoms during the 2011 Northern German *E. coli* serotype O104:H4 outbreak,

Brain 136 (2013) e241. doi: 10.1093/brain/aws361.

24. J. Takanashi, H. Taneichi, T. Misaki, Y. Yahata, A. Okumura, Y. Ishida, T. Miyawaki, N. Okabe,

T. Sata, M. Mizuguchi, Clinical and radiologic features of encephalopathy during 2011 *E. coli*

O111 outbreak in Japan, *Neurology* 82 (2014) 564-572. doi:

10.1212/WNL.000000000000120.

Figure Legends

Fig.1. TNF- α induces decrease in cell proliferation and increase in phosphorylation of NF- κ B in B92 glial cells. (A) The effects of TNF- α on cell growth at 48 h after treatment. (B) The effects of TNF- α on phosphorylation of NF- κ B at 30 min after treatment. Data is representative of at least three independent experiments. Each column represents the mean \pm SEM. ** P < 0.01 versus vehicle control (0 ng/mL).

Fig.2. The combination of STX and TNF- α induces decrease in cell proliferation and increase in phosphorylation of NF- κ B in B92 glial cells. (A) The effects of STX and/or TNF- α on cell growth at 48 h after treatment. (B) The effects of STX and/or TNF- α on phosphorylation of NF- κ B at 30 min after treatment. Data is representative of at least three independent experiments. Each column represents the mean \pm SEM. * P < 0.05 or ** P < 0.01 versus vehicle control, [!] P < 0.01 versus STX alone, ^{##} P < 0.01 versus TNF- α alone.

Fig.3. The combination of STX and TNF- α increases phosphorylation of JNK and p38 in B92 glial cells. The effects of STX and/or TNF- α on phosphorylation of JNK at 30 min after treatment. The effects of STX and/or TNF- α on phosphorylation of p38 at 30 min after treatment. Data is representative of at least three independent experiments. Each column represents the mean \pm SEM.

* $P < 0.05$ or ** $P < 0.01$ versus vehicle control, !! $P < 0.01$ versus STX alone, ### $P < 0.01$ versus TNF- α alone.

Fig.4. The combination of STX and TNF- α increases phosphorylation of NF- κ B in primary glial cells. The effects of STX and/or TNF- α on phosphorylation of NF- κ B at 30 min after treatment. Data is representative of at least three independent experiments. Each column represents the mean \pm SEM.

** $P < 0.01$ versus vehicle control, !! $P < 0.01$ versus STX alone, ### $P < 0.01$ versus TNF- α alone.

Fig.5. The combination of STX and TNF- α increases phosphorylation of JNK and p38 in primary glial cells. The effects of STX and/or TNF- α on phosphorylation of JNK at 30 min after treatment. The effects of STX and/or TNF- α on phosphorylation of p38 at 30 min after treatment. Data is representative of at least three independent experiments. Each column represents the mean \pm SEM.

** $P < 0.01$ versus vehicle control, !! $P < 0.01$ versus STX alone, ### $P < 0.01$ versus TNF- α alone.






Fig. 1






Fig. 2





Fig. 3

Primary glial cells





Fig. 4

Primary glial cells





Fig. 5