

The first total synthesis of ( $\pm$ )-chanoclavine-? acid and an alternative total synthesis of ( $\pm$ )-chanoclavine-?

著者	Somei Masanori, Mukaiyama Harunobu, Nomura Toko, Nakagawa Kyoko
journal or publication title	Heterocycles
volume	31
number	11
page range	1919-1921
year	1990-01-01
URL	<a href="http://hdl.handle.net/2297/4318">http://hdl.handle.net/2297/4318</a>

THE FIRST TOTAL SYNTHESIS OF (±)-CHANOCLAVINE-I ACID AND AN ALTERNATIVE TOTAL SYNTHESIS OF (±)-CHANOCLAVINE-I<sup>1</sup>


Masanori Somei,\* Harunobu Mukaiyama, Yoko Nomura, and Kyoko Nakagawa  
Faculty of Pharmaceutical Sciences, Kanazawa University,  
13-1 Takara-machi, Kanazawa 920, Japan

**Abstract** ——— The total synthesis of (±)-chanoclavine-I acid was achieved for the first time. An alternative total synthesis of (±)-chanoclavine-I was also reported.

Chanoclavine-I acid (1) is a major alkaloid in the seeds of *Ipomea violacea* and its structure was established by H. R. Shough and co-workers in 1977.<sup>2</sup> In this communication, we wish to report the first total synthesis of (±)-chanoclavine-I acid and an alternative synthesis of (±)-chanoclavine-I (9).<sup>3e,i</sup>

In our continuing project for preparing biologically active ergot alkaloid derivatives,<sup>3</sup> (E)-4,5-trans-5-(2-hydroxymethyl-1-propen-1-yl)-4-nitro-1,3,4,5-tetrahydrobenz[cd]indole (3) is now readily available in five steps in 15% overall yield starting from indole-3-carboxaldehyde (2).<sup>3e</sup> Therefore, 3 was converted to the aldehyde (4) (mp 186-187°C) in 55% yield by the treatment with pyridinium chlorochromate in methylene chloride. Subsequent oxidation of 4 with sodium hypochlorite in the presence of 2-methyl-2-butene<sup>4</sup> afforded the corresponding carboxylic acid (5) (mp 264-265°C) in 97% yield. Methylation of 5 with ethereal diazomethane generated 95% yield of the methyl ester (6) (mp 188-189°C), which was then reduced with zinc amalgam and hydrochloric acid to produce the corresponding amine (7) (oil) in 98% yield. Since we have already succeeded in the synthesis of (±)-chanoclavine-I (9),<sup>3e,i</sup> we attempted to convert 7 to 9 in order not only to confirm its structure but also to establish an alternative synthesis route to 9. Thus, treatment of 7 with methyl chloroformate in the presence of triethylamine produced 67% yield of the corresponding methyl carbamate (8) (oil). Reduction of 8 with lithium aluminum hydride in dry tetrahydrofuran afforded 9 in 95% yield. Various attempts<sup>5</sup> to obtain chanoclavine-I acid methyl ester (11) by methylating

the amine (7) were made in vain because of the steric hindrance around the amino group. Finally, methylation of 7 with dimethyl sulfate in the presence of potassium carbonate was found to produce the desired 11 (oil) in 25% yield together with 15% yield of the corresponding dimethylamino derivative (10) (mp 141-142°C). Alkaline hydrolysis of 11 in methanol and subsequent column chromatography on Amberlite IRA 120 afforded (±)-chanoclavine-I acid<sup>6</sup> (1) in 80% yield. Although 1 did not exhibit clear carbonyl absorption band in its ir spectrum, the HCl salt<sup>7</sup> of 1 showed strong absorption band at 1698cm<sup>-1</sup>.<sup>2</sup> For further structural confirmation, methylation of 1 was carried out with ethereal diazomethane to produce 10 and 11 in 7% and 35% yields, respectively. Mass spectral data of 1 and <sup>1</sup>H-nmr spectrum of its HCl salt were identical to those reported in the literature.<sup>2</sup>



## ACKNOWLEDGEMENT

This work is partly supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, and Culture of Japan, which is gratefully acknowledged.

## REFERENCES AND NOTES

1. This paper is part 53 of a series entitled "The Chemistry of Indoles". Part 52: T. Kawasaki and M. Somei, Heterocycles, 1990, 31, 1605.
2. T. C. Choong and H. R. Shough, Tetrahedron Lett., 1977, 3137.
3. a) M. Somei, F. Yamada, Y. Karasawa, and C. Kaneko, Chemistry Lett., 1981, 615; b) M. Somei and M. Tsuchiya, Chem. Pharm. Bull., 1981, 29, 3145; c) M. Somei and F. Yamada, ibid., 1984, 32, 5064; d) F. Yamada, Y. Makita, T. Suzuki, and M. Somei, ibid., 1985, 33, 2162; e) M. Somei, Y. Makita, and F. Yamada, ibid., 1986, 34, 948; f) M. Somei, F. Yamada, and Y. Makita, Heterocycles, 1987, 26, 895; g) M. Somei, F. Yamada, H. Ohnishi, Y. Makita, and M. Kuriki, ibid., 1987, 26, 2823; h) H. Watanabe, M. Somei, S. Sekihara, K. Nakagawa, and F. Yamada, Japan J. Pharmacol., 1987, 45, 501; i) M. Somei, Yakugaku Zasshi, 1988, 108, 361.
4. B. S. Bal, W. E. Childers, Jr., and H. W. Pinnick, Tetrahedron, 1981, 37, 2091.
5. Both Eschweiler-Clark and Leuckart reactions did not work. Attempts to produce imine by the reaction of 7 with various aldehydes were found to be unsuccessful.
6. Colorless powder from ethanol-H<sub>2</sub>O. mp 289-290°C (dec.). Ir (KBr) cm<sup>-1</sup>: 3150, 1607, 1564, 1380, 1084, 760. 400MHz <sup>1</sup>H-nmr (DMSO-d<sub>6</sub>) δ: 1.94 (3H, s), 2.37 (3H, s), 2.65 (1H, dd, J=15 and 8 Hz), 2.86 (1H, dt, J=4 and 8 Hz), 3.16 (1H, dd, J=15 and 4 Hz), 3.92 (1H, dd, J=10 and 8 Hz), 6.49 (1H, d, J=7.5 Hz), 6.55 (1H, d, J=10 Hz), 6.98 (1H, t, J=7.5 Hz), 7.01 (1H, s), 7.15 (1H, d, J=7.5 Hz), 10.66 (1H, s, COOH). Ms m/z (%): 270 (M<sup>+</sup>, 100), 252 (8.8), 197 (14.8), 168 (41.3), 167 (8.4), 155 (35.0), 154 (25.7). High resolution ms m/z: Calcd for C<sub>16</sub>H<sub>18</sub>N<sub>2</sub>O<sub>2</sub>: 270.1367. Found: 270.1363.
7. Colorless prisms from ethanol-H<sub>2</sub>O. mp 288-289°C (dec.). Ir (KBr) cm<sup>-1</sup>: 3330, 2920, 1698, 1581, 1445, 1221, 753. 400MHz <sup>1</sup>H-nmr (DMSO-d<sub>6</sub>+D<sub>2</sub>O) δ: 2.03 (3H, s), 2.65 (3H, s), 3.12 (1H, dd, J=15 and 7 Hz), 3.32 (1H, dd, J=15 and 4 Hz), 3.68 (1H, dt, J=4 and 7 Hz), 4.35 (1H, dd, J=10 and 7 Hz), 6.53 (1H, d, J=10 Hz), 6.68 (1H, d, J=8 Hz), 7.15 (1H, t, J=8 Hz), 7.20 (1H, s), 7.31 (1H, d, J=8 Hz), 10.91 (1H, s, COOH).

Received, 27th September, 1990