

Correlation study of observation data of polycyclic aromatic hydrocarbons and chemical transport model calculation data of other pollutants at Wajima

著者	Yang Xiaoyang, Tang Ning, Okada Yumi, Kameda Takayuki, Toriba Akira, Utanohara Yoichi, Yamaji Kazuyo, Kawanishi Takuya, Tamura Kenji, Uno Itsushi, Hayakawa Kazuichi
journal or publication title	金沢大学 21世紀COEプログラム主催国際シンポジウム「東アジアの大気環境汚と健康・生態系への影響」講演要旨集(金沢)
page range	28-28
year	2006-09-25
URL	http://hdl.handle.net/2297/2450

Correlation study of observation data of polycyclic aromatic hydrocarbons and chemical transport model calculation data of other pollutants at Wajima

Xiaoyang Yang^a, Ning Tang^a, Okada Yumi^a, Takayuki Kameda^a, Akira Toriba^a, Yoichi Utanohara^a, Kazuyo Yamaji^b, Takuya Kawanishi^a, Kenji Tamura^c, Toshimasa Ohara^c, Itsushi Uno^d, Kazuichi Hayakawa^a

^aGraduate School of Natural Science and Technology, Kanazawa University, Japan

^bFrontier Research Center for Global Change, Japan Agency for Marine-Earth Science and Technology Japan

^cNational Institute for Environmental Studies, Japan

^dResearch Institute for Applied Mechanic, Kyushu University, Japan

Abstract

Among the pollutants in the atmosphere, several polycyclic aromatic hydrocarbons (PAHs) and nitropolycyclic aromatic hydrocarbons (NPAHs) have carcinogenicity/mutagenicity or endocrine disrupting activity. PAHs and NPAHs are exhausted from the combustion of petroleum and coal. Coal consumption in China is the largest in the world and represents over 75% of the energy source in China. Long-range transport of atmospheric PAHs and NPAHs from China to Japan was investigated by collecting airborne particulates at Wajima in Noto peninsula from September 17, 2004 to September 16, 2005. As a result, the atmospheric concentration of PAHs showed a high level during the heating period of China. Furthermore, the composition of PAHs at Wajima, during the period, was close to that at Shenyang but not Kanazawa.

On the other hand, it is thought that the simulation model is also a helpful method to predict the long-range transport of atmospheric pollutants such as PAHs and NPAHs. A Community Multi-scale Air Quality model based on the Regional Atmospheric Modeling System for the Asian Domain had been developed and had been used to simulate the emission and transportation of several pollutants such as SO_x and NO_x. Considering the reaction mechanism of PAHs and NPAHs in the atmosphere is not clear enough. We studied the correlations of observation data of PAHs and chemical transport model calculation data of other pollutants at Wajima as the first step of making a new model. As a result, good correlations were found between PAHs and SO₂, PAHs and ANO₃.