

## Bioassay Using Daphnia Daphnia Trajectories

著者	Kimura Koji, Hayashi Yoshishige, Kawanishi Takuya, Ogino Chiaki, Shimizu Nobuaki, Sato Kazuto
journal or publication title	Proceedings, International Symposium of the Kanazawa University 22st-Century COE Program
volume	1
page range	369-370
year	2003-03-16
URL	<a href="http://hdl.handle.net/2297/6431">http://hdl.handle.net/2297/6431</a>

## Bioassay Using *Daphnia* Trajectories

Koji KIMURA, Yoshishige HAYASHI, Takuya KAWANISHI, Chiaki OIGINO, Nobuaki SHIMIZU  
Graduate School of Natural Science and Technology, Kanazawa University, Kanazawa, Ishikawa 920-1192, JAPAN  
Kazuto SATO

Department of Chemistry and Chemical Engineering, Kanazawa University, Kanazawa, Ishikawa 920-8667, JAPAN

**Abstract** - The effect of copper salt concentration on the trajectories of *Daphnia magna* was investigated, and the significant difference was observed between 0.02 mg l<sup>-1</sup> copper solution and the control.

### I. Introduction

*Daphnia* species has been employed extensively in the environmental monitoring of toxic chemicals [1, 2]. Shimizu et al. have shown the possibility of the application of *Daphnia* trajectories to bioassay [3]. In this paper, we investigated the applicability of *Daphnia* trajectory analysis to the detection of toxicity of copper ions.

### II. Materials and Methods.

#### A. Culture Methods

A single clone of *Daphnia magna* was used in this study. Culture conditions were adapted from protocols for ASTM standards (American Society of Testing and Materials, 1986). The culture medium was synthetic freshwater (48 mg of NaHCO<sub>3</sub>, 30mg of CaSO<sub>4</sub>·2H<sub>2</sub>O, 30 mg of MgSO<sub>4</sub>, and 2 mg of KCl per liter of deionized water adjusted to a pH of 7.4) with a hardness of 45 mg of CaCO<sub>3</sub> per liter. Culture vessels were 3-L clear glass jars that were filled with approximately 2 L of culture medium. The density of *Daphnia* in the vessel was kept below 30 animals per liter. *Daphnia* were fed daily with a mixture of commercial fish food and dry yeast, prepared according to an ASTM protocol. *Daphnia* were kept under a light:dark period of 16 h:8 h at 20 ± 1 °C. All salt reagents were of analytical grade.

#### B. Analysis of Swimming Trajectory

Experimental setup is shown in Fig. 1. *Daphnia magna* less than 24 h old was chosen from the stock culture and transferred to a temperature controlled (20 ± 1 °C, with EYELA: CTP101) observation chamber (30 mm in diameter), filled with 4 ml of the ASTM solution containing various amount of copper salt (CuSO<sub>4</sub>·H<sub>2</sub>O; Wako Pure Chemical Industries, Osaka, Japan). *Daphnia* swimming behavior was analyzed with a digital motion analysis system, consisting of digital microscope (KEYENCE : VH-6300), a video capturing board, and a personal computer with motion

analysis software.

At one consecutive measurement, the position of *Daphnia magna* ( $x, y$  coordinate) was recorded for 60 seconds at the interval of 0.1 seconds (600 frames). This procedure was repeated at the interval of 5 to 15 minutes.




Fig. 1 Experimental apparatus.

#### C. Calculation of Fractal Based Parameters

We used a modified dividers method for the calculation of the fractal dimension of the *Daphnia* trajectories.

In the original dividers method, the length of the system  $L$  was evaluated by the number of the steps  $n_\epsilon$  required to cover the system by the divider of length  $\epsilon$ , then the length  $L$  as a function of  $\epsilon$  is expressed as follows.

$$L(\epsilon) = n_\epsilon \epsilon \quad (1)$$

If the system is a fractal, there is a relationship between  $L(\epsilon)$  and the  $\epsilon$  as follows.

$$L(\epsilon) = A \epsilon^{1-D} \quad (2)$$

where  $D$  is the fractal dimension and  $A$  is a constant.

In our experiments, however, we were not able to use dividers of constant lengths. Instead we used the intervals of segment number,  $n$ , for the measure of the length. The length of the *Daphnia* trajectories are evaluated via

$$L = \sum_{j=1}^{N/n} \left\{ (x_{nj} - x_{n(j-1)})^2 + (y_{nj} - y_{n(j-1)})^2 \right\}^{0.5} \quad (3)$$

where  $N$  is the total number of the segments, and was 600 in this research,  $x_i$  and  $y_i$  are the coordinate of the *Daphnia* at  $i$ th segment.

Then, there is a relationship between  $n$  and  $L$  such that

$$L = Bn^{1-D'} \quad (4)$$

where  $B$  is a constant and  $D'$  is a parameter which can be related to  $D$  as follows.

$$D = \frac{1}{2 - D'} \quad (5)$$

### III. Results and Discussion

Fig. 1 shows the relationship between the intervals of segment number  $n$  and the length of the system evaluated by Eq. (3). For  $n$  smaller than 10, the relationship in double-log plot showed linearity, and we were able to estimate the parameter  $D'$  from the slopes in double-log plots.

Figs. 2 shows the change in parameter  $D'$  at various concentration of copper salt. It is clearly seen that the parameter  $D'$  began to increase tens of minutes after *Daphnia magna* was put into the solution containing the copper salt. Also, it is seen that it took more time before the parameter  $D'$  showed significant difference from the control for solutions having less amount of copper salt.




Fig. 1. Determination of parameter  $D'$ .




Fig. 2. Change in  $D'$  at various copper concentration.




Fig. 3. Change in  $D'$  at 0.02 ppm of copper.

Reported  $EC_{50}$  concentrations of copper detected by *Daphnia magna* were at around 0.02 ppm. Fig. 3 shows that the difference of  $D'$  between the copper solution of this concentration and control became significant about 5 hours after *Daphnia magna* was put into the toxic solution. (Note that the traditional  $EC_{50}$  test will take 48 hours before getting results.) This suggests that by using *Daphnia* trajectories, we may be able to establish a more efficient bioassay than  $EC_{50}$  analysis.

### References

- [1] Environmental Protection Agency, *Methods for Measuring the Acute Toxicity of Effluents and Receiving Waters to Freshwater and Marine Organisms*, 4th ed., EPA, pp. 44-69, 148-168, 1991.
- [2] International Organization for Standardization, *Water Quality - Determination of the Inhibition of the Mobility of *Daphnia magna* Straus (Cladocera, Crustacea)- Acute Toxicity Test*, ISO, 1989.
- [3] N. Shimizu, C. Ogino, T. Kawanishi, Y. Hayashi, "Fractal analysis of *Daphnia* motion for acute toxicity bioassay," *Environmental Toxicology* Vol. 17, pp. 441-448, 2002.