

Nucleophilic substitution reactions on indole nucleus: Formation of (3a,8a-cis)-1,2,3,3a,8,8a-hexahydropyrrolo-[2,3-b]indoles having a substituent at the 3a-position

著者	Yamada Fumio, Goto Aya, Hasegawa Masakazu, Kobayashi Kensuke, Somei Masanori
著者別表示	山田 文夫, 染井 正徳
journal or publication title	Heterocycles
volume	95
number	2
page range	844-861
year	2017-12-13
URL	<a href="http://doi.org/10.24517/00049644">http://doi.org/10.24517/00049644</a>

doi: 10.3987/COM-16-S(S)53



## NUCLEOPHILIC SUBSTITUTION REACTIONS ON INDOLE NUCLEUS: FORMATION OF (3a,8a-*cis*)-1,2,3,3a,8,8a-HEXAHYDROPYRROLO-[2,3-*b*]INDOLES HAVING A SUBSTITUENT AT THE 3a-POSITION<sup>1,#</sup>

Fumio Yamada, Aya Goto, Masakazu Hasegawa, Kensuke Kobayashi, and Masanori Somei\*<sup>2</sup>


Faculty of Pharmaceutical Sciences, Graduate School of Natural Science and Technology, Kanazawa University, Kakuma-machi, Kanazawa, 920-1192, Japan  
Corresponding author: e-mail address: somei.home@topaz.plala.or.jp

**Abstract** – Various nucleophiles, such as indole, 1,2,3-trimethoxybenzene, anisole, phenol, and pyrrole, reacted with 1-hydroxy-*Nb*-trifluoroacetyltryptamine under the presence of mesyl chloride to give novel series of (3a,8a-*cis*)-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-*b*]indoles having a substituent at the 3a-position. Their structures and by-products were strictly determined.

### INTRODUCTION

We have opened the door to the chemistry of 1-hydroxyindole and 1-hydroxytryptophan derivatives,<sup>3</sup> and demonstrated that these compounds generally undergo nucleophilic substitution reaction,<sup>4</sup> which was thus far rarely observed in indole chemistry.<sup>4</sup>

In our 1-hydroxyindole hypothesis,<sup>5</sup> we assume the 1-hydroxy group of the general formula (A) in Scheme 1 departs, after being transformed to a good leaving group (B), leaving a resonance stabilized indolyl cation<sup>6</sup> (C). It would be possible to trap it with suitable nucleophiles to give imine<sup>6</sup> (D).




Subsequent cyclization of *Nb*-nitrogen on the side chain results in providing simple and novel methodology for the preparation of (3*a*,8*a*-*cis*)-1,2,3,3*a*,8,8*a*-hexahydropyrrolo[2,3-*b*]indoles (**E**) having an employed nucleophile at the 3*a*-position. According to the idea, we first employed indole as a nucleophile and reported the result as the previous communication.<sup>7</sup> This is its full report together with the results of additionally examined nucleophiles such as 1,2,3-trimethoxybenzene, anisole, phenol, and pyrrole.

## RESULTS AND DISCUSSION

### I. Reaction of 1-hydroxy-*Nb*-trifluoroacetyltryptamine (**2**) with indole

*Nb*-Trifluoroacetyltryptamine (**1**, Scheme 2) was converted to 1-hydroxy-*Nb*-trifluoroacetyltryptamine (**2**) by our 1-hydroxyindole synthetic method.<sup>3</sup> Then, **2** was reacted with mesyl chloride (MsCl) in 1,2-dichloroethane in the presence of indole (3 mol eq) and trimethylamine (Et<sub>3</sub>N) at 0 °C (Table 1, Entry 3). As expected, smooth reaction occurred to provide 1,2,3,8-tetrahydro-1-trifluoroacetylpyrrolo[2,3-*b*]indole<sup>8,9</sup> (**3**), (3*a*,8*a*-*cis*)-1,2,3,3*a*,8,8*a*-hexahydro-3*a*-(indol-2-yl)- (**4**), -3*a*-(indol-3-yl)-1-trifluoroacetylpyrrolo[2,3-*b*]indole (**5**), and 6-mesyloxy-*Nb*-trifluoroacetyltryptamine<sup>8,9</sup> (**6**), in 13, 5, 11, and 3%



**Scheme 2**

yields, respectively.


With an attempt to improve the product yield of nucleophilic reaction and examine solvent effect, 1,2-dichloroethane was changed to benzene, CHCl<sub>3</sub>, THF, DMF, MeCN, MeNHCHO, and EtOAc. The product and their distribution ratio variably changed and their results are summarized in Table 1.

When the reaction was carried out in CHCl<sub>3</sub> (Entry 2), the yield of **5** was improved to 21% together with the formations of **3**, **4**, and **6** in the respective yields of 14, 5, and 4%. Under similar reaction conditions, the use of excess indole (10 mol eq., Entry 8) further raised the yield of **5** up to 30% in addition to the concomitant formations of **3**, **4**, and **6** in 4, 7, and 1% yields, respectively.

In the case of THF as the solvent, various products were formed (Entry 4). Thus, the reaction of **2** with MsCl in THF in the presence of indole (3 mol eq) and Et<sub>3</sub>N at 0 °C gave **3**, **4**, **5**, **6**, 3*H*-3-(indol-3-yl)-*Nb*-trifluoroacetyltryptamine (**7**), and (3*a*,8*a*-*cis*)-3*a*-(4-chlorobutoxy)-1,2,3,3*a*,8,8*a*-hexahydro-1-trifluoroacetylpyrrolo[2,3-*b*]indole<sup>9</sup> (**8**), in 28, 6, 15, 5, 4, and 6% yields, respectively. From the results shown in Table 1, we found that solvent polarity has no effect for the preferred product formation, though

MeNHCHO produced (3a,8a-*cis*)-1,2,3,3a,8,8a-hexahydro-3a-hydroxy-1-trifluoroacetylpyrrolo[2,3-*b*]indole (**9**) as a major product (Entry 7).

**Table 1. Solvent effect on the product formation and distribution**



Entry	Solvent ( $\epsilon$ )	Yield (%) of						
		<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>
1	benzene (2)	18	0	4	0	4	0	0
2	$\text{CHCl}_3$ (4.8)	14	5	21	4	0	0	0
3	$\text{ClCH}_2\text{CH}_2\text{Cl}$ (25)	13	5	11	3	0	0	0
4	THF (30)	28	6	15	5	4	6	0
5	DMF (37)	30	1	7	2	0	0	0
6	MeCN (38)*	10	1	8	0	0	0	0
7	MeNHCHO (182)	2	1	4	0	0	0	20

\* **1** was obtained in 6% yield.


<b>2</b> $\xrightarrow{\text{Indole (10 mol eq.), MsCl, Et}_3\text{N}}$		Entries 8—10						
8	$\text{CHCl}_3$ (4.8)	4	7	30	1	0	0	0
9	$\text{ClCH}_2\text{CH}_2\text{Cl}$ (25)	8	5	18	2	0	0	0
10	EtOAc (30)	6	7	25	3	0	0	0

## II. Reaction of 1-hydroxy-Nb-trifluoroacetyltryptamine (**2**) with nucleophiles

We next examined aromatic electron rich nucleophiles. When 1,2,3-trimethoxybenzene (10 mol eq.) was employed in the reaction of **2** with MsCl in  $\text{CHCl}_3$  in the presence of  $\text{Et}_3\text{N}$  (Scheme 3), (3a,8a-*cis*)-1,2,3,3a,8,8a-hexahydro-1-trifluoroacetyl-3a-(2,3,4-trimethoxyphenyl)pyrrolo[2,3-*b*]indole (**10**), **3**, **6**, **1**, and (3a,8a-*cis*)-1,2,3,3a,8,8a-hexahydro-1-trifluoroacetyl-3a-[3-(Nb-trifluoroacetyl)aminoethylindol-1-yl]pyrrolo[2,3-*b*]indole (**11a**) were formed in 2, 15, 4, 4, and 13% yields, respectively. Further treatment of **11a** with  $\text{NaHCO}_3$  afforded **11b** in 67% yield.

Under similar reaction conditions with anisole as a nucleophile, (3a,8a-*cis*)-1,2,3,3a,8,8a-hexahydro-3a-(4-methoxyphenyl)-1-trifluoroacetylpyrrolo[2,3-*b*]indole (**12**), **3**, **1**, and **11a** were isolated in the

respective yields of 5, 18, 6, and 10%. **12** was easily converted to (3a,8a-*cis*)-1,2,3,3a,8,8a-hexahydro-3a-(4-methoxyphenyl) pyrrolo[2,3-*b*]indole (**13**) in 94% yield by the treatment with aq. NaHCO<sub>3</sub>. In the case of employing phenol as a nucleophile, (3a,8a-*cis*)-1,2,3,3a,8,8a-hexahydro-3a-(4-hydroxyphenyl)- (**14**) and -3a-(2-hydroxyphenyl)-1-trifluoroacetylpyrrolo[2,3-*b*]indole (**15**) were produced in addition to **6**, **1**, and **11a** in 3, 5, 4, 8, and 8% yields, respectively. The compound **14** was derived to **12** in 73% yield by the reaction with CH<sub>2</sub>N<sub>2</sub>.



**Scheme 3**


Since pyrrole is a good nucleophile, expected product, (3a,8a-*cis*)-1,2,3,3a,8,8a-hexahydro-3a-(pyrrol-2-yl)-1-trifluoroacetylpyrrolo[2,3-*b*]indole (**16**), was obtained in rather better yield (29%) compared to the above products (**10–12**, **14**, **15**) together with **3**, **1**, and **11a** in 6, 7, and 8% yields, respectively. Formation of the other expected isomer, pyrrol-3-yl isomer (**17**), was not detected at all. Treatment of **16** with Ac<sub>2</sub>O-pyridine afforded (3a,8a-*cis*)-8-acetyl-1,2,3,3a,8,8a-hexahydro-3a-(pyrrol-2-yl)-1-trifluoroacetylpyrrolo[2,3-*b*]indole (**18**) in 83% yield.

### III. Structural determination of products (Scheme 4)

Structures of various products reported in the above sections were determined spectroscopically. In cases where spectroscopically more than two structures were possible candidates, the product was led to suitable derivative which could prove its structure.

The high resolution MS and other spectral data of **4** and **5** show the presence of an extra indole moiety in both molecules. In the <sup>1</sup>H-NMR spectra, **4** and **5** show characteristic C-(8a) proton signal at δ5.63 and 5.92, respectively, proving the presence of hexahydropyrrolo[2,3-*b*]indole skeleton. In addition, **5** has a long-range coupled doublet proton ( $J=2.5$  Hz) at δ6.93 and is assigned to be C(2')-proton, which is unusually shielded compared to the usual indole C(2)-proton.<sup>10,11</sup> In the spectrum of **4**, a double doublets proton ( $J=2.2$  and 0.7 Hz) resonates at δ6.48, which is attributed to the C(3')-proton. The structures of **4** and **5** were further confirmed by treating them with Ac<sub>2</sub>O and pyridine to provide the acetyl derivatives (**20** and **19**) in the respective yields of 97 and 51% (Scheme 4). From these data, **4** and **5** were deduced to be indol-2-yl and indol-3-yl isomers, respectively.

Repeated recrystallization of **4** formed suitable prisms for X-ray single crystallographic analysis and the structure was determined unequivocally as shown in Figure 1. Since the indol-2-yl structure of **4** is established, then it determines that the other isomer (**5**) is the indol-3-yl isomer. The preferred formation of **5** to **4** is in accord with the well-known positional order 3>2 for the reactivity of unsubstituted indole.



The structure of **6** was proved as reported in the previous paper<sup>9</sup> by converting it to 1-acetyl-6-mesyloxy-*Nb*-trifluoroacetyltryptamine (**21**) in 25% yield by the treatment with NaH-AcCl. The compound (**7**) has a ring opened structure of **5**. It was proved by isolating **7** in 89% yield when **5** was heated in DMSO at 130 °C.

To establish the structure of **9**, it was converted to the common compound for structural determination by series of reactions. First, **9** was led to (3a,8a-*cis*)-3a-acetoxy- (**22**) and -8-acetyl-1,2,3,3a,8,8a-hexahydro-1-trifluoroacetylpyrrolo[2,3-*b*]indole (**23**) in 71 and 17% yields, respectively, by the reaction with Ac<sub>2</sub>O-pyridine at rt. Treatment of **22** with Ac<sub>2</sub>O-pyridine at 55 °C afforded **23** in 62% yield together with 26% yield of recovery. Hydrolysis of trifluoroacetyl group of **23** with aq. NaHCO<sub>3</sub> at rt provided (3a,8a-*cis*)-3a-acetoxy-8-acetyl-1,2,3,3a,8,8a-hexahydro- (**24**) and (3a,8a-*cis*)-8-acetyl-1,2,3,3a,8,8a-hexahydro-3a-hydroxypyrrolo[2,3-*b*]indole (**25**) in 80 and 19% yields, respectively. At the reflux conditions **23** gave 98% yield of **25**. Treatment of both **24** and **25** with Ac<sub>2</sub>O-pyridine at rt furnished (3a,8a-*cis*)-3a-acetoxy-1,8-diacetyl-1,2,3,3a,8,8a-hexahydroxypyrrolo[2,3-*b*]indole (**26**) in the respective yields of 99 and 96%.

On the other hand, **26** was obtained from **27**<sup>3,4</sup> by the treatment with Ac<sub>2</sub>O-NaOAc. Aside from this, **26** was produced by the treatment with Ac<sub>2</sub>O-pyridine in 80% yield from (3a,8a-*cis*)-1-acetyl-1,2,3,3a,8,8a-hexahydro-3a-hydroxypyrrolo[2,3-*b*]indole (**29**), which was derived in 80% yield from 8b-(2-acetyl-aminoethyl)-2,2-dimethyl-4*H*-1,3-dioxolo[4,5-*b*]indole (**28**) by the treatment with K<sub>2</sub>CO<sub>3</sub>. Since the structure of a derivative of **28** is determined by X-ray single crystallographic analysis as reported in our previous paper,<sup>12</sup> the structure of common compound (**26**) is established.

In conclusion, 1-hydroxy-*Nb*-trifluoroacetyltryptamine is a suitable starting material for obtaining novel type of (3a,8a-*cis*)-1,2,3,3a,8,8a-hexahydroxypyrrolo[2,3-*b*]indoles carrying aromatic and/or heteroaromatic substituent at the 3a position. Among this family members are core structures of Leptosins A–F,<sup>13</sup> which are cytotoxic substances against P-388 lymphocytic leukemia cell line comparable to that of mytomycin C. Therefore, we expect that compounds shown in this paper would have a useful biological activity.

## Experimental

Melting points were determined on a Yanagimoto micro melting point apparatus and are uncorrected. Infrared (IR) spectra with a Shimadzu IR-420, a Shimadzu IR-460, and a Horiba FT-720 spectrophotometer and proton nuclear magnetic resonance (<sup>1</sup>H-NMR) spectra with a JEOL JNM-GSX 500 spectrometer with tetramethylsilane as an internal standard. Mass spectra (MS) were recorded on a JEOL SX-102A spectrometer. Column chromatography was performed on silica gel (SiO<sub>2</sub>, 100-200 mesh, from Kanto Chemical Co. Inc.) throughout the present study. The solution of diazomethane (CH<sub>2</sub>N<sub>2</sub>) in diethylether (Et<sub>2</sub>O) was prepared as follows: a solution of potassium hydroxide (KOH) (5.50 g, 98.0

mmol) in H<sub>2</sub>O (8.0 mL) was placed in a 500 mL round bottom flask and cooled in an ice bath. The 95% EtOH (25 mL), Et<sub>2</sub>O (60.0 mL), and *p*-tolylsulfonylmethylnitrosoamide (21.5 g, 100 mmol) were added and the whole was slowly distilled to give the Et<sub>2</sub>O solution including about 3 g of CH<sub>2</sub>N<sub>2</sub>. Anhydrous *N,N*-dimethylformamide (DMF), tetrahydrofuran (THF), and CHCl<sub>3</sub> were prepared by distillation over calcium hydride, sodium, and calcium chloride, respectively.

**Reaction of 1-hydroxy-Nb-trifluoroacetyltryptamine (2) with indole as a nucleophile: 1,2,3,8-Tetrahydro-1-trifluoroacetylpyrrolo[2,3-*b*]indole (3), (3a,8a-*cis*)-1,2,3,3a,8,8a-hexahydro-3a-(indol-2-yl)-1-trifluoroacetyl- (4), -3a-(indol-3-yl)-1-trifluoroacetyl- (5), (3a,8a-*cis*)-3a-(4-chlorobutoxy)-1,2,3,3a,8,8a-hexahydro-1-trifluoroacetylpyrrolo[2,3-*b*]indole (8), 6-mesyloxy-Nb-trifluoroacetyltryptamin (6), 3*H*-3-(indol-3-yl)-Nb-trifluoroacetyltryptamine (7) from 2** — [Table 1, Entry 4]: A solution of MsCl (232.9 mg, 1.99 mmol) in anhydrous THF (2.0 mL) was added to a solution of **2** (419.9 mg, 1.54 mmol) and indole (542.6 mg, 4.73 mmol) in anhydrous THF (14.0 mL) and Et<sub>3</sub>N (1.6 mL, 11.5 mmol) at 0 °C and the mixture was stirred at 0 °C for 1 h. After addition of H<sub>2</sub>O under ice cooling, the whole was extracted with CHCl<sub>3</sub>–MeOH (95:5, v/v). The extract was washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and evaporated under reduced pressure to leave an oil, which was column-chromatographed on SiO<sub>2</sub> successively with CHCl<sub>3</sub>–hexane (1:1, v/v), CHCl<sub>3</sub>, CHCl<sub>3</sub>–MeOH (95:5, v/v), EtOAc–hexane (1:5, v/v), and EtOAc–hexane (1:2, v/v) to give **3** (107.7 mg, 28%), **8** (31.9 mg, 6%), **4** (31.4 mg, 6%), **5** (86.3 mg, 15%), **7** (23.4 mg, 4%), and **6** (29.4 mg, 5%) in the order of elution. **3**: mp 238.0–240.0 °C (decomp., colorless plates, recrystallized from CH<sub>2</sub>Cl<sub>2</sub>–hexane). IR (KBr): 3370, 1670, 1446, 1351, 1278, 1233, 1203, 1139, 1069, 746 cm<sup>-1</sup>. <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 3.30 (2H, t, *J*=7.4 Hz), 4.71 (2H, t, *J*=7.4 Hz), 7.15 (1H, dt, *J*=1.6, 6.9 Hz), 7.18 (1H, dt, *J*=1.6, 6.9 Hz), 7.36 (1H, dd, *J*=1.6, 6.9 Hz), 7.42 (1H, dd, *J*=1.6, 6.9 Hz), 9.11 (1H, br s). High resolution MS *m/z*: Calcd for C<sub>12</sub>H<sub>9</sub>F<sub>3</sub>N<sub>2</sub>O: 254.0666. Found: 254.0662. **4**: mp 223.0–225.0 °C (decomp., colorless prisms, recrystallized from CHCl<sub>3</sub>). IR (KBr): 3365, 1676, 1605, 1482, 1465, 1453, 1205, 1183, 1179, 745 cm<sup>-1</sup>. <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 2.60 (1/11H, dd, *J*=12.5, 5.6 Hz), 2.76 (10/11H, dd, *J*=12.5, 5.6 Hz), 2.81 (1/11H, td, *J*=12.5, 7.8 Hz), 2.93 (10/11H, td, *J*=12.5, 7.8 Hz), 3.34 (1/11H, td, *J*=12.5, 5.6 Hz), 3.45 (10/11H, td, *J*=12.5, 5.6 Hz), 4.10 (10/11H, m), 4.32 (1/11H, m), 4.54 (1/11H, s, disappeared on addition of D<sub>2</sub>O), 5.30 (10/11H, s, disappeared on addition of D<sub>2</sub>O), 5.63 (10/11H, s), 5.75 (1/11H, s), 6.48 (10/11H, dd, *J*=2.2, 0.7 Hz), 6.50 (1/11H, dd, *J*=2.2, 0.7 Hz), 6.76 (1/11H, d, *J*=7.6 Hz), 6.78 (10/11H, d, *J*=7.6 Hz), 6.83 (10/11H, dt, *J*=7.6, 1.0 Hz), 6.86 (1/11H, dt, *J*=7.6, 1.0 Hz), 7.07 (10/11H, dt, *J*=7.6, 1.0 Hz), 7.10 (10/11H, br d, *J*=7.6 Hz), 7.13 (10/11H, td, *J*=7.6, 1.0 Hz), 7.19 (10/11H, td, *J*=7.6, 1.0 Hz), 7.22 (10/11H, dd, *J*=7.6, 1.0 Hz), 7.07–7.24 (5/11H, m), 7.56 (10/11H, dd, *J*=7.6, 0.7 Hz), 7.58 (1/11H, dd, *J*=7.6, 0.7 Hz), 7.77 (1/11H, br s), 7.94 (10/11H, br s). High-resolution MS *m/z*: Calcd for C<sub>20</sub>H<sub>16</sub>F<sub>3</sub>N<sub>3</sub>O: 371.1246. Found: 371.1244.



**5**: colorless oil. IR (film): 3405, 1681, 1467, 1460, 1204, 1185, 1145, 744  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 2.47—2.53 (1/6H, m), 2.64—2.70 (5/6H, m), 2.87—2.96 (1/6H, m), 3.02—3.11 (5/6H, m), 3.32—3.40 (1/6H, m), 3.45—3.52 (5/6H, m), 4.19 (5/6H, m), 4.30—4.36 (1/6H, m), 4.71 (1/6H, br s, disappeared on addition of  $\text{D}_2\text{O}$ ), 5.25 (5/6H, br s, disappeared on addition of  $\text{D}_2\text{O}$ ), 5.92 (5/6H, s), 6.00 (1/6H, br s), 6.90 (5/6H, d,  $J=2.5$  Hz), 6.93 (1/6H, d,  $J=2.5$  Hz), 7.06—7.12 (1/6H, m), 7.09 (5/6H, ddd,  $J=8.1, 7.1, 1.0$  Hz), 7.12—7.26 (8/6H, m), 7.16 (5/6H, td,  $J=7.6, 1.2$  Hz), 7.22 (5/6H, d,  $J=7.6$  Hz), 7.36 (5/6H, d,  $J=8.1$  Hz), 7.38 (1/6H, d,  $J=8.1$  Hz), 7.39 (1/6H, d,  $J=8.1$  Hz), 7.54 (5/6H, d,  $J=8.1$  Hz), 8.02 (5/6H, br s), 8.05 (1/6H, br s). High-resolution MS  $m/z$ : Calcd for  $\text{C}_{20}\text{H}_{16}\text{F}_3\text{N}_3\text{O}$ : 371.1245. Found: 371.1246. **6**: mp 114.5—115.5  $^\circ\text{C}$  (colorless needles, recrystallized from  $\text{CH}_2\text{Cl}_2$ –hexane). IR (KBr): 3430, 3340, 1700, 1563, 1349, 1206, 1172, 1119, 976, 952, 870  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 3.04 (2H, t,  $J=6.6$  Hz), 3.15 (3H, s), 3.67 (2H, q,  $J=6.6$  Hz), 6.37 (1H, br s), 7.05 (1H, d,  $J=8.8$  Hz), 7.11 (1H, s), 7.37 (1H, s), 7.58 (1H, d,  $J=8.8$  Hz), 8.26 (1H, br s). High resolution MS  $m/z$ : Calcd for  $\text{C}_{13}\text{H}_{13}\text{F}_3\text{N}_2\text{O}_4\text{S}$ : 350.0547. Found: 350.0539. **7**: very pale yellow oil. IR (film): 3402, 1709, 1213, 1180, 1167, 746  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  ( $\text{DMSO}-d_6$ )  $\delta$ : 3.00 (2H, tm,  $J=8.1$  Hz), 3.39—3.41 (2H, m), 7.02 (1H, ddd,  $J=8.0, 7.0, 1.1$  Hz), 7.07 (1H, ddd,  $J=8.0, 7.0, 1.1$  Hz), 7.11 (1H, ddd,  $J=8.0, 7.0, 1.1$  Hz), 7.19 (1H, ddd,  $J=8.0, 7.0, 1.1$  Hz), 7.38 (1H, d,  $J=8.0$  Hz), 7.48 (1H, d,  $J=8.0$  Hz), 7.55 (1H, d,  $J=8.0$  Hz), 7.64 (1H, d,  $J=2.4$  Hz), 7.72 (1H, d,  $J=8.0$  Hz), 9.62 (1H, br t,  $J=5.6$  Hz disappeared on addition of  $\text{D}_2\text{O}$ ), 11.0 (1H, s), 11.5 (1H, br s, disappeared on addition of  $\text{D}_2\text{O}$ ). High-resolution MS  $m/z$ : Calcd for  $\text{C}_{20}\text{H}_{16}\text{F}_3\text{N}_3\text{O}$ : 371.1245. Found: 371.1248. **8**: colorless oil. IR (film): 3370, 2940, 1694, 1612, 1486, 1471, 1255, 1206, 1145, 1101, 1066, 750  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 1.60—1.69 (2H, m), 1.75—1.83 (2H, m), 2.34—2.41 (2/6H, m), 2.47—2.59 (10/6H, m), 3.15 (1H, dt,  $J=8.8, 6.4$  Hz), 3.30 (1H, dt,  $J=8.8, 6.4$  Hz), 3.36 (1H, dt,  $J=6.4, 11.2$  Hz), 3.49 (2H, t,  $J=7.8$  Hz), 3.95—3.98 (5/6H, m), 4.14—4.18 (1/6H, m), 5.52 (5/6H, s), 5.64 (1/6H, d,  $J=2.0$  Hz), 6.65 (1H, d,  $J=7.8$  Hz), 6.85 (1H, t,  $J=7.8$  Hz), 7.22 (1H, t,  $J=7.8$  Hz), 7.23 (1H, d,  $J=7.8$  Hz). High resolution MS  $m/z$ : Calcd for  $\text{C}_{16}\text{H}_{18}\text{ClF}_3\text{N}_2\text{O}_2$ : 364.0978 and 362.1008. Found: 364.1003 and 362.1022.

**[Entry 1]** A solution of MsCl (67.0 mg, 0.59 mmol) in benzene (1.0 mL) was added to a solution of **2** (119.0 mg, 0.44 mmol) and indole (155.0 mg, 1.32 mmol) in benzene (3.0 mL) and  $\text{Et}_3\text{N}$  (0.4 mL, 2.87 mmol) at 0  $^\circ\text{C}$  and the mixture was stirred at 0 $^\circ\text{C}$  for 1 h. After the same work-up and separation as described in Entry 4, **3** (19.9 mg, 18%), **5** (6.7 mg, 4%), and **7** (6.7 mg, 4%) were obtained in the order of elution.

**[Entry 2]** A solution of MsCl (73.4 mg, 0.64 mmol) in anhydrous  $\text{CHCl}_3$  (1.0 mL) was added to a solution of **2** (113.5 mg, 0.42 mmol) and indole (146.1 mg, 1.25 mmol) in anhydrous  $\text{CHCl}_3$  (3.0 mL) and  $\text{Et}_3\text{N}$  (0.4 mL, 2.87 mmol) at 0  $^\circ\text{C}$  and the mixture was stirred at 0 $^\circ\text{C}$  for 1 h. After the same work-up and separation as described in Entry 4, **3** (15.2 mg, 14%), **4** (7.1 mg, 5%), **5** (33.1 mg, 21%), and **6** (5.7 mg,

4%) were obtained in the order of elution.

**[Entry 3]** A solution of MsCl (60.3 mg, 0.53 mmol) in anhydrous ClCH<sub>2</sub>CH<sub>2</sub>Cl (1.0 mL) was added to a solution of **2** (111.6 mg, 0.41 mmol) and indole (143.7 mg, 1.23 mmol) in anhydrous ClCH<sub>2</sub>CH<sub>2</sub>Cl (3.0 mL) and Et<sub>3</sub>N (0.4 mL, 2.87 mmol) at 0 °C and the mixture was stirred at 0 °C for 1 h. After the same work-up and separation as described in Entry 4, **3** (13.8 mg, 13%), **4** (7.6 mg, 5%), **5** (16.4 mg, 11%), and **6** (3.8 mg, 3%) were obtained in the order of elution.

**[Entry 5]** A solution of MsCl (59.4 mg, 0.52 mmol) in anhydrous DMF (1.0 mL) was added to a solution of **2** (101.0 mg, 0.37 mmol) and indole (131.9 mg, 1.13 mmol) in anhydrous DMF (3.0 mL) and Et<sub>3</sub>N (0.4 mL, 2.87 mmol) at 0 °C and the mixture was stirred at 0 °C for 1 h. After the same work-up and separation as described in Entry 4, **3** (28.2 mg, 30%), **4** (1.6 mg, 1%), **5** (9.8 mg, 7%), and **6** (1.9 mg, 2%) were obtained in the order of elution.

**[Entry 6]** A solution of MsCl (59.2 mg, 0.52 mmol) in MeCN (1.0 mL) was added to a solution of **2** (107.8 mg, 0.39 mmol) and indole (137.8 mg, 1.18 mmol) in MeCN (3.0 mL) and Et<sub>3</sub>N (0.4 mL, 2.87 mmol) at 0 °C and the mixture was stirred at 0 °C for 1 h. After the same work-up and separation as described in Entry 4, **3** (10.0 mg, 10%), **4** (1.5 mg, 1%), **5** (12.2 mg, 8%), and **1** (5.8 mg, 6%) were obtained in the order of elution.

**[Entry 7]** A solution of MsCl (57.5 mg, 0.50 mmol) in anhydrous MeNHCHO (1.0 mL) was added to a solution of **2** (108.2 mg, 0.39 mmol) and indole (140.5 mg, 1.20 mmol) in anhydrous MeNHCHO (3.0 mL) and Et<sub>3</sub>N (0.4 mL, 2.87 mmol) at 0 °C and the mixture was stirred at 0 °C for 1 h. After the same work-up and separation as described in Entry 4, **3** (1.7 mg, 2%), **4** (1.1 mg, 1%), **5** (4.4 mg, 4%), (3a,8a-*cis*)-1,2,3,3a,8,8a-hexahydro-3a-hydroxy-1-trifluoroacetylpyrrolo[2,3-*b*]indole (**9**) (21.2 mg, 20%), and unreacted **2** (22.0 mg, 20%) were obtained in the order of elution. **9**: mp 115.0—115.5 °C (colorless prisms, recrystallized from CHCl<sub>3</sub>–hexane). IR (KBr): 3336, 3282, 1697, 1685, 1469, 1250, 1201, 1147, 750 cm<sup>-1</sup>. <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>, 120°C) δ: 2.14—2.53 (2H, m), 3.22—3.43 (1H, m), 3.84—4.03 (1H, m), 5.36 (1H, br s), 5.56 (1H, br s), 6.26 (1H, br s), 6.60 (1H, d, *J*=7.6 Hz), 6.69 (1H, br t, *J*=7.6 Hz), 7.06 (1H, t, *J*=7.6 Hz), 7.21 (1H, d, *J*=7.6 Hz). High-resolution MS *m/z*: Calcd for C<sub>12</sub>H<sub>11</sub>F<sub>3</sub>N<sub>2</sub>O<sub>2</sub>: 272.0773. Found: 272.0772.

**[Entry 8]** A solution of MsCl (63.3 mg, 0.55 mmol) in anhydrous CHCl<sub>3</sub> (1.0 mL) was added to a solution of **2** (111.7 mg, 0.41 mmol) and indole (481.0 mg, 4.11 mmol) in anhydrous CHCl<sub>3</sub> (3.0 mL) and Et<sub>3</sub>N (0.4 mL, 2.87 mmol) at 0 °C and the mixture was stirred at 0 °C for 1 h. After the same work-up and separation as described in Entry 4, **3** (4.4 mg, 4%), **4** (10.3 mg, 7%), **5** (46.1 mg, 30%), and **6** (1.9 mg, 1%) were obtained in the order of elution.

**[Entry 9]** A solution of MsCl (57.4 mg, 0.50 mmol) in anhydrous ClCH<sub>2</sub>CH<sub>2</sub>Cl (1.0 mL) was added to a solution of **2** (103.8 mg, 0.38 mmol) and indole (444.7 mg, 3.80 mmol) in anhydrous ClCH<sub>2</sub>CH<sub>2</sub>Cl (3.0

mL) and Et<sub>3</sub>N (0.4 mL, 2.87 mmol) at 0 °C and the mixture was stirred at 0 °C for 1 h. After the same work-up and separation as described in Entry 4, **3** (7.4 mg, 8%), **4** (7.3 mg, 5%), **5** (25.5 mg, 18%), and **6** (2.3 mg, 2%) were obtained in the order of elution.

**[Entry 10]** A solution of MsCl (59.3 mg, 0.52 mmol) in anhydrous EtOAc (1.0 mL) was added to a solution of **2** (111.7 mg, 0.41 mmol) and indole (479.8 mg, 4.10 mmol) in anhydrous EtOAc (3.0 mL) and Et<sub>3</sub>N (0.4 mL, 2.87 mmol) at 0 °C and the mixture was stirred at 0 °C for 1 h. After the same work-up and separation as described in Entry 4, **3** (6.4 mg, 6%), **4** (11.1 mg, 7%), **5** (38.6 mg, 25%), and **6** (4.9 mg, 3%) were obtained in the order of elution.

**7 from 5** — A solution of **5** (10.0 mg, 0.03 mmol) in DMSO (2.0 mL) was stirred at 130 °C for 3 h. After addition of H<sub>2</sub>O and EtOAc, the organic layer was washed with H<sub>2</sub>O and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and evaporated under reduced pressure to leave an oil, which was column-chromatographed on SiO<sub>2</sub> with EtOAc–hexane (1:5, v/v) to give **7** (8.9 mg, 89%).

**(3a,8a-cis)-1,2,3,3a,8,8a-Hexahydro-1-trifluoroacetyl-3a-(2,3,4-trimethoxyphenyl)pyrrolo[2,3-b]indole (10), and (3a,8a-cis)-1,2,3,3a,8,8a-hexahydro-1-trifluoroacetyl-3a-[3-(Nb-trifluoroacetyl)-aminoethylindol-1-yl]pyrrolo[2,3-b]indole (11a) from 2** — A solution of MsCl (55.1 mg, 0.48 mmol) in anhydrous CHCl<sub>3</sub> (1.0 mL) was added to a solution of **2** (100.1 mg, 0.37 mmol) and 1,2,3-trimethoxybenzene (619.0 mg, 3.69 mmol) in anhydrous CHCl<sub>3</sub> (3.0 mL) and Et<sub>3</sub>N (0.4 mL, 2.87 mmol) at 0 °C and the mixture was stirred at 0 °C for 1 h. After addition of H<sub>2</sub>O under ice cooling, the whole was extracted with CHCl<sub>3</sub>–MeOH (95:5, v/v). The extract was washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and evaporated under reduced pressure to leave an oil, which was column-chromatographed on SiO<sub>2</sub> successively with CHCl<sub>3</sub>–hexane (1:1, v/v), CHCl<sub>3</sub>–MeOH (98:2, v/v), and EtOAc–hexane (1:5, v/v) to give **3** (14.2 mg, 15%), **10** (2.4 mg, 2%), **11a** (11.8 mg, 13%), **1** (4.0 mg, 4%), and **6** (5.4 mg, 4%) in the order of elution. **10**: colorless oil. IR (film): 1684, 1466, 1203, 1144, 1103, 752 cm<sup>-1</sup>. <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>, 120 °C) δ: 2.36–2.58 (1H, m), 2.71–2.92 (1H, m), 3.19–3.34 (1H, m), 3.65 (3H, br s), 3.74 (3H, s), 3.77 (3H, s), 3.90–4.10 (1H, m), 5.86 (1H, br s), 6.37 (1H, br s), 6.61–6.70 (2H, m), 6.66 (1H, d, *J*=8.8 Hz), 6.87 (1H, d, *J*=8.8 Hz), 7.01 (1H, t, *J*=7.3 Hz), 6.99–7.10 (1H, m). High-resolution MS *m/z*: Calcd for C<sub>21</sub>H<sub>21</sub>F<sub>3</sub>N<sub>2</sub>O<sub>4</sub>: 422.1453. Found: 422.1448. **11a**: colorless oil. IR (film): 1689, 1209, 1186, 1153, 752 cm<sup>-1</sup>. <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>, 120 °C) δ: 2.58–3.03 (4H, m), 3.27–3.47 (1H, m), 3.42 (2H, q, *J*=6.6 Hz), 4.02–4.12 (1H, m), 5.87 (1H, br s), 6.64–6.75 (3H, m), 7.00–7.07 (2H, m), 7.11 (1H, t, *J*=8.2 Hz), 7.16 (1H, t, *J*=8.2 Hz), 7.21 (1H, d, *J*=8.2 Hz), 7.26 (1H, d, *J*=8.2 Hz), 7.49 (1H, d, *J*=8.2 Hz), 8.96 (1H, br s). High-resolution MS *m/z*: Calcd for C<sub>24</sub>H<sub>20</sub>F<sub>6</sub>N<sub>4</sub>O<sub>2</sub>: 510.1490. Found: 510.1486.

**(3a,8a-cis)-1,2,3,3a,8,8a-Hexahydro-3a-[3-(Nb-trifluoroacetyl)aminoethylindol-1-yl]pyrrolo[2,3-b]indole (11b) from (11a)** — Sat. aq. NaHCO<sub>3</sub> (2.0 mL, 2.1 mmol) was added to a solution of **11a** (25.3 mg, 0.05 mmol) in MeOH (4.0 mL) and the mixture was stirred at rt for 3 h. After addition of H<sub>2</sub>O, the

whole was extracted with  $\text{CHCl}_3$ . The extract was washed with brine, dried over  $\text{Na}_2\text{SO}_4$ , and evaporated under reduced pressure to leave an oil. Then  $\text{H}_2\text{O}$  layer was evaporated under reduced pressure to leave an oil. These oils were combined and column-chromatographed on  $\text{SiO}_2$  with  $\text{CHCl}_3$ -MeOH (97:3, v/v) to give **11b** (13.8 mg, 67%). **11b**: colorless oil. IR (film): 1709, 1213, 1182, 1161, 746  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 2.36—2.42 (1H, m), 2.61—2.70 (1H, m), 2.85 (2H, t,  $J=6.6$  Hz), 2.94—3.02 (1H, m), 3.22—3.30 (1H, m), 3.48—3.60 (2H, m), 4.20 (1H, br s, disappeared on addition of  $\text{D}_2\text{O}$ ), 5.31 (1H, s), 6.26 (1H, br s), 6.59 (1H, d,  $J=7.7$  Hz), 6.64 (1H, s), 6.83 (1H, t,  $J=7.7$  Hz), 7.11 (1H, t,  $J=7.7$  Hz), 7.19 (1H, t,  $J=7.7$  Hz), 7.24 (1H, t,  $J=7.7$  Hz), 7.33 (1H, d,  $J=7.7$  Hz), 7.38 (1H, d,  $J=7.7$  Hz), 7.46 (1H, d,  $J=7.7$  Hz). High-resolution MS  $m/z$ : Calcd for  $\text{C}_{22}\text{H}_{21}\text{F}_3\text{N}_4\text{O}$ : 414.1668. Found: 414.1647.

**(3a,8a-cis)-1,2,3,3a,8,8a-Hexahydro-3a-(4-methoxyphenyl)-1-trifluoroacetylpyrrolo[2,3-b]indole**

**(12) from 2** — A solution of  $\text{MsCl}$  (56.8 mg, 0.49 mmol) in anhydrous  $\text{CHCl}_3$  (1.0 mL) was added to a solution of **2** (99.4 mg, 0.37 mmol) and anisole (2 mL, 18.4 mmol) in anhydrous  $\text{CHCl}_3$  (1.0 mL) and  $\text{Et}_3\text{N}$  (0.4 mL, 2.87 mmol) at 0 °C and the mixture was stirred at 0 °C for 1 h. After addition of  $\text{H}_2\text{O}$  under ice cooling, the whole was extracted with  $\text{CHCl}_3$ -MeOH (95:5, v/v). The extract was washed with brine, dried over  $\text{Na}_2\text{SO}_4$ , and evaporated under reduced pressure to leave an oil, which was column-chromatographed on  $\text{SiO}_2$  with  $\text{EtOAc}$ -hexane (1:7, v/v) to give **3** (16.5 mg, 18%), **12** (6.0 mg, 5%), **11a** (9.6 mg, 10%), and **1** (6.0 mg, 6%) in the order of elution. **12**: colorless oil. IR (film): 1684, 1252, 1186, 1144  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 2.52—2.58 (1/11H, m), 2.61—2.67 (1/11H, m), 2.67—2.80 (20/11H, m), 3.22—3.31 (1/11H, m), 3.38 (10/11H, td,  $J=11.3, 6.2$  Hz), 3.79 (30/11H, s), 3.88 (3/11H, s), 4.01—4.09 (10/11H, m), 4.24—4.30 (1/11H, m), 4.65 (1/11H, br s, disappeared on addition of  $\text{D}_2\text{O}$ ), 5.22 (10/11H, br s, disappeared on addition of  $\text{D}_2\text{O}$ ), 5.65 (10/11H, s), 5.71 (1/11H, br s), 6.68 (1H, d,  $J=7.5$  Hz), 6.79 (10/11H, td,  $J=7.5, 0.8$  Hz), 6.77—6.87 (1/11H, m), 6.83 (20/11H, dm,  $J=8.9$  Hz), 6.85 (2/11H, dm,  $J=8.9$  Hz), 7.01 (1/11H, d,  $J=7.5$  Hz), 7.06 (10/11H, d,  $J=7.5$  Hz), 7.12 (10/11H, td,  $J=7.5, 1.3$  Hz), 7.10—7.15 (1/11H, m), 7.20 (2/11H, dm,  $J=8.9$  Hz), 7.26 (20/11H, dm,  $J=8.9$  Hz). High-resolution MS  $m/z$ : Calcd for  $\text{C}_{19}\text{H}_{17}\text{F}_3\text{N}_2\text{O}_2$ : 362.1242. Found: 362.1244.

**(3a,8a-cis)-1,2,3,3a,8,8a-Hexahydro-3a-(4-methoxyphenyl)pyrrolo[2,3-b]indole (13) from 12** — Sat. aq.  $\text{NaHCO}_3$  (0.5 mL, 0.53 mmol) was added to a solution of **12** (6.2 mg, 0.02 mmol) in MeOH (1.0 mL) and the mixture was refluxed for 40 min with stirring. The solvent was evaporated under reduced pressure to leave a solid, which was column-chromatographed on  $\text{SiO}_2$  with  $\text{CHCl}_3$ -MeOH (95:5, v/v) to give **13** (4.3 mg, 94%). **13**: pale yellow oil. IR (film): 2929, 1606, 1512, 1250, 746  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 1.56—2.05 (2H, m, disappeared on addition of  $\text{D}_2\text{O}$ ), 2.41 (1H, dd,  $J=11.5, 5.1$  Hz), 2.49 (1H, td,  $J=11.5, 6.6$  Hz), 2.81 (1H, td,  $J=11.5, 5.1$  Hz), 3.21 (1H, dd,  $J=11.5, 6.6$  Hz), 3.77 (3H, s), 5.12 (1H, s), 6.63 (1H, d,  $J=7.6$  Hz), 6.70 (1H, t,  $J=7.6$  Hz), 6.82 (2H, dm,  $J=8.7$  Hz), 6.93 (1H, dd,  $J=7.6, 0.9$  Hz), 7.04 (1H, td,

$J=7.6, 0.9$  Hz), 7.25 (2H, dm,  $J=8.7$  Hz). High-resolution MS  $m/z$ : Calcd for  $C_{17}H_{18}N_2O$ : 266.1419. Found: 266.1412.

**(3a,8a-cis)-1,2,3,3a,8,8a-Hexahydro-3a-(4-hydroxyphenyl)- (14) and -3a-(2-hydroxyphenyl)-1-trifluoroacetylpyrrolo[2,3-*b*]indole (15) from 2** — A solution of MsCl (226.7 mg, 1.99 mmol) in anhydrous  $CHCl_3$  (1.0 mL) was added to a solution of **2** (107.2 mg, 0.39 mmol) and phenol (370.5 mg, 3.94 mmol) in anhydrous  $CHCl_3$  (3.0 mL) and  $Et_3N$  (0.27 mL, 1.94 mmol) at 0 °C and the mixture was stirred at 0 °C for 1 h. After addition of  $H_2O$  under ice cooling, the whole was extracted with  $CHCl_3$ -MeOH (95:5, v/v). The extract was washed with brine, dried over  $Na_2SO_4$ , and evaporated under reduced pressure to leave an oil, which was column-chromatographed on  $SiO_2$  successively with  $CHCl_3$ -hexane (2:1, v/v) and EtOAc-hexane (1:5, v/v) to give **11a** (8.0 mg, 8%), **1** (7.6 mg, 8%), **15** (7.4 mg, 5%), **14** (4.0mg, 3%), and **6** (5.1 mg, 4%) in the order of elution. **14**: colorless oil. IR (film): 1678, 1203, 1188, 1151, 754  $cm^{-1}$ .  $^1H$ -NMR ( $CDCl_3$ )  $\delta$ : 2.66—2.79 (2H, m), 3.32—3.41 (1H, m), 4.00—4.09 (1H, m), 4.91 (1H, br s, disappeared on addition of  $D_2O$ ), 5.21 (1H, br s, disappeared on addition of  $D_2O$ ), 5.64 (1H, s), 6.68 (1H, d,  $J=7.6$  Hz), 6.76 (2H, m), 6.79 (1H, td,  $J=7.6, 0.6$  Hz), 7.05 (1H, d,  $J=7.6$  Hz), 7.12 (1H, td,  $J=7.6, 1.1$  Hz), 7.26 (2H, m). High-resolution MS  $m/z$ : Calcd for  $C_{18}H_{15}F_3N_2O_2$ : 348.1086. Found: 348.1086. **15**: colorless oil. IR (film): 1709, 1211, 1184, 1165, 750  $cm^{-1}$ .  $^1H$ -NMR ( $CDCl_3$ )  $\delta$ : 2.34—2.43 (1H, m), 2.43—2.52 (1H, m), 3.22—3.32 (1H, m), 3.35—3.45 (1H, m), 5.04 (1H, br s), 6.21 (1H, d,  $J=1.7$  Hz, collapsed to s on addition of  $D_2O$ ), 6.31 (1H, br s), 6.70 (1H, d,  $J=7.6$  Hz), 6.78 (1H, d,  $J=7.6$  Hz), 6.81 (1H, td,  $J=7.6, 0.9$  Hz), 6.91 (1H, td,  $J=7.6, 0.9$  Hz), 7.09 (1H, td,  $J=7.6, 1.3$  Hz), 7.12 (1H, td,  $J=7.6, 1.3$  Hz), 7.19 (1H, d,  $J=7.6$  Hz), 7.32 (1H, dd,  $J=7.6, 1.3$  Hz). High-resolution MS  $m/z$ : Calcd for  $C_{18}H_{15}F_3N_2O_2$ : 348.1085. Found: 348.1084.

**12 from 14** — Excess  $CH_2N_2$  in  $Et_2O$  was added to a solution of **14** (3.7 mg, 0.01 mmol) in MeOH (0.5 mL) and the mixture was stirred at rt for 30 min. The solvent was evaporated under reduced pressure to leave an oil, which was column-chromatographed on  $SiO_2$  with  $CHCl_3$ -hexane (2:1, v/v) to give **12** (2.8 mg, 73%).

**(3a,8a-cis)-1,2,3,3a,8,8a-Hexahydro-3a-(pyrrol-2-yl)-1-trifluoroacetylpyrrolo[2,3-*b*]indole (16) from 2** — A solution of MsCl (60.1 mg, 0.53 mmol) in anhydrous  $CHCl_3$  (1.0 mL) was added to a solution of **2** (106.2 mg, 0.39 mmol) and pyrrole (263.6 mg, 3.93 mmol) in anhydrous  $CHCl_3$  (3.0 mL) and  $Et_3N$  (0.4 mL, 2.87 mmol) at 0 °C and the mixture was stirred at 0 °C for 1 h. After addition of  $H_2O$  under ice cooling, the whole was extracted with  $CHCl_3$ -MeOH (95:5, v/v). The extract was washed with brine, dried over  $Na_2SO_4$ , and evaporated under reduced pressure to leave an oil, which was column-chromatographed on  $SiO_2$  with EtOAc-hexane (1:5, v/v) and  $CHCl_3$ -MeOH (99:1, v/v) to give **3** (5.6mg, 6%), **16** (36.5mg, 29%), **11a** (8.3 mg, 8%), and **1** (6.9mg, 7%) in the order of elution. **16**:

colorless oil. IR (film): 1684, 1483, 1468, 1205, 1188, 1747, 754  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  ( $\text{DMSO-}d_6$ )  $\delta$ : 2.34—2.70 (2H, m), 3.04—3.13 (1/5H, m), 3.16—3.27 (4/5H, m), 3.90—3.98 (4/5H, m), 3.99—4.06 (1/5H, m), 5.61 (4/5H, s), 5.62—5.67 (1H, m), 5.67—5.71 (1/5H, m), 5.83—5.89 (1H, m), 6.62 (4/5H, d,  $J=7.6$  Hz), 6.65 (1/5H, d,  $J=7.6$  Hz), 6.67—6.76 (3H, m), 7.04 (4/5H, td,  $J=7.6, 1.2$  Hz), 7.07 (1/5H, td,  $J=7.6, 1.2$  Hz), 7.41 (1/5H, d,  $J=7.6$  Hz), 7.22 (4/5H, d,  $J=7.6$  Hz), 10.81 (1H, br s). High-resolution MS  $m/z$ : Calcd for  $\text{C}_{16}\text{H}_{14}\text{F}_3\text{N}_3\text{O}$ : 321.1089. Found: 321.1083.

**(3a,8a-cis)-8-Acetyl-1,2,3,3a,8,8a-hexahydro-3a-(pyrrol-2-yl)-1-trifluoroacetylpyrrolo[2,3-b]indole (18) from 16** —  $\text{Ac}_2\text{O}$  (2.0 mL) was added to a solution of **16** (36.2 mg, 0.11 mmol) in pyridine (2.0 mL) and the mixture was stirred at 65 °C for 10 h. The solvent was evaporated under reduced pressure to leave an oil, which was column-chromatographed on  $\text{SiO}_2$  with EtOAc–hexane (1:4, v/v) to give **18** (33.9 mg, 83%). **18**: mp 218.0—220.0 °C (colorless powder, recrystallized from EtOAc–hexane). IR (KBr): 3263, 1705, 1662, 1151, 760  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 2.47 (3H, s), 2.62 (1H, dd,  $J=12.7, 5.3$  Hz), 2.78 (1H, td,  $J=12.7, 7.5$  Hz), 3.22 (1H, td,  $J=12.7, 5.3$  Hz), 4.02 (1H, m), 6.02 (1H, s), 6.16—6.22 (2H, m), 6.74—6.78 (1H, m), 7.19—7.28 (2H, m), 7.39 (1H, ddd,  $J=8.1, 7.1, 1.8$  Hz), 7.79 (1H, br s, disappeared on addition of  $\text{D}_2\text{O}$ ), 8.06 (1H, d,  $J=8.1$  Hz). *Anal.* Calcd for  $\text{C}_{18}\text{H}_{16}\text{F}_3\text{N}_3\text{O}_2$ : C, 59.50; H, 4.44; N, 11.57. Found: C, 59.61; H, 4.43; N, 11.56.

**(3a,8a-cis)-8-Acetyl-1,2,3,3a,8,8a-hexahydro-3a-(indol-3-yl)-1-trifluoroacetylpyrrolo[2,3-b]indole (19) from 5** —  $\text{Ac}_2\text{O}$  (3.0 mL) was added to a solution of **5** (22.0 mg, 0.06 mmol) in pyridine (3.0 mL) and the mixture was stirred at 62 °C for 9.5 h. The solvent was evaporated under reduced pressure to leave an oil, which was column-chromatographed on  $\text{SiO}_2$  with EtOAc–hexane (1:2, v/v) to give **19** (12.5 mg, 51%). **19**: mp 219.0—220.5 °C (colorless prisms, recrystallized from  $\text{CHCl}_3$ ). IR (KBr): 3360, 1679, 1479, 1462, 1388, 1206, 1142, 740  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 2.41 (3H, s), 2.58 (1H, dd,  $J=12.5, 5.1$  Hz), 3.04 (1H, td,  $J=12.5, 7.3$  Hz), 3.30 (1H, td,  $J=12.5, 5.1$  Hz), 4.09 (1H, m), 6.43 (1H, br s), 6.71 (1H, d,  $J=2.7$  Hz), 7.13 (1H, t,  $J=8.1$  Hz), 7.22 (1H, t,  $J=8.1$  Hz), 7.23 (1H, t,  $J=8.1$  Hz), 7.30—7.34 (2H, m), 7.38 (1H, t,  $J=8.1$  Hz), 7.39 (1H, d,  $J=8.1$  Hz), 8.10 (1H, br s), 8.16 (1H, br s). High-resolution MS  $m/z$ : Calcd for  $\text{C}_{22}\text{H}_{18}\text{F}_3\text{N}_3\text{O}_2$ : 413.1351. Found: 413.1353. *Anal.* Calcd for  $\text{C}_{22}\text{H}_{18}\text{F}_3\text{N}_3\text{O}_2 \cdot 1/4\text{H}_2\text{O}$ : C, 63.23; H, 4.34; N, 10.05. Found: C, 63.00; H, 4.37; N, 9.81.

**(3a,8a-cis)-8-Acetyl-1,2,3,3a,8,8a-hexahydro-3a-(indol-2-yl)-1-trifluoroacetylpyrrolo[2,3-b]indole (20) from 4** —  $\text{Ac}_2\text{O}$  (2.0 mL) was added to a solution of **4** (20.5 mg, 0.03 mmol) in pyridine (2.0 mL) and the mixture was stirred at 63 °C for 10 h. The solvent was evaporated under reduced pressure to leave an oil, which was column-chromatographed on  $\text{SiO}_2$  with EtOAc–hexane (1:5, v/v) to give **20** (22.1 mg, 97 %). **20**: mp 147.0—150.0 °C (colorless fine needles, recrystallized from  $\text{CHCl}_3$ –hexane). IR (KBr): 1709, 1662, 1479, 1394, 1147, 1142, 1124, 750  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 2.46 (3H, s), 2.71 (1H, dd,

$J=12.5$ , 5.1 Hz), 2.91 (1H, td,  $J=12.5$ , 7.2 Hz), 3.27 (1H, td,  $J=12.5$ , 5.1 Hz), 4.09 (1H, m), 6.19 (1H, s), 6.51 (1H, dd,  $J=2.2$ , 0.7 Hz), 7.11 (1H, ddd,  $J=8.1$ , 7.0, 1.1 Hz), 7.17 (1H, ddd,  $J=8.1$ , 7.0, 1.1 Hz), 7.21—7.29 (3H, m), 7.38—7.45 (1H, m), 7.58 (1H, dd,  $J=8.1$ , 1.1 Hz), 7.94 (1H, br s), 8.11 (1H, d,  $J=8.1$ Hz). High-resolution MS  $m/z$ : Calcd for  $C_{22}H_{18}F_3N_3O_2$ : 413.1351. Found: 413.1351.

**1-Acetyl-6-mesyloxy-Nb-trifluoroacetyltryptamine (21) from 6** — Reported in our previous paper.<sup>9</sup>

**(3a,8a-cis)-3a-Acetoxy- (22) and (3a,8a-cis)-3a-acetoxy-8-acetyl-1,2,3,3a,8,8a-hexahydro-1-trifluoroacetylpyrrolo[2,3-b]indole (23) from 9** —  $Ac_2O$  (5.0 mL) was added to a solution of **9** (40.9 mg, 0.15 mmol) in pyridine (5.0 mL) and the mixture was stirred at rt for 18 h. The solvent was evaporated under reduced pressure to leave an oil, which was column-chromatographed on  $SiO_2$  successively with  $CHCl_3$ –hexane (1:2, v/v) and  $CHCl_3$  to give **22** (33.5 mg, 71%) and **23** (9.0 mg, 17%) in the order of elution. **22**: colorless oil. IR (film): 1741, 1693, 1240, 1205, 1146  $cm^{-1}$ .  $^1H$ -NMR ( $CDCl_3$ )  $\delta$ : 2.04 (12/5H, s), 2.05 (3/5H, s), 2.51—2.60 (1/5H, m), 2.68 (4/5H, ddd,  $J=12.9$ , 11.6, 8.3 Hz), 2.77 (1/5H, dd,  $J=12.4$ , 6.0 Hz), 3.04 (4/5H, ddd,  $J=12.9$ , 6.2, 1.5 Hz), 3.17 (1/5H, td,  $J=12.4$ , 6.0 Hz), 3.41 (4/5H, td,  $J=11.6$ , 6.2 Hz), 4.02 (4/5H, m), 4.22 (1/5H, dd,  $J=12.4$ , 8.3 Hz), 4.81 (1/5H, br d,  $J=4.0$  Hz disappeared on addition of  $D_2O$ ), 5.18 (4/5H, br s, disappeared on addition of  $D_2O$ ), 5.81 (4/5H, d,  $J=2.0$  Hz, collapsed to s on addition of  $D_2O$ ), 5.95—5.98 (1/5H, m), 6.67 (4/5H, d,  $J=7.6$  Hz), 6.69 (1/5H, d,  $J=7.6$  Hz), 6.82 (4/5H, td,  $J=7.6$ , 1.1 Hz), 6.86 (1/5H, td,  $J=7.6$ , 1.1 Hz), 7.22 (4/5H, td,  $J=7.6$ , 1.3 Hz), 7.23 (1/5H, td,  $J=7.6$ , 1.3 Hz), 7.41 (1/5H, d,  $J=7.6$  Hz), 7.51(4/5H, d,  $J=7.6$  Hz). High-resolution MS  $m/z$ : Calcd for  $C_{14}H_{13}F_3N_2O_3$ : 314.0878. Found: 314.0881. **23**: mp 117.5—118.0 °C (colorless prisms, recrystallized from EtOAc–hexane). IR (KBr): 1745, 1701, 1685, 1373, 1242, 1133, 758  $cm^{-1}$ .  $^1H$ -NMR ( $CDCl_3$ )  $\delta$ : 2.05 (3H, s), 2.59 (3H, s), 2.59 (1H, td,  $J=12.7$ , 7.8 Hz), 2.90 (1H, dd,  $J=12.7$ , 5.1 Hz), 3.13 (1H, ddd,  $J=12.7$ , 11.7, 5.1 Hz), 4.00 (1H, m), 6.40 (1H, br s), 7.19 (1H, td,  $J=7.4$ , 1.0 Hz), 7.42 (1H, ddd,  $J=8.1$ , 7.4, 1.2 Hz), 7.53 (1H, dd,  $J=8.1$ , 1.0 Hz), 8.04 (1H, br d,  $J=7.4$  Hz). High-resolution MS  $m/z$ : Calcd for  $C_{16}H_{15}F_3N_2O_4$ : 356.0984. Found: 356.0994. *Anal.* Calcd for  $C_{16}H_{15}F_3N_2O_4$ : C, 53.94; H, 4.24; N, 7.86. Found: C, 53.98; H, 4.18; N, 7.62.

**23 from 22** —  $Ac_2O$  (5.0 mL) was added to a solution of **22** (33.5 mg, 0.10 mmol) in pyridine (5.0 mL) and the mixture was stirred at 55 °C for 32 h. The solvent was evaporated under reduced pressure to leave an oil, which was column-chromatographed on  $SiO_2$  with  $CHCl_3$ –hexane (1:1, v/v) to give unreacted **22** (8.7 mg, 26%) and **23** (23.7 mg, 62%) in the order of elution.

**(3a,8a-cis)-3a-Acetoxy-8-acety-1,2,3,3a,8,8a-hexahydro- (24) and (3a,8a-cis)-8-acety-1,2,3,3a,8,8a-hexahydro-3a-hydroxypyrrolo[2,3-b]indole (25) from 23** — Sat. aq.  $NaHCO_3$  (4.0 mL, 4.2 mmol) was added to a solution of **23** (39.7 mg, 0.11 mmol) in MeOH (5.0 mL) and the mixture was stirred at rt for 20 min. The solvent was evaporated under reduced pressure to leave a solid, which was column-chromatographed on  $SiO_2$  successively with  $CHCl_3$ –MeOH–AcOH (46:1:0.1, v/v) and

CHCl<sub>3</sub>–MeOH–AcOH (46:10:1, v/v) to give **24** (23.3 mg, 80%) and **25** (4.5 mg, 19%) in the order of elution. **24**: mp 125.0–126.0 °C (very pale yellow prisms, recrystallized from EtOAc–hexane). IR (KBr): 3315, 1739, 1649, 1483, 1408, 1238, 1047 cm<sup>-1</sup>. <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>) δ: 1.99 (3H, s), 2.24 (3H, s), 2.29–2.54 (3H, m), 2.97–3.09 (1H, m), 3.37 (1H, br s, disappeared on addition of D<sub>2</sub>O), 5.63 (1H, br d, *J*=2.2 Hz, collapsed to s on addition of D<sub>2</sub>O), 7.06 (1H, td, *J*=7.6, 1.1 Hz), 7.28 (1H, ddd, *J*=8.3, 7.6, 1.1 Hz), 7.45 (1H, dd, *J*=7.6, 1.1 Hz), 8.01 (1H, d, *J*=8.3 Hz). *Anal.* Calcd for C<sub>14</sub>H<sub>16</sub>N<sub>2</sub>O<sub>3</sub>: C, 64.60; H, 6.20; N, 10.76. Found: C, 64.50; H, 6.26; N, 10.63. **25**: mp 196.0–197.0 °C (colorless prisms, recrystallized from MeOH–EtOAc). IR (KBr): 3342, 3294, 1641, 1483, 1406, 762 cm<sup>-1</sup>. <sup>1</sup>H-NMR (CD<sub>3</sub>OD) δ: 2.28–2.34 (2H, m), 2.31 (3H, s), 2.53–2.63 (1H, m), 3.06–3.14 (1H, m), 5.25 (1H, s), 7.16 (1H, td, *J*=7.4, 1.0 Hz), 7.30 (1H, ddd, *J*=8.3, 7.4, 1.0 Hz), 7.44 (1H, d, *J*=7.4 Hz), 8.12 (1H, d, *J*=8.3 Hz). *Anal.* Calcd for C<sub>12</sub>H<sub>14</sub>N<sub>2</sub>O<sub>2</sub>: C, 66.03; H, 6.47; N, 12.84. Found: C, 66.01; H, 6.48; N, 12.82.

**25 from 23** — Sat. aq. NaHCO<sub>3</sub> (4.0 mL, 4.2 mmol) was added to a solution of **23** (40.2 mg, 0.11 mmol) in MeOH (5.0 mL) and the mixture was refluxed for 30 min with stirring. The solvent was evaporated under reduced pressure to leave a solid, which was column-chromatographed on SiO<sub>2</sub> with CHCl<sub>3</sub>–MeOH–AcOH (46:5:0.5, v/v) to give **25** (24.2 mg, 98%).

**(3a,8a-cis)-3a-Acetoxy-1,8-diacetyl-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-*b*]indole (26) from 24** — Ac<sub>2</sub>O (2.0 mL) was added to a solution of **24** (18.6 mg, 0.07 mmol) in pyridine (4.0 mL) and the mixture was stirred at rt for 1 h. The solvent was evaporated under reduced pressure to leave an oil, which was column-chromatographed on SiO<sub>2</sub> with CHCl<sub>3</sub>–MeOH (98:2, v/v) to give **26** (20.8 mg, 96%).

**26 from 25** — Ac<sub>2</sub>O (3.0 mL) was added to a solution of **25** (29.7 mg, 0.14 mmol) in pyridine (6.0 mL) and the mixture was stirred at rt for 16 h. The solvent was evaporated under reduced pressure to leave an oil, which was column-chromatographed on SiO<sub>2</sub> with CHCl<sub>3</sub>–MeOH (98:2, v/v) to give **26** (40.9 mg, 99%).

**26 from Nb-acetyl-1-hydroxytryptamine (27)** — NaOAc (23.9 mg, 0.29 mmol) was added to a solution of **27** (31.3 mg, 0.14 mmol) in Ac<sub>2</sub>O (2.0 mL) and the mixture was stirred at 118–122 °C for 4.5 h. After addition of H<sub>2</sub>O under ice cooling, the whole was extracted with CHCl<sub>3</sub>–MeOH (95:5, v/v). The extract was washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and evaporated under reduced pressure to leave an oil, which was column-chromatographed on SiO<sub>2</sub> with CHCl<sub>3</sub>–MeOH (98:2, v/v) to give **26** (20.5 mg, 47%). **26**: mp 190.0–191.0 °C (colorless prisms, recrystallized from CHCl<sub>3</sub>–hexane). IR (KBr): 3535, 2875, 1742, 1623, 1603, 1477, 1404, 1239, 1043, 904, 789, 769 cm<sup>-1</sup>. <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>, 60 °C) δ: 1.99, (3H, s), 2.04 (3H, br s), 2.43 (3H, s), 2.45–2.58 (1H, m), 2.64 (1H, br dd, *J*=11.5, 4.4 Hz), 2.82 (1H, m), 3.84 (1H, m), 6.34 (1H, br s), 7.16 (1H, t, *J*=7.6 Hz), 7.35 (1H, t, *J*=7.6 Hz), 7.52 (1H, d, *J*=7.6 Hz), 7.86 (1H, br s). MS *m/z*: 302 (M<sup>+</sup>). *Anal.* Calcd for C<sub>16</sub>H<sub>18</sub>N<sub>2</sub>O<sub>4</sub>: C, 63.56; H, 6.00; N, 9.27. Found: C, 63.42; H,



6.00; N, 9.17.

**(3a,8a-cis)-1-Acetyl-1,2,3,3a,8,8a-hexahydro-3a-hydroxypyrrolo[2,3-*b*]indole (29) from 8b-(2-acetylaminoethyl)-2,2-dimethyl-4*H*-1,3-dioxolo[4,5-*b*]indole (28)** — K<sub>2</sub>CO<sub>3</sub> (16.6 mg, 0.12 mmol) was added to a solution of **28** (6.5 mg, 0.02 mmol) in MeOH (1.0 mL) and the mixture was stirred at rt for 45 min. After addition of H<sub>2</sub>O, the whole was extracted with CHCl<sub>3</sub>–MeOH (95:5, v/v). The extract was washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and evaporated under reduced pressure to leave an oil, which was column-chromatographed on SiO<sub>2</sub> with CHCl<sub>3</sub>–MeOH (97:3, v/v) to give **29** (4.1 mg, 80%). **29**: colorless oil. IR (film): 3320, 1613, 1470, 1449, 1423, 1060, 752 cm<sup>-1</sup>. <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 2.03 (3H, s), 2.45 (1H, s, disappeared on addition of D<sub>2</sub>O), 2.41–2.57 (2H, m), 3.25–3.33 (1H, m), 3.67–3.75 (1H, m), 5.27 (1H, br s, disappeared on addition of D<sub>2</sub>O), 5.33 (1H, s), 6.63 (1H, d, *J*=7.6, 1.0 Hz), 6.81 (1H, td, *J*=7.6, 1.0 Hz), 7.19 (1H, td, *J*=7.6, 1.2 Hz), 7.31 (1H, d, *J*=7.6, 1.2 Hz). High-resolution MS *m/z*: Calcd for C<sub>12</sub>H<sub>14</sub>N<sub>2</sub>O<sub>2</sub>: 218.1056. Found: 218.1056.

**26 from 29** — Ac<sub>2</sub>O (0.5 mL) was added to a solution of **29** (6.5 mg, 0.03 mmol) in pyridine (1.0 mL) at 0 °C and the mixture was stirred at rt for 8 h. The solvent was evaporated under reduced pressure to leave an oil, which was column-chromatographed on SiO<sub>2</sub> with EtOAc to give **26** (7.2 mg, 80%).

**X-Ray Crystallographic Analysis of 4** — A single crystal (0.20x0.20x0.20 mm) of **4** was obtained by recrystallization from CHCl<sub>3</sub>. All measurements was made on a Rigaku AFC5R diffractometer with graphite monochromated Cu-*K*α radiation ( $\lambda=1.54178$  Å). Crystal data: C<sub>20</sub>H<sub>16</sub>F<sub>3</sub>N<sub>3</sub>O, *M*=454.52, monoclinic, space group *P*2<sub>1</sub>/*n* (#14), *a*=8.8339 (5) Å, *b*=12.1938 (8) Å, *c*=15.7993 (9) Å,  $\beta=93.072$  (5)°, *V*=1699.4 (2) Å<sup>3</sup>, *Z*=4, *D*<sub>calc</sub>=1.451 g/cm<sup>3</sup>, *F*(000)=768, and  $\mu(\text{CuK}\alpha)=9.40$  cm<sup>-1</sup>. The structure was solved by direct methods using MITHRIL.<sup>14</sup> The non-hydrogen atoms were refined anisotropically. The final cycle of full-matrix least-squares refinement was based on 1866 observed reflections (*I*>3.00σ(*I*), 2θ<120.2°) and 308 variable parameters. The final refinement converged with *R*=0.046 and *R*<sub>w</sub>=0.056.

Numbering  
of Atoms





Figure 1



ORTEP Drawing  
of **4** (*R* = 0.046)

**Table 2. Positional Parameters and B (eq) for 4**

atom	x	y	z	B (eq)	atom	x	y	z	B (eq)
F (1)	0.8894 (4)	0.3539 (2)	-0.0913 (1)	9.5 (2)	C (16)	1.2211 (5)	-0.2951 (4)	0.3706 (3)	5.2 (2)
F (2)	0.9224 (3)	0.4006 (2)	0.0360 (1)	6.9 (1)	C (17)	1.1484 (4)	-0.2638 (3)	0.2948 (2)	4.2 (2)
F (3)	0.7035 (3)	0.3547 (2)	-0.0115 (2)	8.4 (2)	C (18)	1.0776 (3)	-0.1627 (3)	0.2925 (2)	3.2 (1)
O (1)	0.9367 (3)	0.1526 (2)	-0.0456 (1)	4.5 (1)	C (19)	0.8815 (3)	0.2115 (3)	0.0073 (2)	3.5 (1)
N (1)	0.8435 (3)	0.1775 (2)	0.0829 (1)	3.2 (1)	C (20)	0.8490 (5)	0.3308 (3)	-0.0157 (2)	4.9 (2)
N (2)	0.7651 (3)	-0.0136 (2)	0.0658 (2)	3.7 (1)	H (1)	0.659 (4)	0.235 (3)	0.141 (2)	4.85 (2)
N (3)	0.9933 (3)	-0.1117 (2)	0.2279 (2)	3.3 (1)	H (2)	0.802 (4)	0.313 (3)	0.143 (2)	4.82 (2)
C (1)	0.7749 (5)	0.2386 (3)	0.1515 (2)	4.0 (2)	H (3)	0.775 (4)	0.191 (3)	0.278 (2)	4.65 (2)
C (2)	0.8370 (4)	0.1792 (3)	0.2289 (2)	3.8 (2)	H (4)	0.943 (4)	0.202 (3)	0.246 (2)	4.46 (2)
C (3)	0.8378 (3)	0.0585 (2)	0.2022 (2)	3.0 (1)	H (5)	0.611 (3)	0.023 (3)	0.318 (2)	4.11 (2)
C (4)	0.6797 (3)	0.0077 (2)	0.1983 (2)	3.0 (1)	H (6)	0.377 (4)	-0.067 (3)	0.288 (2)	5.31 (2)
C (5)	0.5790 (4)	-0.0033 (3)	0.2614 (2)	3.8 (2)	H (7)	0.319 (4)	-0.138 (3)	0.150 (2)	6.04 (2)
C (6)	0.4432 (4)	-0.0568 (3)	0.2429 (3)	4.6 (2)	H (8)	0.489 (4)	-0.117 (3)	0.044 (2)	4.73 (2)
C (7)	0.4104 (4)	-0.0976 (3)	0.1633 (3)	5.1 (2)	H (9)	0.755 (4)	-0.019 (3)	0.021 (2)	4.46 (2)
C (8)	0.5101 (4)	-0.0868 (3)	0.0987 (3)	4.5 (2)	H (10)	0.972 (3)	0.038 (2)	0.097 (2)	3.09 (2)
C (9)	0.6449 (3)	-0.0331 (2)	0.1181 (2)	3.3 (1)	H (11)	0.985 (3)	-0.136 (2)	0.175 (2)	3.58 (2)
C (10)	0.8709 (3)	0.0615 (2)	0.1066 (2)	3.0 (1)	H (12)	0.976 (3)	0.066 (3)	0.373 (2)	4.43 (2)
C (11)	0.9450 (3)	-0.0108 (3)	0.2557 (2)	3.2 (1)	H (13)	1.156 (4)	-0.078 (3)	0.489 (2)	5.69 (2)
C (12)	0.9957 (4)	0.0028 (3)	0.3377 (2)	3.8 (2)	H (14)	1.271 (4)	-0.249 (3)	0.494 (2)	6.35 (3)
C (13)	1.0796 (3)	-0.0923 (3)	0.3630 (2)	3.6 (1)	H (15)	1.266 (4)	-0.362 (3)	0.373 (2)	5.70 (3)
C (14)	1.1544 (4)	-0.1271 (3)	0.4388 (2)	4.7 (2)	H (16)	1.143 (3)	-0.310 (3)	0.246 (2)	4.57 (2)
C (15)	1.2235 (5)	-0.2281 (4)	0.4415 (3)	5.4 (2)					

## REFERENCES AND NOTES

1. a) This is Part 143 of a series entitled "The Chemistry of Indoles"; b) Part 142: T. Iwaki, Y. Fukui, M. Okigawa, F. Yamada, Y. Nagahama, S. Ogasawara, S. Tanaka, S. Funaki, and M. Somei, *Heterocycles*, 2016, **93**, in press. COM-15-S(T)40 .
2. Professor Emeritus of Kanazawa University. Present address: 56-7 Matsuhidai, Matsudo, Chiba 270-2214, Japan.
3. a) T. Kawasaki, M. Tabata, K. Nakagawa, K. Kobayashi, A. Kodama, T. Kobayashi, M. Hasegawa, K. Tanii, and M. Somei, *Heterocycles*, 2015, **90**, 1038, and references cited therein; b) Recent application of 1-hydroxyindole to the synthesis of benzofuroindolines: T. Tomakinian, C. Kouklovsky, and G. Vincent, *Synlett.*, 2015, 1269.
4. a) F. Yamada, D. Shinmyo, M. Nakajou, and M. Somei, *Heterocycles*, 2012, **86**, 435, and references cited therein. b) K. Yoshino, F. Yamada, and M. Somei, *Heterocycles*, 2008, **76**, 989, and references cited therein.
5. a) M. Somei, *Yakugaku Zasshi*, 2008, **128**, 527, and references cited therein; b) M. Somei, Topics in Heterocyclic Chemistry, Vol. **6**, ed. by S. Eguchi, Springer-Verlag, Berlin, 2006, pp. 77—111; c) M. Somei, Advances in Heterocyclic Chemistry, Vol. **82**, ed. by A. R. Katritzky, Elsevier Science (USA),

- 2002, pp. 101—155; d) M. Somei, *Heterocycles*, 1999, **50**, 1157; e) M. Somei, *J. Synth. Org. Chem. Jpn*, 1991, **49**, 205.
6. There is a possibility that the elimination of 1-hydroxy group and an attack by nucleophiles follow a concerted mechanism.
  7. F. Yamada, A. Goto, and M. Somei, *Heterocycles*, 2000, **53**, 1255.
  8. M. Somei, N. Oshikiri, M. Hasegawa, and F. Yamada, *Heterocycles*, 1999, **51**, 1237.
  9. M. Hasegawa, Y. Nagahama, K. Kobayashi, M. Hayashi, and M. Somei, *Heterocycles*, 2000, **52**, 483.
  10. H. Minato, M. Matsumoto, and T. Katayama, *J. Chem. Soc., Perkin Trans. 1*, 1973, 1819.
  11. D. Crich, E. Fredette, and W. J. Flosi, *Heterocycles*, 1998, **48**, 545.
  12. M. Somei, F. Yamada, A. Goto, M. Hayashi, and M. Hasegawa, *Heterocycles*, 2000, **53**, 2487.
  13. C. Takahashi, A. Numata, Y. Ito, E. Matsumura, H. Iwaki, and K. Kushida, *J. Chem. Soc., Perkin Trans. 1*, 1994, 1859.
  14. C. J. Gilmore, *J. Appl. Cryst.*, 1984, **17**, 42.