

## 神経移植の核医学的評価に関する基礎的研究

著者	森 厚文
著者別表示	Mori Hirofumi
雑誌名	平成9(1997)年度 科学研究費補助金 基盤研究(C) 研究成果報告書概要
巻	1996-1997
ページ	2p.
発行年	1999-03-15
URL	<a href="http://doi.org/10.24517/00056878">http://doi.org/10.24517/00056878</a>



# 1997 Fiscal Year Final Research Report Summary

---

## Fundamental Study on Assessment of Neurotransplantation Using Nuclear Medicine Imaging

Research Project

### Project/Area Number

---

08671011

### Research Category

---

Grant-in-Aid for Scientific Research (C)

### Allocation Type

---

Single-year Grants

### Section

---

一般

### Research Field

---

Radiation science

### Research Institution

---

KANAZAWA UNIVERSITY

### Principal Investigator

---

**MORI Hirofumi** Kanazawa University, Radioisotope Center, Professor, アイソトープ総合センター, 教授 (90019604)

### Co-Investigator(Kenkyū-buntansha)

---

TSJI Siro Kanazawa University, School of Medicine, Associate Professor, 医学部, 助教授 (70227388)

SHIBA Kazuhiro Kanazawa University, Radioisotope Center, Assistant, アイソトープ総合センター, 助手 (40143929)

### Project Period (FY)

---

1996 - 1997

### Keywords

---

Alzheimer's disease / Lesion of Nucleus Basalis Magnocellularis / Acetylcholine / Receptor / Neurotransplantation / Transporter

### Research Abstract

---

Alzheimer's disease is one of most troublesome problems encountered in the elderly population. One proposed hypothesis is that a deficit in cholinergic neurotransmission in Alzheimer's disease underlies this serious symptom of the disease. Cholinergic denervation rat model by producing an unilateral lesion of nucleus basalis magnocellularis (NBM) are reported to a model of cognitive deficits, one of instructive models of Alzheimer's disease (AD). The neurotransplantation is now promising as an effective strategy for functional repair in a variety of neural systems in disorders such as Parkinson's disease. In this study autotransplantation of vagal ganglion was performed to NBM lesioned rats. The effects of cholinergic grafts on cholinergic systems evaluated by autoradiographic images in duration of 1,2 and 4 weeks after surgery. Cerebral blood flow (CBF), muscarinic acetylcholine receptor (mAChR), m1 and m2 subtype of AChR-mRNA images were obtained using <sup>99m</sup>Tc-hexamethyl-propyleneamine ... More

## Research Products (4 results)

All Other

All Publications (4 results)

[Publications] Shiba K: "In vitro characterization of radioiodinated (-) m-Iodovesamicol in rat cerebral membranes" Life Sciences. 59 · 13. 1039-1045 (1996) ▼

[Publications] Kuji I: "Discrepancy between blood flow and muscarinic receptor distribution in rat brain after middle cerebral occlusion" Eur J Nuclear Medicine. 24 · 6. 665-669 (1997) ▼

[Publications] Shiba, K.: "In vitro characterization of radioiodinated (-) m-Iodovesamicol in rat cerebral membranes." Life Sciences. 59 (13). 1039-1045 (1996) ▼

[Publications] Kuji, I.: "Discrepancy between blood flow and nuscarinic receptor distribution in rat brain after middle cerebral occlusion." Eur J Nuclear Medicine. 24 (6). 665-669 (1997) ▼

URL: [https://kaken.nii.ac.jp/report/KAKENHI-PROJECT-08671011/086710111997kenkyu\\_seika\\_hokoku\\_](https://kaken.nii.ac.jp/report/KAKENHI-PROJECT-08671011/086710111997kenkyu_seika_hokoku_)

Published: 1999-03-15