

Autologous serum and serum components

著者	Higuchi Akihiro
著者別表示	樋口 明弘
journal or publication title	Investigative Ophthalmology and Visual Science
volume	59
number	14 Special Issue
page range	DES121-DES129
year	2018-11
URL	http://doi.org/10.24517/00062809

doi: 10.1167/iovs.17-23760



Autologous Serum and Serum Components

Akihiro Higuchi

Institute for Research Promotion, Oita University, Oita, Japan

Correspondence: Akihiro Higuchi, Institute for Research Promotion, Oita University, 1-1 Idaigaoka, Hasama-machi, Yufu City, Oita, 879-5593, Japan; higuchi-akihiro@oita-u.ac.jp.

Submitted: January 4, 2018
Accepted: March 26, 2018

Citation: Higuchi A. Autologous serum and serum components. *Invest Ophthalmol Vis Sci.* 2018;59:DES121-DES129. <https://doi.org/10.1167/iops.17-23760>

Dry eye syndrome is a multifactorial condition on the tear and ocular surface. Autologous serum eye drop is an effective method for treating dry eye. Autologous serum eye drops are now widely used by specialists since a first report in 1975. The results of a systematic study showed that the efficacy of autologous serum eye drops remains ambiguous because its preparation methods and clinical application have not been standardized. To elucidate the efficacy of autologous serum eye drops, well-designed, large-scale, high-quality randomized controlled trials need to be conducted with standardized treatment and use. Since serum components are partially similar to tear components, autologous serum eye drops improve dry eye by supplying tear components such as growth factors, proteins, and vitamins. Adding to the evidence based on the treatment of dry eye, we have found a new treatment candidate from serum: selenoprotein P (SeP). The efficacy of SeP as a treatment for dry eye was revealed by applying SeP eye drops to a dry eye rat model. Compared with phosphate-buffered saline treatment, SeP eye drops significantly reduced the fluorescein score of the cornea and suppressed the oxidative stress in the cornea, which is related to onset of dry eye, leading to improved corneal disorder. We have developed a new dry eye model caused by oxidative stress that will be used to screen candidate molecules for antioxidative activity.

Keywords: autologous serum, serum components, selenium, oxidative stress

Dry eye syndrome is a multifactorial disease of the tears and ocular surface caused by endogenous and environmental (i.e., physical,¹⁻⁴ chemical,^{5,6} and biological⁷) factors. At present, several effective methods for treating dry eye are known, and autologous serum eye drops are among these. The number of articles on the use of autologous serum as a treatment for ocular surface disease increases each year (Fig. 1). A search of PubMed in November 2017 retrieved 134 reports on “autologous serum ‘eye drops’” and 119 reports on “autologous serum ‘dry eye’.” Of these 134 reports, 44 that were published in 2015 to 2017 are summarized in Table 1.⁸⁻⁵¹ Twelve were review articles, 11 were basic research articles, and the remainder were clinical studies (Table 1). The first report on autologous serum for the treatment of ocular surface diseases was published by Ralph et al.⁵² in 1975. They continuously delivered a serum or drug solution to the ocular surface of patients with ocular surface disease using a mobile ocular perfusion pump, and serum treatment was shown to be partially effective. The efficacy of autologous serum eye drops for the treatment of dry eye was first mentioned by Fox et al.⁵³ in 1984. Many studies from various countries on the efficacy of autologous serum were subsequently reported. In Japan, studies on autologous serum eye drops were conducted by Tsubota's group⁵⁴⁻⁵⁶ and Takamura et al.,⁵⁷ who showed the usefulness of autologous serum eye drops in the treatment of dry eye caused by several diseases.⁵⁴⁻⁵⁷ At present, autologous serum eye drops are globally used as a therapeutic intervention for dry eye.

AUTOLOGOUS SERUM EYE DROPS AND OTHER THERAPEUTIC INTERVENTIONS FOR DRY EYE SYNDROME

Sy et al.³⁸ conducted an international survey to identify the most common treatments for dry eye used by specialists. The

most commonly prescribed topical treatments for aqueous-deficient dry eye included 0.05% cyclosporine A (68%), 0.1% fluorometholone (60%), 0.5% loteprednol etabonate (51%), and autologous serum eye drops³³ (49%), and nontopical medications included essential fatty acid supplements (69%), low-dose doxycycline (oral; 61%), and flaxseed supplements (33%), as well as punctal plugs³³ (75%). The top three signs and symptoms that were indicators of treatment response were fluorescein staining of the cornea, reduced foreign body sensation, and reduced burning sensation.

In Australia, the Red Cross Blood Service manufactures and distributes autologous serum eye drops to eligible patients with dry eye, persistent corneal epithelial defects,




FIGURE 1. The number of published articles, by year, retrieved from a PubMed search conducted in November 2017 using the terms “autologous serum ‘eye drops’” (open circle) and “autologous serum ‘dry eye’” (closed circle). The vertical axis shows the number of articles.



TABLE 1. Articles on Autologous Serum Eye Drops Published in 2015 Through 2017 (October). The Articles Were Searched in PubMed Using the Keyword “Autologous Serum ‘Eye Drops”

Year	Authors	Journal	Article Type	Ref. No.
2017	Giannaccare G, et al.	<i>Transfus Apber Sci.</i> 2017;56:595-604.	Review	8
2017	Alio JL, et al.	<i>Ophthalmol Ther.</i> 2017;6:1-9.	Clinical study	9
2017	Wu MF, et al.	<i>Int J Ophthalmol.</i> 2017;10:908-913.	Basic research	10
2017	Akcem HT, et al.	<i>Clin Exp Optom.</i> 2017;101:34-37.	Clinical study	11
2017	Sanchez-Avila RM, et al.	<i>Int Ophthalmol.</i> 2018;38:1193-1204.	Clinical study	12
2017	Marks DC, et al.	<i>Vox Sang.</i> 2017;112:310-317.	Basic research	13
2017	Jian-Wei L, et al.	<i>Cutan Ocul Toxicol.</i> 2017;36:377-380.	Clinical study	14
2017	Etxebarria J, et al.	<i>Acta Ophthalmol.</i> 2017;95:e693-e705.	Basic research	15
2017	Pan Q, et al.	<i>Cochrane Database Syst Rev.</i> 2017;2:CD009327.	Review	16
2017	Stachon T, et al.	<i>Klin Monbl Augenbeilkd.</i> 2017;234:1015-1018.	Basic research	17
2017	Mahelková G, et al.	<i>Clin Exp Optom.</i> 2017;100:243-249.	Clinical study	18
2017	Piccin A, et al.	<i>Blood Transfus.</i> 2017;15:333-340.	Review	19
2017	Azari AA, et al.	<i>Cutan Ocul Toxicol.</i> 2017;36:152-156.	Clinical study	20
2017	Tahmaz V, et al.	<i>Br J Ophthalmol.</i> 2017;101:322-326.	Clinical study	21
2017	Wang WY, et al.	<i>Eye Contact Lens.</i> 2017;43:225-229.	Clinical study	22
2016	Semeraro F, et al.	<i>In Vivo.</i> 2016;30:931-938.	Clinical study	23
2016	Asena L, et al.	<i>Curr Eye Res.</i> 2017;42:506-512.	Basic research	24
2016	Gus PI, et al.	<i>Oxid Med Cell Longev.</i> 2016;2016:9780193.	Clinical study	25
2016	Anitua E, et al.	<i>Exp Eye Res.</i> 2016;151:115-121.	Basic research	26
2016	Lee YK, et al.	<i>Cont Lens Anterior Eye.</i> 2016;39:425-430.	Clinical study	27
2016	Tseng CL, et al.	<i>PLoS One.</i> 2016;11:e0153573.	Basic research	28
2016	Lee JH, et al.	<i>Korean J Ophthalmol.</i> 2016;30:101-107.	Clinical trial	29
2016	von Hofsten J, et al.	<i>Int Med Case Rep J.</i> 2016;9:47-54.	Clinical study	30
2016	Rybickova I, et al.	<i>Biomed Pap.</i> 2016;160:271-275.	Clinical study	31
2016	van der Meer PF, et al.	<i>Transfus Apber Sci.</i> 2016;54:164-167.	Review	32
2016	Marshall LL, et al.	<i>Consult Pharm.</i> 2016;31:96-106.	Review	33
2016	López-García JS, et al.	<i>Cornea.</i> 2016;35:336-341.	Clinical study	34
2016	Hondur AM, et al.	<i>Curr Eye Res.</i> 2016;41:15-19.	Clinical study	35
2015	Katsakoulas I, et al.	<i>Int J Pharm Compd.</i> 2015;19:252-260.	Basic research	36
2015	Mondy P, et al.	<i>Transfus Apber Sci.</i> 2015;53:404-411.	Review	37
2015	Sy A, et al.	<i>BMC Ophthalmol.</i> 2015;15:133.	Basic research	38
2015	Mahelková G, et al.	<i>Cesk Slov Oftalmol.</i> 2015;71:184-188.	Clinical study	39
2015	Liu Y, et al.	<i>Cornea.</i> 2015;34:1214-1220.	Clinical study	40
2015	Stenwall PA, et al.	<i>Acta Ophthalmol.</i> 2015;93:654-657.	Clinical study	41
2015	Li J, et al.	<i>Cornea.</i> 2015;34:1072-1078.	Clinical trial	42
2015	van der Meer PF, et al.	<i>Transfus Apber Sci.</i> 2015;53:99-100.	Review	43
2015	Marks DC, et al.	<i>Transfus Apber Sci.</i> 2015;53:92-94.	Review	44
2015	Lin SJ, et al.	<i>Taiwan J Ophthalmol.</i> 2015;5:109-113.	Clinical study	45
2015	Wirotko B, et al.	<i>Ocul Surf.</i> 2015;13:204-212.	Review	46
2015	Espinosa A, et al.	<i>Transfus Apber Sci.</i> 2015;53:88-91.	Review	47
2015	Anitua E, et al.	<i>Acta Ophthalmol.</i> 2015;93:e605-e614.	Review	48
2015	Anitua E, et al.	<i>Exp Eye Res.</i> 2015;135:118-126.	Basic research	49
2015	Azari AA, et al.	<i>Eye Contact Lens.</i> 2015;41:133-140.	Review	50
2015	Blasetti F, et al.	<i>J Infect Dev Ctries.</i> 2015;9:55-59.	Basic research	51

and nonhealing corneal ulcers.⁴⁴ Since 2005, the Blood Service has collected whole blood from patients for the manufacturing of serum eye drops using a national, standardized process to produce 20% of serum drops. Demand for serum eye drops in Australia is increasing every year, with a 30% increase in 12 months during 2014 to 2015.⁴⁴ Based on the results of these studies, it can be concluded that autologous serum eye drops are a well-established treatment for dry eye disease.

CLINICAL STUDIES OF AUTOLOGOUS SERUM EYE DROPS

A small cohort study on patients with mixed ocular surface disease treated with autologous serum eye drops was

conducted at a Swedish tertiary referral center.³⁰ All cases ($n = 32$ eyes, 24 patients) were treated with autologous serum eye drops. The results of the study showed that patients with a persistent epithelial defect or superficial punctate keratitis responded well to treatment with autologous serum eye drops. However, large-scale randomized clinical trials are needed to fully comprehend the role of the eye drops in the treatment of ocular surface disease.

Pan et al.^{16,58} reported two systematic reviews in Cochrane Database of Systematic Reviews. In a more recent report,¹⁶ they conducted a review to evaluate the efficacy and safety of autologous serum eye drops for adults with dry eye administered alone or in combination with artificial tears compared with artificial tears alone, saline, placebo, or no treatment. The results showed that in 20% of patients, autologous serum eye drops provided some benefits in improving patient-reported

TABLE 2. Comparison of Tear and Serum Components

Components	Concentration	
	Tear, Basal	Serum†
Proteins		
Total protein	7.51 mg/mL	66–81 mg/mL
Lysozyme	2.36 mg/mL	5.0–10.2 µg/mL
Lactoferrin	1.84 mg/mL	0.17–0.28 mg/mL
Albumin	1.3 mg/mL	41–51 mg/mL
IgA	0.30 mg/mL	0.93–3.93 mg/mL
IgD	ND	0.03 mg/mL
IgE	0.1 µg/mL	0.4 µg/mL
IgG	0.126 mg/L	8.61–17.47 mg/mL
IgM	0.86 µg/mL	0.33–1.83 mg/mL
CuZn-SOD	103 ng/mg protein	ND‡
Growth factors		
EGF	1.66 ng/mL	0.72 ng/mL
TGF-α, male	247 pg/mL	147 pg/mL
TGF-α, female	180 pg/mL	147 pg/mL
TGF-β1	Not detected	140.3 ng/mL
TGF-β1*	2.32 ng/mL	-
TGF-β2	55 pg/mL	-
Vitamins		
Vitamin A	16 ng/mL	200–500 ng/mL
Vitamin C	117 µg/mL	5–9 µg/mL
Antioxidants		
Tyrosine	45 µM	77 µM
Glutathione	107 µM	ND‡
Carbohydrate		
Glucose	26 mg/L	0.6–1.2 g/L
Electrolytes		
Na ⁺	148.5 mM	138–145 mM
K ⁺	18.7 mM	3.6–4.8 mM
Ca ⁺⁺	1.73 mM	8.8–10.1 mM
Cl ⁻	112 mM	101–108 mM
HCO ₃ ⁻	26 mM	21–29 mM
NO ₃ ⁻	0.14 mM	0.19 mM
PO ₄ ³⁻	0.22 mM	1.42 mM
SO ₄ ²⁻	0.39 mM	0.53 mM

* Acid-activated tear.

† Each value is of normal concentration in the serum.

‡ These components are present in red blood cells at high concentration. See Refs. 46 through 51.

symptoms in the short term (2 weeks), but longer periods of follow-up indicated no evidence of long-term improvement. Objective clinical measures of the ocular surface showed no clear effect. The authors concluded that well-designed, large-scale, high-quality randomized controlled trials are needed to clarify the efficacy of autologous serum eye drops for the treatment of dry eye.

Despite this, autologous serum eye drops have been found to have some value in the treatment of dry eye. To elucidate the efficacy of the eye drops, the methods for its preparation, manufacturing, and storage (such as clotting time, centrifugation condition, dilution ratio, and the most suitable diluent) must be well established, and its clinical application for dry eye patients should be optimized.

COMPONENTS OF SERUM AND TEARS

Autologous serum eye drops are thought to alleviate dry eye by supplying tear components, such as various growth factors, vitamins, and proteins, that aid in the maintenance of the

Method of preparation for autologous serum eye drops




FIGURE 2. Method of preparation for autologous serum eye drops. Autologous serum eye drops were prepared using autoserum collected from each patient in sterile conditions.

ocular surface. Table 2 contains a comparison of tear^{59–62} and serum components.^{59,63,64} The concentration of total tear protein is 7.51 mg/mL, which is approximately 10% lower than that of serum. Lysozyme, lactoferrin, and albumin are the major proteins in tears, whereas albumin and IgG are the major proteins in serum. Since lysozyme protects against infection on the ocular surface via antimicrobial activity and the lactoferrin transports iron to the cornea, those proteins are thought to be secreted at high concentration in tears. Autologous serum eye drops are prepared using autologous serum under sterile conditions (Fig. 2). Blood collected from a patient is clotted and centrifuged to prepare the serum. The prepared serum is diluted to 20% to 100% with saline and preserved in a refrigerator or freezer in case of the need for long-term preservation. Because infection via the serum can occur, it is necessary to prepare autologous serum eye drops for every patient to prevent infection. Another problem is that serum is physically unstable and probably degraded by serum internal enzymes. Use of serum components for eye drops, such as growth factors, albumin, and lactoferrin, could minimize these problems. Studies with a focus on the efficacy of serum components for the treatment of dry eye have been increasing since about 2007.

GROWTH FACTORS

Tears contain several growth factors that are also contained in serum (Table 2). We previously reported an in vitro experiment using human conjunctival epithelial cell lines (CCLs) that demonstrated that epidermal growth factors (EGFs) and retinoic acid play key roles in the maintenance of the ocular surface.⁶⁵ In an animal dry eye model experiment, EGF eye drops improved tear breakup time (BUT) and fluorescein staining, but not inflammatory index and tear volume.⁶⁶ The authors concluded that EGF has potential as a therapeutic agent in the clinical treatment of dry eye.⁶⁶ Lou-Bonafonte et al.⁶⁷ reported a systematic review on clinical studies on the posology, efficacy, and safety of EGF eye drops in the treatment of certain human corneal disorders. EGF eye drops were found to be approximately 50 times more effective than vehicle eye drops in achieving accelerated and successful corneal wound healing. EGF eye drops (50–1000 ng, two or three times/day) could be a useful treatment for promoting postoperative refractive surgery and reversing cases of keratopathy secondary to systematic EGF receptor inhibitors, diabetic keratopathy, and other corneal and conjunctival disorders.




FIGURE 3. Lactoferrin uptake into corneal epithelial cells. (A, B) Immunohistochemical staining of lactoferrin receptor LRP1 in a rat cornea was performed using anti-LRP1 rabbit antibody (A) and normal rabbit IgG (B). Scale bar: 20 μ m. Original magnification: $\times 20$. (C) Lactoferrin uptake into corneal epithelial cells. The cells were incubated in the presence of 500 ng/mL lactoferrin for 0, 1, 2, or 4 hours. Results are expressed as mean \pm standard deviation ($n = 6$). Reprinted with permission from Higuchi A, Inoue H, Kaneko Y, Oonishi E, Tsubota K. Selenium-binding lactoferrin is taken into corneal epithelial cells by a receptor and prevents corneal damage in dry eye model animals. *Sci Rep.* 2016;6:36903. \copyright 2016 Higuchi et al. This is an open access article distributed under the terms of the Creative Commons Attribution License.

The efficacy of growth factor eye drops has been supported by clinical studies that have used eye drops prepared from umbilical cord blood serum (UCBS)⁸ or platelet-rich plasma (PLP).⁹ UCBS contains more growth factors than other blood-derived preparations, with higher levels of EGF, TGF- β , NGF, and VEGF.⁶⁸ Platelets are blood-derived components fundamental to the wound healing process that deliver many growth factors contained in their α -granules, such as platelet-derived growth factor (PDGF) and TGF- β . UCBS eye drops were more effective than autologous serum in the treatment of severe dry

eye, particularly when secondary to graft-versus-host disease⁶⁹ and Sjögren's syndrome.^{8,70} Autologous PLP eye drops were an effective treatment in improving the signs and symptoms of patients suffering from moderate-to-severe chronic dry eye disease.^{8,9,71,72}

Growth factors promote the proliferation and differentiation of ocular surface cells. Loo et al.⁷³ showed EGF-induced bcl-2 expression in serum-free mouse embryo cell lines. One of the mechanisms of the effectiveness of EGF relates to the antiapoptotic effect caused by bcl-2 induction. In the author's opinion, the efficacy of growth factors in the treatment of dry eye is at least partially responsible for maintaining cellular homeostasis.

Selenoprotein P (SeP)




FIGURE 4. Schematic of selenoprotein P (SeP). SeP is a selenium (Se)-transfer plasma glycoprotein that contains 10 selenocysteine (Sec) residues. One of the Secs is located in the N-terminal, and the other nine residues are localized in the C-terminal. Sec residues in the C-terminal function as a supplier of Se to the cells. *In a square frame:* structural formula Sec. Sec, the 21st amino acid, is a cysteine analogue with a sulfur atom replaced by a selenium atom. Twenty-five selenoprotein genes are present in the human genome, such as glutathione peroxidase (GPx) and thioredoxin reductase (TR). GPx and TR are implicated in the regulation of oxidative stress in cells. Since Sec is located in the active site of GPx, Sec and selenium are essential for GPx activity.

VITAMINS

Vitamins are likewise found in tears and serum (Table 2). Vitamin A is important in maintaining the health of the ocular surface, affecting cellular regulation and differentiation, and keratomalacia occurs when the diet is grossly deficient in vitamin A.⁷⁴ Vitamin A deficiency adversely affects epithelial cells in the eyelid, conjunctiva, and cornea. Hence, an absence of vitamin A causes loss of goblet cells and leads to increased epidermal keratinization and squamous metaplasia of the mucous membranes, which generally includes the cornea and conjunctiva.⁷⁵ In previous clinical studies, retinoic acid has been shown to be useful in the treatment of dry eye^{76,77} and Stevens-Johnson syndrome.⁷⁸ Kim et al.⁷⁹ conducted a comparison of the efficacy of vitamin A and 0.05% cyclosporine A eye drops in treating patients with dry eye and showed that both eye drops were effective. Vitamins A, C, and E have antioxidative properties. Antioxidative vitamins protect the cornea, lens, and retina against oxidative stress and maintain the functions of those tissues.⁷⁴ However, there have not been any remarkable reports on the efficacy of vitamin C eye drops in the treatment of dry eye.




FIGURE 5. Concentration of selenoprotein P (SeP) in patients' tears. SeP concentration with or without corneal disorder was measured. Scores are indices of dry eye (A, fluorescein scores of ≥ 1 ; B, rose bengal score of ≥ 3). Reprinted with permission from Higuchi A, Takahashi K, Hirashima M, Kawakita T, Tsubota K. Selenoprotein P controls oxidative stress in cornea. *PLoS One*. 2010;5:e9911. © 2010 Higuchi et al. This is an open access article distributed under the terms of the Creative Commons Attribution License.

ALBUMIN

Albumin is a principal protein that is plentiful in tears and serum (Table 2). Human serum albumin (HSA) is a monomeric single-chain protein constituted from 585 amino acids that does not have any prosthetic group. HSA is a remarkably stable protein that is widely distributed in body fluids. It is a multifunctional carrier protein, and a number of albumin-binding proteins have been identified from the endothelium in various tissues.^{80,81} Our clinical pilot study⁸² showed that 5% HSA eye drops for 4 weeks statistically significantly improved fluorescein staining, rose bengal staining, BUT, and symptomatic face score. An in vitro experiment using CCL and human corneal epithelial (CEPI) cell lines showed that HSA suppressed apoptosis, accompanied by the inhibition of caspase-3 activity.⁸³

Recently, to elucidate the efficacy of albumin eye drops in the treatment of severe ocular surface defects, a case-control study that used 5% albumin and 0.1% hyaluronan eye drops was performed.⁸⁴ Although healing times for both treatments were similar, albumin eye drops seemed to reduce the recurrence of sterile corneal ulcers. The authors concluded that the efficacy of both treatments was comparable, but that additional large-scale studies were necessary.

LACTOFERRIN

Lactoferrin is an iron-binding glycoprotein that is found in most exocrine fluids, such as saliva, bile, pancreatic fluid, amniotic fluid, and tears.⁸⁵ The most common metal ion associated with lactoferrin in vivo is iron in its ferric (III) form. Lactoferrin can also bind to other metal ions, such as copper and magnesium. Tear fluid contains many kinds of antioxidative stress compounds, and lactoferrin is one of these, which protects the corneal epithelium against UV irradiation.⁸⁶ Previous studies have demonstrated that the concentration of lacrimal lactoferrin reduced⁸⁷ and oral administration of lactoferrin improved the symptoms of dry eye patients with Sjögren's syndrome.⁸⁸ Using a rabbit dry eye model, Fujiwara et al.⁸⁹ showed that lactoferrin eye drops rescued corneal damage and that this effect was more remarkable using 1% apolactoferrin not saturated with iron, compared with iron-saturated lactoferrin.⁸⁹ However, in our recent study that used a dry eye model rat caused by the removal of the lacrimal glands, 1% apolactoferrin eye drops were found to be only slightly effective.⁹⁰ Apolactoferrin eye drops at a high dose (5%) showed a weak, and not statistically significant, improvement in corneal damage. Since lactoferrin is a physiological iron carrier to the corneal epithelium, it is hypothesized that a




FIGURE 6. Evaluation of selenoprotein P (SeP) eye drops using dry eye model rats. (A) Fluorescein score of a dry eye model rat treated with PBS or SeP eye drops. Results are expressed as mean \pm standard deviation ($n = 10$). Dunnett's test was used to determine the significance of differences. * and *** Indicate significant differences from the PBS treatment result ($P < 0.05$ and < 0.005 , respectively). (B) Ratio of 8-OHdG content in the corneas of dry eye model rats between PBS and SeP eye drops. Results are expressed as mean \pm standard deviation ($n = 10$). *t*-test was used to determine the significance of the differences. ** Indicates a significant difference from the PBS treatment result, $P < 0.05$. Reprinted with permission from Higuchi A, Takahashi K, Hirashima M, Kawakita T, Tsubota K. Selenoprotein P controls oxidative stress in cornea. *PLoS One*. 2010;5:e9911. © 2010 Higuchi et al. This is an open access article distributed under the terms of the Creative Commons Attribution License.




FIGURE 7. Tobacco smoke exposure rat model, prepared as described in our previous studies.^{55,64} The rat was placed in an experimental smoking chamber (60 × 40 × 35 cm) (A) with continuous fresh air ventilation for 3 hours per day for 5 days. Three hundred milliliters of mainstream cigarette smoke prepared from cigarettes containing 14 mg tar (Seven Stars; Japan Tobacco, Tokyo, Japan) was injected into the smoking chamber (B, C) every 30 minutes for a total of six times over a 3-hour period per day for 5 days during the exposure period (D).

physiological mechanism, for example, a ligand-uptake receptor system, is essential for iron uptake to the corneal epithelium. Recently, we reported that a lactoferrin uptake receptor, LRP-1, was expressed in the corneal epithelium (Figs. 3A, 3B) and that lactoferrin was taken up into the CEPI cells (Fig. 3C). Lactoferrin was promptly taken up into the CEPI cells via the lactoferrin receptor.⁹¹

SELENOPROTEIN P

Although the above-mentioned serum components have been found to be useful in the treatment of dry eye, there are no candidates with excellent efficacy. We tried to find a new candidate from serum for the treatment of dry eye.⁹² Human serum was fractionated using column chromatography, and the index of purification of the candidate was cellular viability. The details can be found in Higuchi et al.⁹² The fraction finally obtained was separated using SDS-polyacrylamide gel electrophoresis (SDS-PAGE) and the candidate band of SDS-PAGE was cut out for identification using the Edman method. We finally identified selenoprotein P (SeP) as a new candidate for the treatment of dry eye. Figure 4 shows a schematic of SeP.

SeP is a selenium-transfer plasma glycoprotein⁹³ that is present in extracellular fluids such as plasma⁹⁴ and milk.⁹⁵ The principal source of SeP is thought to be the liver.⁹⁶ SeP is a bifunctional protein,⁹⁷ as the Sec residue in the N-terminal shows GPx activity⁹⁸ and Sec residues in the C-terminal function as a cellular selenium supplier.⁹⁹ Selenium is an essential trace element for the human body and performs physiological functions as a selenocysteine (Fig. 4), which regulate oxidative stress in the cells. We have reported that SeP is synthesized in the lacrimal gland and secreted in tears to supply selenium to the cornea.⁹² Human tears (2 μL) were obtained by capillary collection from patients ($n = 49$) with or without dry eye symptoms. The concentration of SeP in patients without corneal disorder was significantly higher than

that in patients with corneal disorder ($P = 0.01$), as evaluated using fluorescein scoring (Fig. 5A), and a similar result was obtained using rose bengal scoring (Fig. 5B). The efficacy of SeP in the treatment of dry eye was evaluated by applying SeP eye drops to a dry eye rat model. The fluorescein scores of corneas treated with SeP eye drops were significantly lower than in those treated with PBS (Fig. 6A). SeP eye drops suppressed the production of the oxidative stress marker 8-OHdG in the cornea (Fig. 6B). SeP eye drops improved corneal disorder associated with the suppression of oxidative stress in the cornea of a dry eye model rat.

OXIDATIVE STRESS AND DRY EYE

Ocular surface is strongly affected by oxidative stress caused by several factors, such as light exposure (including ultraviolet irradiation), molecular oxygen, chemical compounds, and direct contact with airflow. Oxidative stress causes and/or exacerbates many ocular diseases, including dry eye. Serum and tears contain antioxidative components: SeP, lactoferrin, glutathione, and vitamins (Table 2). To evaluate the efficacy of antioxidative components in the treatment of dry eye, we attempted to prepare a new dry eye model principally caused by oxidative stress. Since smoking elevates oxidative stress in the whole body, we placed rats in the experimental smoking chamber and systemically exposed them to the main tobacco smoke, that is, a rat tobacco smoke exposure model¹⁰⁰ (Fig. 7). We plan to use the tobacco smoke exposure dry eye model to screen candidate molecules for antioxidative effects.

CONCLUSIONS

At the present time, autologous serum eye drops are a widely used treatment for dry eye syndrome both in Japan and internationally. However, there are several problems relating to the methods used for the preparation and preservation of

autologous serum that need to be addressed. Furthermore, research on autologous serum eye drops will lead to the development of new drugs for the treatment of dry eye based on compounds derived from serum components.

Acknowledgments

The author thanks Kazuo Tsubota of the Department of Ophthalmology, Keio University, School of Medicine; Masaki Hirashima of the Chemo-Sero-Therapeutic Research Institute; and Yoshio Kaneko of Tokyo New Drug Research Laboratories, Kowa Co., Ltd. for their valuable suggestions.

Supported in part by a grant from Adaptable and Seamless Technology Transfer Program through target-driven R&D, Japan Science and Technology Agency, and Smoking Research Foundation. Funding of the publication fee and administration was provided by the Dry Eye Society, Tokyo, Japan. The Dry Eye Society had no role in the contents or writing of the manuscript.

Disclosure: **A. Higuchi**, None

References

- Craig JP, Singh I, Tomlinson A, Morgan PB, Efron N. The role of tear physiology in ocular surface temperature. *Eye*. 2000; 14:635-641.
- Cekic O, Ohji M, Hayashi A, Fang XY, Kusaka S, Tano Y. Effects of humidified and dry air on corneal endothelial cells during vitreal fluid-air exchange. *Am J Ophthalmol*. 2002; 134:75-80.
- Zuclich JA, Connolly JS. Ocular damage induced by near-ultraviolet laser radiation. *Invest Ophthalmol Vis Sci*. 1976; 15:760-764.
- Nakamura S, Shibuya M, Nakashima H, et al. Involvement of oxidative stress on corneal epithelial alterations in a blink-suppressed dry eye. *Invest Ophthalmol Vis Sci*. 2007;48: 1552-1558.
- Higuchi A, Ito K, Dogru M, et al. Corneal damage and lacrimal glands dysfunction in a smoking rat model. *Free Radic Biol Med*. 2011;51:2210-2216.
- Rios JL, Boechat JL, Gioda A, dos Santos CY, de Aquino Neto FR, Lapa e Silva JR. Symptoms prevalence among office workers of a sealed versus a non-sealed building: associations to indoor air quality. *Environ Int*. 2009;35:1136-1141.
- Ayaki M, Kawashima M, Uchino M, Tsubota K, Negishi K. Possible association between subtypes of dry eye disease and seasonal variation. *Clin Ophthalmol*. 2017;11:1769-1775.
- Giannaccare G, Versura P, Buzzi M, Primavera L, Pellegrini M, Campos EC. Blood derived eye drops for the treatment of cornea and ocular surface diseases. *Transfus Apher Sci*. 2017;56:595-604.
- Alio JL, Rodriguez AE, Ferreira-Oliveira R, Wróbel-Dudzińska D, Abdelghany AA. Treatment of dry eye disease with autologous platelet-rich plasma: a prospective, interventional, non-randomized study. *Ophthalmol Ther*. 2017;6:1-9.
- Wu MF, Stachon T, Seitz B, Langenbucher A, Szentmáry N. Effect of human autologous serum and fetal bovine serum on human corneal epithelial cell viability, migration and proliferation in vitro. *Int J Ophthalmol*. 2017;10:908-913.
- Akcam HT, Unlu M, Karaca EE, Yazici H, Aydin B, Hondur AM. Autologous serum eye-drops and enhanced epithelial healing time after photorefractive keratectomy. *Clin Exp Optom*. 2017;101:34-37.
- Sanchez-Avila RM, Merayo-Llives J, Riestra AC, et al. Treatment of patients with neurotrophic keratitis stages 2 and 3 with plasma rich in growth factors (PRGF-Endoret) eye-drops. *Int Ophthalmol*. 2018;38:1193-1204.
- Marks DC, van der Meer PF; Biomedical Excellence for Safer Transfusion (BEST) Collaborative. Serum eye drops: a survey of international production methods. *Vox Sang*. 2017;112: 310-317.
- Jian-Wei L, Xiu-Yun L, Ai-Jun D. Effectiveness of heparin eye drops in paraquat-induced ocular injury. *Cutan Ocul Toxicol*. 2017;36:377-380.
- Etxebarria J, Sanz-Lázaro S, Hernández-Moya R, et al. Serum from plasma rich in growth factors regenerates rabbit corneas by promoting cell proliferation, migration, differentiation, adhesion and limbal stemness. *Acta Ophthalmol*. 2017;95:e693-e705.
- Pan Q, Angelina A, Marrone M, Stark WJ, Akpek EK. Autologous serum eye drops for dry eye. *Cochrane Database Syst Rev*. 2017;2:CD009327.
- Stachon T, Wu MF, Bischoff M, et al. Amniotic membrane suspension and autologous serum - are they important for wound healing? [in German]. *Klin Monbl Augenheilkd*. 2017;234:1015-1018.
- Mahelková G, Jirsova K, Seidler Stangova P, et al. Using corneal confocal microscopy to track changes in the corneal layers of dry eye patients after autologous serum treatment. *Clin Exp Optom*. 2017;100:243-249.
- Piccin A, Di Pierro AM, Canzian L, et al. Platelet gel: a new therapeutic tool with great potential. *Blood Transfus*. 2017; 15:333-340.
- Azari AA, Karadag R, Kanavi MR, et al. Safety and efficacy of autologous serum eye drop for treatment of dry eyes in graft-versus-host disease. *Cutan Ocul Toxicol*. 2017;36:152-156.
- Tahmaz V, Gehlsen U, Sauerbier L, et al. Treatment of severe chronic ocular graft-versus-host disease using 100% autologous serum eye drops from a sealed manufacturing system: a retrospective cohort study. *Br J Ophthalmol*. 2017;101:322-326.
- Wang WY, Lee YK, Tsai SH, Lin YC, Chen YM. Autologous serum eye drops combined with silicone hydrogen lenses for the treatment of postinfectious corneal persistent epithelial defects. *Eye Contact Lens*. 2017;43:225-229.
- Semeraro F, Forbice E, Nascimbeni G, et al. Effect of autologous serum eye drops in patients with sjögren syndrome-related dry eye: clinical and in vivo confocal microscopy evaluation of the ocular surface. *In Vivo*. 2016; 30:931-938.
- Asena L, Suveren EH, Karabay G, Dursun Altinors D. Human breast milk drops promote corneal epithelial wound healing. *Curr Eye Res*. 2017;42:506-512.
- Gus PI, Marinho D, Zelanis S, et al. A case-control study on the oxidative balance of 50% autologous serum eye drops. *Oxid Med Cell Longev*. 2016;2016:9780193.
- Anitua E, Muruzabal F, de la Fuente M, Riestra A, Merayo-Llives J, Orive G. PRGF exerts more potent proliferative and anti-inflammatory effects than autologous serum on a cell culture inflammatory model. *Exp Eye Res*. 2016;151:115-121.
- Lee YK, Lin YC, Tsai SH, Chen WL, Chen YM. Therapeutic outcomes of combined topical autologous serum eye drops with silicone-hydrogel soft contact lenses in the treatment of corneal persistent epithelial defects: a preliminary study. *Cont Lens Anterior Eye*. 2016;39:425-430.
- Tseng CL, Chen ZY, Renn TY, Hsiao SH, Burnouf T. Solvent/detergent virally inactivated serum eye drops restore healthy ocular epithelium in a rabbit model of dry-eye syndrome. *PLoS One*. 2016;11:e0153573.
- Lee JH, Kim MJ, Ha SW, Kim HK. Autologous platelet-rich plasma eye drops in the treatment of recurrent corneal erosions. *Korean J Ophthalmol*. 2016;30:101-107.
- von Hofsten J, Egardt M, Zetterberg M. The use of autologous serum for the treatment of ocular surface disease at a Swedish tertiary referral center. *Int Med Case Rep J*. 2016;9: 47-54.

31. Rybickova I, Vesela V, Fales I, Skalicka P, Jirsova K. Apoptosis of conjunctival epithelial cells before and after the application of autologous serum eye drops in severe dry eye disease. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub.* 2016;160:271-275.
32. van der Meer PF, Seghatchian J, Marks DC. Quality standards, safety and efficacy of blood-derived serum eye drops: a review. *Transfus Apher Sci.* 2016;54:164-167.
33. Marshall LL, Roach JM. Treatment of dry eye disease. *Consult Pharm.* 2016;31:96-106.
34. López-García JS, García-Lozano I, Rivas L, Giménez C, Acera A, Suárez-Cortés T. Effects of autologous serum eye drops on conjunctival expression of MUC5AC in patients with ocular surface disorders. *Cornea.* 2016;35:336-341.
35. Hondur AM, Akcam HT, Karaca EE, Yazici Eroglu H, Aydin B. Autologous serum eye drops accelerate epithelial healing after LASEK. *Curr Eye Res.* 2016;41:15-19.
36. Katsakoulas I, Lougovi C, Paraskevopoulou P, Vougioukas N. Protocol of blood serum eye drops. *Int J Pharm Compd.* 2015;19:252-260.
37. Mondy P, Brama T, Fisher J, et al. Sustained benefits of autologous serum eye drops on self-reported ocular symptoms and vision-related quality of life in Australian patients with dry eye and corneal epithelial defects. *Transfus Apher Sci.* 2015;53:404-411.
38. Sy A, O'Brien KS, Liu MP, et al. Expert opinion in the management of aqueous deficient dry eye disease (DED). *BMC Ophthalmol.* 2015;15:133.
39. Mahelková G, Veselá V, Seidler Štangová P, et al. Tear osmolarity in patients with severe dry eye syndrome before and after autologous serum treatment: a comparison with tear osmolarity in healthy volunteers [in Czech]. *Cesk Slov Oftalmol.* 2015;71:184-188.
40. Liu Y, Hirayama M, Cui X, Connell S, Kawakita T, Tsubota K. Effectiveness of autologous serum eye drops combined with punctal plugs for the treatment of sjögren syndrome-related dry eye. *Cornea.* 2015;34:1214-1220.
41. Stenwall PA, Bergström M, Seiron P, et al. Improving the anti-inflammatory effect of serum eye drops using allogeneic serum permissive for regulatory T cell induction. *Acta Ophthalmol.* 2015;93:654-657.
42. Li J, Zhang X, Zheng Q, et al. Comparative evaluation of silicone hydrogel contact lenses and autologous serum for management of Sjögren syndrome-associated dry eye. *Cornea.* 2015;34:1072-1078.
43. van der Meer PF, Seghatchian J, de Korte D. Autologous and allogeneic serum eye drops. The Dutch perspective. *Transfus Apher Sci.* 2015;53:99-100.
44. Marks DC, Fisher J, Mondy P, Seghatchian J, Dennington PM. Serum eye drop preparation in Australia: current manufacturing practice. *Transfus Apher Sci.* 2015;53:92-94.
45. Lin SJ, Su CC, Chang DCK, et al. Autologous serum therapy in recalcitrant laser-assisted in situ keratomileusis-induced neurotrophic epitheliopathy. *Taiwan J Ophthalmol.* 2015;5:109-113.
46. Wirostko B, Rafii M, Sullivan DA, Morelli J, Ding J. Novel therapy to treat corneal epithelial defects: a hypothesis with growth hormone. *Ocul Surf.* 2015;13:204-212.
47. Espinosa A, Hjorth-Hansen H, Aasly K, Teigum I, Sivertsen G, Seghatchian J. Implementation of a standardised method for the production of allogeneic serum eye drops from regular blood donors in a Norwegian University Hospital: some methodological aspects and clinical considerations. *Transfus Apher Sci.* 2015;53:88-91.
48. Anitua E, Muruzabal F, Tayebba A, et al. Autologous serum and plasma rich in growth factors in ophthalmology: preclinical and clinical studies. *Acta Ophthalmol.* 2015;93:e605-e614.
49. Anitua E, de la Fuente M, Muruzabal F, Riestra A, Merayo-Llodes J, Orive G. Plasma rich in growth factors (PRGF) eye drops stimulates scarless regeneration compared to autologous serum in the ocular surface stromal fibroblasts. *Exp Eye Res.* 2015;135:118-126.
50. Azari AA, Rapuano CJ. Autologous serum eye drops for the treatment of ocular surface disease. *Eye Contact Lens.* 2015;41:133-140.
51. Blasetti F, Usai D, Sotgia S, Carru C, Zanetti S, Pinna A. A protocol for microbiologically safe preparation, storage, and use of autologous serum eye-drops in low-income countries. *J Infect Dev Ctries.* 2015;9:55-59.
52. Ralph RA, Doane MG, Dohlman CH. Clinical experience with a mobile ocular perfusion pump. *Arch Ophthalmol.* 1975;93:1039-1043.
53. Fox RI, Chan R, Michelson JB, Belmont JB, Michelson PE. Beneficial effect of artificial tears made with autologous serum in patients with keratoconjunctivitis sicca. *Arthritis Rheum.* 1984;27:459-461.
54. Tsubota K, Goto E, Fujita H, et al. Treatment of dry eye by autologous serum application in Sjögren's syndrome. *Br J Ophthalmol.* 1999;83:390-395.
55. Ogawa Y, Okamoto S, Mori T, et al. Autologous serum eye drops for the treatment of severe dry eye in patients with chronic graft-versus-host disease. *Bone Marrow Transplant.* 2003;31:579-583.
56. Noda-Tsuruya T, Asano-Kato N, Toda I, Tsubota K. Autologous serum eye drops for dry eye after LASIK. *J Refract Surg.* 2006;22:61-66.
57. Takamura E, Shinozaki K, Hata H, Yukari J, Hori S. Efficacy of autologous serum treatment in patients with severe dry eye. *Adv Exp Med Biol.* 2002;506(pt B):1247-1250.
58. Pan Q, Angelina A, Zambrano A, et al. Autologous serum eye drops for dry eye. *Cochrane Database Syst Rev.* 2013;8:CD009327.
59. Tsubota K, Higuchi A. Serum application for the treatment of ocular surface disorders. *Int Ophthalmol Clin.* 2000;40:113-122.
60. Sariri R, Ghafoori H. Tear proteins in health, disease, and contact lens wear. *Biochemistry (Mosc).* 2008;73:381-392.
61. Bright, AM, Tighe, BJ. The composition and interfacial proteins of tears, tear substitute and tear models. *J Br Contact Lens Assoc.* 1993;16:57-66.
62. Lew H, Yun YS, Lee SY. Electrolytes and electrophoretic studies of tear proteins in tears of patients with nasolacrimal duct obstruction. *Ophthalmologica.* 2005;219:142-146.
63. Muruganand D, Daniel E, Ebenezer GJ, Rabboni SE, Segar P, Job CK. Mycobacterium leprae infection and serum lactoferrin levels. *Lepr Rev.* 2004;75:282-288.
64. Barthe C, Galabert C, Guy-Crotte O, Figarella C. Plasma and serum lactoferrin levels in cystic fibrosis. Relationship with the presence of cystic fibrosis protein. *Clin Chim Acta.* 1989;181:183-188.
65. Higuchi A, Shimmura S, Takeuchi T, Suematsu M, Tsubota K. Elucidation of apoptosis induced by serum deprivation in cultured conjunctival epithelial cells. *Br J Ophthalmol.* 2006;90:760-764.
66. Xiao X, He H, Lin Z, et al. Therapeutic effects of epidermal growth factor on benzalkonium chloride-induced dry eye in a mouse model. *Invest Ophthalmol Vis Sci.* 2012;53:191-197.
67. Lou-Bonafonte JM, Bonafonte-Marquez E, Bonafonte-Royo S, Martínez-Carpio PA. Posology, efficacy, and safety of epidermal growth factor eye drops in 305 patients: logistic regression and group-wise odds of published data. *J Ocul Pharmacol Ther.* 2012;28:467-472.

68. Versura P, Profazio V, Buzzi M, et al. Efficacy of standardized and quality-controlled cord blood serum eye drop therapy in the healing of severe corneal epithelial damage in dry eye. *Cornea*. 2013;32:412-418.
69. Yoon KC, Jeong IY, Im SK, Park YG, Kim HJ, Choi J. Therapeutic effect of umbilical cord serum eyedrops for the treatment of dry eye associated with graft-versus-host disease. *Bone Marrow Transplant*. 2007;39:231-235.
70. Yoon KC, Heo H, Im SK, You IC, Kim YH, Park YG. Comparison of autologous serum and umbilical cord serum eye drops for dry eye syndrome. *Am J Ophthalmol*. 2007;144:86-92.
71. Alio JL, Colecha JR, Pastor S, Rodriguez S, Artola A. Symptomatic dry eye treatment with autologous platelet-rich plasma. *Ophthalmic Res*. 2007;39:124-129.
72. Sanchez-Avila RM, Merayo-Llodes J, Riestra AC, et al. The effect of immunologically safe plasma rich in growth factor eye drops in patients with Sjögren syndrome. *J Ocul Pharmacol Ther*. 2017;33:391-399.
73. Loo DT, Bradford S, Helmrich A, et al. Bcl-2 inhibits cell death of serum-free mouse embryo cells caused by epidermal growth factor deprivation. *Cell Biol Toxicol*. 1998;14:375-382.
74. Brown NA, Bron AJ, Harding JJ, Dewar HM. Nutrition supplements and the eye. *Eye (Lond)*. 1998;12(pt 1):127-133.
75. Hatchell DL, Sommer A. Detection of ocular surface abnormalities in experimental vitamin A deficiency. *Arch Ophthalmol*. 1984;102:1389-1393.
76. Tseng SC, Maumenee AE, Stark WJ, et al. Topical retinoid treatment for various dry-eye disorders. *Ophthalmology*. 1985;92:717-727.
77. Selek H, Unlu N, Orhan M, et al. Evaluation of retinoic acid ophthalmic emulsion in dry eye. *Eur J Ophthalmol*. 2000;10:121-127.
78. Singer L, Brook U, Romem M, et al. Vitamin A in Stevens-Johnson syndrome. *Ann Ophthalmol*. 1989;21:209-210.
79. Kim EC, Choi JS, Joo CK. A comparison of vitamin a and cyclosporine a 0.05% eye drops for treatment of dry eye syndrome. *Am J Ophthalmol*. 2009;147:206-213.
80. Schnitzer JE, Oh P. Albondin-mediated capillary permeability to albumin. Differential role of receptors in endothelial transcytosis and endocytosis of native and modified albumins. *J Biol Chem*. 1994;269:6072-6082.
81. Brunskill NJ. Molecular interactions between albumin and proximal tubular cells. *Exp Nephrol*. 1998;6:491-495.
82. Shimmura S, Ueno R, Matsumoto Y, et al. Albumin as a tear supplement in the treatment of severe dry eye. *Br J Ophthalmol*. 2003;87:1279-1283.
83. Higuchi A, Ueno R, Shimmura S, Suematsu M, Dogru M, Tsubota K. Albumin rescues ocular epithelial cells from cell death in dry eye. *Curr Eye Res*. 2007;32:83-88.
84. Schargus M, Kohlhaas M, Unterlauff JD. Treatment of severe ocular surface disorders with albumin eye drops. *J Ocul Pharmacol Ther*. 2015;31:291-295.
85. Masson PL, Heremans JF, Dive C. An iron-binding protein common to many external secretion. *Clin Chim Acta*. 1996;14:735-739.
86. Fujihara T, Nagano T, Endo K, Nakamura M, Nakata K. Lactoferrin protects against UV-B irradiation-induced corneal epithelial damage in rats. *Cornea*. 2000;19:207-211.
87. Solomon A, Dursun D, Liu Z, Xie Y, Macri A, Pflugfelder SC. Pro- and anti-inflammatory forms of interleukin-1 in the tear fluid and conjunctiva of patients with dry-eye disease. *Invest Ophthalmol Vis Sci*. 2001;42:2283-2292.
88. Dogru M, Matsumoto Y, Yamamoto Y, et al. Lactoferrin in Sjögren's syndrome. *Ophthalmology*. 2007;114:2366-2367.
89. Fujihara T, Nagano T, Nakamura M, Shirasawa E. Lactoferrin suppresses loss of corneal epithelial integrity in a rabbit short-term dry eye model. *J Ocul Pharmacol Ther*. 1998;14:99-107.
90. Higuchi A, Inoue H, Kawakita T, Ogishima T, Tsubota K. Selenium compound protects corneal epithelium against oxidative stress. *PLoS One*. 2012;7:e45612.
91. Higuchi A, Inoue H, Kaneko Y, Oonishi E, Tsubota K. Selenium-binding lactoferrin is taken into corneal epithelial cells by a receptor and prevents corneal damage in dry eye model animals. *Sci Rep*. 2016;6:36903.
92. Higuchi A, Takahashi K, Hirashima M, Kawakita T, Tsubota K. Selenoprotein P controls oxidative stress in cornea. *PLoS One*. 2010;5:e9911.
93. Motsenbocker MA, Tappel AL. A selenocysteine-containing selenium-transport protein in rat plasma. *Biochim Biophys Acta*. 1982;719:147-153.
94. Burk RF, Hill KE. Selenoprotein P. A selenium-rich extracellular glycoprotein. *J Nutr*. 1994;124:1891-1897.
95. Schweizer U, Michaelis M, Kohrle J, Schomburg L. Efficient selenium transfer from mother to offspring in selenoprotein-P-deficient mice enables dose-dependent rescue of phenotypes associated with selenium deficiency. *Biochem J*. 2004;378:21-26.
96. Carlson BA, Novoselov SV, Kumaraswamy E, et al. Specific excision of the selenocysteine tRNA[Ser]Sec (Trsp) gene in mouse liver demonstrates an essential role of selenoproteins in liver function. *J Biol Chem*. 2004;279:8011-8017.
97. Saito Y, Sato N, Hirashima M, et al. Domain structure of bi-functional selenoprotein P. *Biochem J*. 2004;381:841-846.
98. Saito Y, Hayashi T, Tanaka A, et al. Selenoprotein P in human plasma as an extracellular phospholipid hydroperoxide glutathione peroxidase. Isolation and enzymatic characterization of human selenoprotein P. *J Biol Chem*. 1999;274:2866-2871.
99. Saito Y, Takahashi K. Characterization of selenoprotein P as a selenium supply protein. *Eur J Biochem*. 2002;269:5746-5751.
100. Higuchi A, Ito K, Dogru M, et al. Corneal damage and lacrimal gland dysfunction in a smoking rat model. *Free Radic Biol Med*. 2011;51:2210-2216.