

高温気相反応速度は超臨界水領域へ適用できるか： OHラジカルの反応を例として

著者	高橋 憲司
著者別表示	Takahashi Kenji
雑誌名	平成16(2004)年度 科学研究費補助金 基盤研究(C) 研究成果報告書概要
巻	2003 2004
ページ	2p.
発行年	2006-07-10
URL	http://doi.org/10.24517/00063169



2004 Fiscal Year Final Research Report Summary

Applicability of reaction rate in gas phase data to supercritical water region

Research Project

Project/Area Number

15560660

Research Category

Grant-in-Aid for Scientific Research (C)

Allocation Type

Single-year Grants

Section

一般

Research Field

Reaction engineering/Process system

Research Institution

Kanazawa University

Principal Investigator

TAKAHASHI Kenji Kanazawa University, Graduate School of Natural Science and Technology, Associate Professor, 自然科学研究科, 助教授 (00216714)

Project Period (FY)

2003 - 2004

Keywords

hydrated electron / Hydroxyl radical / solvation / dissociative electron attachment / electron affinity / electron transfer / supercritical water / pulse radiolysis

Research Abstract

To study reaction rate of OH radical in high temperature water, the most direct method is to observe the transient absorption of OH radical itself. At room temperature there are a several demonstrations for such direct measurements, however, there are few example under high temperature conditions. In the direct UV absorption measurements, the absorption by sapphire windows prevents the measurement of OH signal. The actual absorption intensity due to OH radical was determined by subtraction of the sapphire window signals. The absorption spectra of OH radical in high temperature water shifted to blue. The absorption coefficient of OH radical at 250 nm decreased with increasing the temperature. At room temperature, the absorption coefficient of OH radical is known as to be 500. At 300 C, we found that the absorption coefficients decreased to 200.

The temperature dependence of OH radical reaction rate was examined. The rates are followed by the Arrhenius temperature dependence up to 125 C. Above 125 C, however, the reaction rates are independent on the water temperature. The reaction rate at 125 C was $1 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$.

Research Products (4 results)

All 2005 2004

All Journal Article

- [Journal Article] Pulse Radiolysis of Supercritical Water. III. Spectrum and thermodynamics of the hydrated electron 2005 ▾
- [Journal Article] Pulse Radiolysis of Supercritical Water. III. Spectrum and thermodynamics of the hydrated electron 2005 ▾
- [Journal Article] Reaction rates of the hydrated electron with N₂O in high temperature water and potential surface of the N₂O anion 2004 ▾
- [Journal Article] Reaction rates of the hydrated electron with N₂O in high temperature water and potential surface of the N₂O anion 2004 ▾

URL: https://kaken.nii.ac.jp/report/KAKENHI-PROJECT-15560660/155606602004kenkyu_seika_hokoku_

Published: 2006-07-10