

細胞増殖因子の心筋内投与による心筋梗塞の血管新生療法の研究

著者	川筋 道雄
著者別表示	Kawasuji Michio
雑誌名	平成12(2000)年度 科学研究費補助金 基盤研究(C) 研究成果報告書概要
巻	1999 2000
ページ	2p.
発行年	2002-03-25
URL	http://doi.org/10.24517/00063967



2000 Fiscal Year Final Research Report Summary

Research of angiogenic therapy of myocardial infarction with intramyocardial administration of basic fibroblast growth factor

Research Project

Project/Area Number

11671311

Research Category

Grant-in-Aid for Scientific Research (C)

Allocation Type

Single-year Grants

Section

一般

Research Field

Thoracic surgery

Research Institution

Kanazawa University

Principal Investigator

KAWASUJI Michio Kanazawa Univ. School of Med. Associate Professor, 医学部, 助教授 (40135067)

Co-Investigator(Kenkyū-buntansha)

YADUDA Tamotsu Kanazawa Univ. School of Med. Lecturer, 医学部・附属病院, 助手 (80324119)

FUJII Susumu Kanazawa Univ. School of Med. Lecturer

Project Period (FY)

1999 - 2000

Keywords

Myocardial infarction / Basic fibroblast growth factor / Angiogenesis

Research Abstract

This study was designed to evaluate the effects of intramyocardial administration of basic fibroblast growth factor (bFGF) on myocardial blood flow, angiogenesis, and ventricular function in a canine acute myocardial infarction model. Myocardial infarction was induced in dogs by ligation of the left anterior descending coronary artery. Within 5 minutes after coronary occlusion, 100µg of bFGF in 1 mL of saline was injected into the infarct and border zone in 6 dogs, whereas saline alone was used in 6 control dogs. Myocardial blood flow was determined with colored microsphere before and immediately after coronary ligation and again 3, 7, 14, and 28 days after treatment. Angiogenesis was evaluated by immunohistochemical studies 28 days later. Treatment with bFGF significantly increased the endocardial blood flow in the

border zone as well as epicardial blood flow in the infarcted zone. Treatment with bFGF significantly increased the capillary density as well as arteriolar density in the border zone. Treatment with bFGF significantly reduced the change in ratio of thickness of the infarcted wall to the normal wall. It improved the left ventricular ejection fraction. Intramyocardial administration of bFGF increased the regional myocardial blood flow, reduced thinning of the infarcted region, and improved ventricular function. Intramyocardial administration of bFGF may be a new therapeutic approach for patients with acute myocardial infarction.

Research Products (13 results)

All Other
All Publications

[Publications] Michio Kawasuji: "Arterial revascularization - 18 year experience with coronary artery bypass grafting in patients with familial hypercholesterolemia."Jpn.J.Thorac.Cardiovasc.Surg.. 47. 330-334 (1999) ▼

[Publications] 川筋道雄: "不安定狭心症に対する外科治療"循環器科. 46. 573-578 (1999) ▼

[Publications] Yasuda Tamotsu: "Coronary artery bypass grafting in patients with hypopituitarism"Jpn.Circulation J.. 64. 207-208 (2000) ▼

[Publications] Michio Kawasuji: "Therapeutic angiogenesis with intramyocardial administration of basic fibroblast growth factor"Ann.Thorac.Surg.. 69. 1155-1161 (2000) ▼

[Publications] Michio Kawasuji: "Coronary artery bypass surgery with arterial grafts in familial hypercholesterolemia"J.Thorac.Cardiovasc.Surg.. 119. 1008-1014 (2000) ▼

[Publications] Michio Kawasuji: "Near-infrared monitoring of myocardial oxygenation during ischemic preconditioning"Ann.Thorac.Surg.. 69. 1806-1810 (2000) ▼

[Publications] 川筋道雄: "胸部外科手術のpitfallと最新術式(分担執筆)"日本胸部外科学会・学術企画印刷. 372 (1999) ▼

[Publications] Michio Kawasuji et al.: "Arterial revascularization -18-year experience with coronary artery bypass grafting in patients with familial hypercholesterolemia."Jpn.Circulation J.. 47. 330-334 (1999) ▼

[Publications] Michio Kawasuji: "Surgical treatment of unstable angina"Junkankika. 46. 573-578 (1999) ▼

[Publications] Tamotsu Yasuda et al.: "Coronary artery bypass grafting in patients with hypopituitarism."Jpn.Circulation J.. 64. 207-208 (2000) ▼

[Publications] Michio Kawasuji et al.: "Therapeutic angiogenesis with intramyocardial administration of basic fibroblast growth factor."Ann.Thorac.Surg.. 69. 1155-1161 (2000) ▼

[Publications] Michio Kawasuji et al.: "Coronary artery bypass surgery with arterial grafts in familial hypercholesterolemia."J.Thorac.Cardiovasc.Surg.. 119. 1008-1014 (2000) ▼

[Publications] Michio Kawasuji et al.: "Near-infrared monitoring of myocardial oxygenation during ischemic preconditioning."Ann.Thorac.Surg.. 69. 1806-1810 (2000) ▼

URL: https://kaken.nii.ac.jp/report/KAKENHI-PROJECT-11671311/116713112000kenkyu_seika_hokoku

Published: 2002-03-25