

# Estimates for singular integrals along surfaces of revolution

著者	Sato Shuichi
著者別表示	佐藤 秀一
journal or publication title	Journal of the Australian Mathematical Society
volume	86
number	3
page range	413-430
year	2009-06-01
URL	<a href="http://hdl.handle.net/2297/19609">http://hdl.handle.net/2297/19609</a>

doi: 10.1017/S1446788708000773

# ESTIMATES FOR SINGULAR INTEGRALS ALONG SURFACES OF REVOLUTION

SHUICHI SATO

ABSTRACT. We prove certain  $L^p$  estimates ( $1 < p < \infty$ ) for nonisotropic singular integrals along surfaces of revolution. The singular integrals are defined by rough kernels. As an application we obtain  $L^p$  boundedness of the singular integrals under a sharp size condition on their kernels. We also prove a certain estimate for a trigonometric integral, which is useful in studying nonisotropic singular integrals.

## 1. INTRODUCTION

Let  $P$  be an  $n \times n$  real matrix whose eigenvalues have positive real parts. Let  $\gamma = \text{trace } P$ . Define a dilation group  $\{A_t\}_{t>0}$  on  $\mathbb{R}^n$  by  $A_t = t^P = \exp((\log t)P)$ . We assume  $n \geq 2$ . There is a non-negative function  $r$  on  $\mathbb{R}^n$  associated with  $\{A_t\}_{t>0}$ . The function  $r$  is continuous on  $\mathbb{R}^n$  and infinitely differentiable in  $\mathbb{R}^n \setminus \{0\}$ ; furthermore it satisfies

- (1)  $r(A_t x) = tr(x)$  for all  $t > 0$  and  $x \in \mathbb{R}^n$ ;
- (2)  $r(x + y) \leq C(r(x) + r(y))$  for some  $C > 0$ ;
- (3) if  $\Sigma = \{x \in \mathbb{R}^n : r(x) = 1\}$ , then  $\Sigma = \{\theta \in \mathbb{R}^n : \langle B\theta, \theta \rangle = 1\}$  for a positive symmetric matrix  $B$ , where  $\langle \cdot, \cdot \rangle$  denotes the inner product in  $\mathbb{R}^n$ .

Also, we have  $dx = t^{\gamma-1} d\sigma dt$ , that is,

$$\int_{\mathbb{R}^n} f(x) dx = \int_0^\infty \int_{\Sigma} f(A_t \theta) t^{\gamma-1} d\sigma(\theta) dt$$

for appropriate functions  $f$ , where  $d\sigma$  is a  $C^\infty$  measure on  $\Sigma$ . See [2, 13, 17] for more details.

Let  $\Omega$  be locally integrable in  $\mathbb{R}^n \setminus \{0\}$  and homogeneous of degree 0 with respect to the dilation group  $\{A_t\}$ , that is,  $\Omega(A_t x) = \Omega(x)$  for  $x \neq 0$ . We assume that

$$\int_{\Sigma} \Omega(\theta) d\sigma(\theta) = 0.$$

For  $s \geq 1$ , let  $\Delta_s$  denote the collection of measurable functions  $h$  on  $\mathbb{R}_+ = \{t \in \mathbb{R} : t > 0\}$  satisfying

$$\|h\|_{\Delta_s} = \sup_{j \in \mathbb{Z}} \left( \int_{2^j}^{2^{j+1}} |h(t)|^s dt/t \right)^{1/s} < \infty,$$

where  $\mathbb{Z}$  denotes the set of integers. We define  $\|h\|_{\Delta_\infty}$  as usual ( $\|h\|_{\Delta_\infty} = \|h\|_{L^\infty(\mathbb{R}_+)}$ ).

---

2000 *Mathematics Subject Classification.* Primary 42B20, 42B25.

*Key Words and Phrases.* Nonisotropic singular integrals, extrapolation, trigonometric integrals.

Let  $\Gamma : [0, \infty) \rightarrow \mathbb{R}^m$  be a continuous mapping satisfying  $\Gamma(0) = 0$ . We define a singular integral operator along the surface  $(y, \Gamma(r(y)))$  by

$$(1.1) \quad \begin{aligned} Tf(x, z) &= \text{p.v.} \int_{\mathbb{R}^n} f(x - y, z - \Gamma(r(y))) K(y) dy \\ &= \lim_{\epsilon \rightarrow 0} \int_{r(y) > \epsilon} f(x - y, z - \Gamma(r(y))) K(y) dy, \end{aligned}$$

where  $K(y) = h(r(y))\Omega(y')r(y)^{-\gamma}$ ,  $y' = A_{r(y)^{-1}}y$  and  $h \in \Delta_1$ . We assume that the principal value integral in (1.1) exists for every  $(x, z) \in \mathbb{R}^n \times \mathbb{R}^m$  and  $f \in \mathcal{S}(\mathbb{R}^n \times \mathbb{R}^m)$  (the Schwartz class).

We denote by  $L \log L(\Sigma)$  the Zygmund class of all those functions  $\Omega$  on  $\Sigma$  which satisfy

$$\int_{\Sigma} |\Omega(\theta)| \log(2 + |\Omega(\theta)|) d\sigma(\theta) < \infty.$$

Also, we consider the  $L^q(\Sigma)$  spaces and write  $\|\Omega\|_q = (\int_{\Sigma} |\Omega(\theta)|^q d\sigma(\theta))^{1/q}$  for  $\Omega \in L^q(\Sigma)$  ( $\|\Omega\|_{\infty}$  is defined as usual).

Let

$$M_{\Gamma}g(z) = \sup_{R > 0} R^{-1} \int_0^R |g(z - \Gamma(t))| dt.$$

We assume that the maximal operator  $M_{\Gamma}$  is bounded on  $L^p(\mathbb{R}^m)$  for all  $p > 1$ . See [15, 17] for examples of such functions  $\Gamma$ .

In this note we prove the following.

**Theorem 1.** *Let  $T$  be as in (1.1). Suppose that  $\Omega \in L^q(\Sigma)$  for some  $q \in (1, 2]$  and  $h \in \Delta_s$  for some  $s > 1$ . Then, we have*

$$\|Tf\|_{L^p(\mathbb{R}^{n+m})} \leq C_p (q-1)^{-1} \|\Omega\|_q \|h\|_{\Delta_s} \|f\|_{L^p(\mathbb{R}^{n+m})}$$

if  $|1/p - 1/2| < \min(1/s', 1/2)$ , where  $1/s' + 1/s = 1$  and the constant  $C_p$  is independent of  $q$  and  $\Omega$ .

**Theorem 2.** *Suppose  $\Omega \in L \log L(\Sigma)$  and  $h \in \Delta_s$  for some  $s > 1$ . Then,  $T$  is bounded on  $L^p(\mathbb{R}^{n+m})$  if  $|1/p - 1/2| < \min(1/s', 1/2)$ .*

Theorem 2 follows from Theorem 1 by an extrapolation method. When  $r(x) = |x|$  (the Euclid norm),  $m = 1$  and  $\Gamma$  is a  $C^2$ , convex, increasing function, Theorem 2 was proved in A. Al-Salman and Y. Pan [1] (see [1, Theorem 4.1] and also [10] for a related result). In [1], it is noted that the estimates as  $q \rightarrow 1$  of Theorem 1 (in their setting) can be used through extrapolation to prove the  $L^p$  boundedness of [1, Theorem 4.1], although such estimates are yet to be proved. In this note, we are able to prove Theorem 1 and apply it to prove Theorem 2.

If  $\Gamma \equiv 0$  ( $\Gamma$  is identically 0), then  $T$  essentially reduces to the lower dimensional singular integral

$$(1.2) \quad Sf(x) = \text{p.v.} \int_{\mathbb{R}^n} f(x - y) K(y) dy.$$

For this singular integral we have the following.

**Theorem 3.** *Let  $\Omega \in L^q(\Sigma)$  and  $h \in \Delta_s$  for some  $q, s \in (1, 2]$ . Then we have*

$$\|Sf\|_{L^p(\mathbb{R}^n)} \leq C_p (q-1)^{-1} (s-1)^{-1} \|\Omega\|_q \|h\|_{\Delta_s} \|f\|_{L^p(\mathbb{R}^n)}$$

for all  $p \in (1, \infty)$ , where the constant  $C_p$  is independent of  $q, s, \Omega$  and  $h$ .

For  $a > 0$ , let

$$L_a(h) = \sup_{j \in \mathbb{Z}} \int_{2^j}^{2^{j+1}} |h(r)| (\log(2 + |h(r)|))^a dr/r.$$

We define a class  $\mathcal{L}_a$  to be the space of all those measurable functions  $h$  on  $\mathbb{R}_+$  which satisfy  $L_a(h) < \infty$ .

By Theorem 3 and an extrapolation we have the following.

**Theorem 4.** *Suppose  $\Omega \in L \log L(\Sigma)$  and  $h \in \mathcal{L}_a$  for some  $a > 2$ . Then  $S$  is bounded on  $L^p(\mathbb{R}^n)$  for all  $p \in (1, \infty)$ .*

It is noted in [5] that  $S$  is bounded on  $L^p$ ,  $1 < p < \infty$ , if  $\Omega \in L^q$  for some  $q > 1$  and  $h \in \Delta_2$  (see [5, Corollary 4.5]). Theorem 4 improves that result. See [13, 16] for nonisotropic singular integrals  $S$  with  $h \equiv 1$  and also [3, 7, 9, 12] for related results.

In Section 2, we prove Theorems 1 and 3. The proofs are based on the method of [5]. As in [14], a key idea of the proof of Theorem 1 is to use a Littlewood–Paley decomposition depending on  $q$  for which  $\Omega \in L^q$ . Theorem 3 is proved in a similar fashion. Applying an extrapolation argument, we can prove Theorems 2 and 4 from Theorems 1 and 3, respectively. We give a proof of Theorem 4 in Section 3. In Section 4, we prove an estimate for a trigonometric integral, a corollary of which is used in proving Theorems 1 and 3.

Throughout this note, the letter  $C$  will be used to denote non-negative constants which may be different in different occurrences.

## 2. PROOFS OF THEOREMS 1 AND 3

Let  $A^*$  denote the adjoint of a matrix  $A$ . Then  $A_t^* = \exp((\log t)P^*)$ . We write  $A_t^* = B_t$ . We can define a non-negative function  $s$  from  $\{B_t\}$  exactly in the same way as we define  $r$  from  $\{A_t\}$ .

There are positive constants  $c_1, c_2, c_3, c_4, \alpha_1, \alpha_2, \beta_1$  and  $\beta_2$  such that

$$\begin{aligned} c_1|x|^{\alpha_1} < r(x) < c_2|x|^{\alpha_2} & \text{ if } r(x) \geq 1, \\ c_3|x|^{\beta_1} < r(x) < c_4|x|^{\beta_2} & \text{ if } 0 < r(x) \leq 1. \end{aligned}$$

Also, we have

$$\begin{aligned} d_1|\xi|^{a_1} < s(\xi) < d_2|\xi|^{a_2} & \text{ if } s(\xi) \geq 1, \\ d_3|\xi|^{b_1} < s(\xi) < d_4|\xi|^{b_2} & \text{ if } 0 < s(\xi) \leq 1 \end{aligned}$$

for some positive numbers  $d_1, d_2, d_3, d_4, a_1, a_2, b_1$  and  $b_2$  (see [17]). These estimates are useful in the following.

We consider the singular integral operator  $T$  defined in (1.1). Let  $E_j = \{x \in \mathbb{R}^n : \beta^j < r(x) \leq \beta^{j+1}\}$ , where  $\beta \geq 2$  and  $j \in \mathbb{Z}$ . We define a sequence of Borel measures  $\{\sigma_j\}$  on  $\mathbb{R}^n \times \mathbb{R}^m$  by

$$\hat{\sigma}_j(\xi, \eta) = \int_{E_j} e^{-2\pi i \langle y, \xi \rangle} e^{-2\pi i \langle \Gamma(r(y)), \eta \rangle} K(y) dy,$$

where  $\hat{\sigma}_j$  denotes the Fourier transform of  $\sigma_j$  defined by

$$\hat{\sigma}_j(\xi, \eta) = \int e^{-2\pi i \langle (x, z), (\xi, \eta) \rangle} d\sigma_j(x, z).$$

Then  $Tf(x) = \sum_{-\infty}^{\infty} \sigma_k * f(x)$ .

Let  $\mu_k = |\sigma_k|$ , where  $|\sigma_k|$  denotes the total variation of  $\sigma_k$ . Let  $\Omega \in L^q$ ,  $h \in \Delta_s$ ,  $q, s \in (1, 2]$ . We prove the following estimates (2.1)–(2.5):

$$(2.1) \quad \|\sigma_k\| \leq C(\log \beta) \|\Omega\|_1 \|h\|_{\Delta_1} \leq C(\log \beta) \|\Omega\|_q \|h\|_{\Delta_s},$$

where  $\|\sigma_k\| = |\sigma_k|(\mathbb{R}^{n+m})$ ;

$$(2.2) \quad |\hat{\sigma}_k(\xi, \eta)| \leq C \|\Omega\|_q \|h\|_{\Delta_s} (\beta^{k+d} s(\xi))^{1/b_1},$$

where  $d = b_1/\alpha_1$ ;

$$(2.3) \quad |\hat{\sigma}_k(\xi, \eta)| \leq C(\log \beta) \|\Omega\|_q \|h\|_{\Delta_s} (\beta^k s(\xi))^{-\epsilon_0/(q's')}$$

for some  $\epsilon_0 > 0$ ;

$$(2.4) \quad |\hat{\mu}_k(\xi, \eta)| \leq C(\log \beta) \|\Omega\|_q \|h\|_{\Delta_s} (\beta^k s(\xi))^{-\epsilon_0/(q's')},$$

where  $\epsilon_0$  is as in (2.3);

$$(2.5) \quad |\hat{\mu}_k(\xi, \eta) - \hat{\mu}_k(0, \eta)| \leq C \|\Omega\|_q \|h\|_{\Delta_s} (\beta^{k+d} s(\xi))^{1/b_1},$$

where  $d$  is as in (2.2).

First we see that

$$(2.6) \quad \|\sigma_k\|_1 = \int_{\beta^k}^{\beta^{k+1}} |h(r)| \|\Omega\|_1 dr/r \leq C(\log \beta) \|\Omega\|_1 \|h\|_{\Delta_1}.$$

From this, (2.1) follows. Next, we show (2.2). Take  $\nu \in \mathbb{Z}$  so that  $2^\nu < \beta \leq 2^{\nu+1}$ . Note that

$$\hat{\sigma}_k(\xi, \eta) = \int_{\beta^k < r(x) \leq \beta^{k+1}} e^{-2\pi i \langle \Gamma(r(x)), \eta \rangle} (e^{-2\pi i \langle x, \xi \rangle} - 1) h(r(x)) \Omega(x') r(x)^{-\gamma} dx.$$

Thus

$$(2.7) \quad \begin{aligned} |\hat{\sigma}_k(\xi, \eta)| &\leq C \int_{1 < r(x) \leq \beta} |x|_{B_{\beta^k} \xi} |h(\beta^k r(x)) \Omega(x')| r(x)^{-\gamma} dx \\ &\leq C \sum_{j=0}^{\nu} |B_{\beta^k} \xi| \|\Omega\|_1 2^{j/\alpha_1} \int_{2^j}^{2^{j+1}} |h(\beta^k r)| dr/r \\ &\leq C \beta^{1/\alpha_1} |B_{\beta^k} \xi| \|\Omega\|_1 \|h\|_{\Delta_1}. \end{aligned}$$

Combining (2.6) and (2.7), we have

$$(2.8) \quad |\hat{\sigma}_k(\xi, \eta)| \leq C \|\Omega\|_1 \|h\|_{\Delta_1} \min \left( \log \beta, \beta^{1/\alpha_1} |B_{\beta^k} \xi| \right).$$

If  $s(B_{\beta^k} \xi) < 1$ , then  $|B_{\beta^k} \xi| \leq C(\beta^k s(\xi))^{1/b_1}$ . Therefore,

$$\min \left( \log \beta, \beta^{1/\alpha_1} |B_{\beta^k} \xi| \right) \leq C(\beta^{k+d} s(\xi))^{1/b_1}.$$

Using this in (2.8), we have (2.2). We can prove (2.5) in the same way.

Next we prove (2.3). We use a method similar to that of [5, p. 551]. Define

$$\tau(\xi) = \int_{\Sigma} \Omega(\theta) e^{-2\pi i \langle \xi, \theta \rangle} d\sigma(\theta).$$

We need the following estimates.

**Lemma 1.** *Let  $L$  be the degree of the minimal polynomial of  $P$ . Then, if  $0 < \epsilon_0 < a_2^{-1} \min(1/2, q'/L)$ , we have*

$$\int_{\beta^k}^{\beta^{k+1}} |\tau(B_r \xi)|^2 dr/r \leq C(\log \beta)(\beta^k s(\xi))^{-\epsilon_0/q'} \|\Omega\|_q^2,$$

where  $C$  is independent of  $\Omega \in L^q$ ,  $q \in (1, 2]$  and  $\beta$ .

In proving Lemma 1 we use the following estimate, which follows from the corollary to Theorem 5 in Section 4 via an integration by parts argument.

**Lemma 2.** *Let  $L$  be as in Lemma 1. Then, for  $\eta, \zeta \in \mathbb{R}^n \setminus \{0\}$  we have*

$$\left| \int_1^2 \exp(i\langle B_t \eta, \zeta \rangle) dt/t \right| \leq C |\langle \eta, P\zeta \rangle|^{-1/L}$$

for some positive constant  $C$  independent of  $\eta$  and  $\zeta$ .

Proof of Lemma 1. Choose  $\nu \in \mathbb{Z}$  such that  $2^\nu < \beta \leq 2^{\nu+1}$ . Then, we have

$$\begin{aligned} \int_{\beta^k}^{\beta^{k+1}} |\tau(B_r \xi)|^2 dr/r &\leq \sum_{j=0}^{\nu} \int_{\beta^{k+2j}}^{\beta^{k+2j+1}} |\tau(B_r \xi)|^2 dr/r \\ &= \sum_{j=0}^{\nu} \iint_{\Sigma \times \Sigma} \left( \int_1^2 \exp(-2\pi i \langle B_{\beta^{k+2j} r} \xi, \theta - \omega \rangle) dr/r \right) \Omega(\theta) \bar{\Omega}(\omega) d\sigma(\theta) d\sigma(\omega). \end{aligned}$$

By Lemma 2 we have

$$\left| \int_1^2 \exp(-2\pi i \langle B_{\beta^{k+2j} r} \xi, \theta - \omega \rangle) dr/r \right| \leq C |\langle B_{\beta^{k+2j} \xi}, P(\theta - \omega) \rangle|^{-\epsilon},$$

where  $0 < \epsilon \leq 1/L$ . Using Hölder's inequality, if  $0 < \epsilon < \min(1/(2q'), 1/L)$ , we see that

$$\begin{aligned} &\iint_{\Sigma \times \Sigma} |\langle B_{\beta^{k+2j} \xi}, P(\theta - \omega) \rangle|^{-\epsilon} |\Omega(\theta) \bar{\Omega}(\omega)| d\sigma(\theta) d\sigma(\omega) \\ &\leq \left( \iint_{\Sigma \times \Sigma} |P^* B_{\beta^{k+2j} \xi}, \theta - \omega|^{-\epsilon q'} d\sigma(\theta) d\sigma(\omega) \right)^{1/q'} \|\Omega\|_q^2 \leq C |B_{\beta^{k+2j} \xi}|^{-\epsilon} \|\Omega\|_q^2, \end{aligned}$$

where the last inequality follows from (3) of Section 1 (see [5, p. 553]). Therefore

(2.9)

$$\int_{\beta^k}^{\beta^{k+1}} |\tau(B_r \xi)|^2 dr/r \leq C \|\Omega\|_q^2 \sum_{j=0}^{\nu} |B_{\beta^{k+2j} \xi}|^{-\epsilon} \quad (0 < \epsilon < \min(1/(2q'), 1/L)).$$

If  $s(B_{\beta^k} \xi) \geq 1$ ,  $|B_{\beta^{k+2j} \xi}| \geq C(\beta^{k+2j} s(\xi))^{1/a_2}$  ( $0 \leq j \leq \nu$ ). Thus we see that

$$(2.10) \quad \sum_{j=0}^{\nu} |B_{\beta^{k+2j} \xi}|^{-\epsilon} \leq \sum_{j=0}^{\nu} C(\beta^{k+2j} s(\xi))^{-\epsilon/a_2} \leq C(\log \beta)(\beta^k s(\xi))^{-\epsilon/a_2},$$

where  $C$  is independent of  $q$ . By (2.9) and (2.10) we have the estimate of Lemma 1 when  $s(B_{\beta^k} \xi) \geq 1$ . If  $s(B_{\beta^k} \xi) < 1$ , the estimate of Lemma 1 follows from the inequality  $|\tau(\xi)| \leq \|\Omega\|_1$ . This completes the proof of Lemma 1.

Now, by Hölder's inequality we have

$$\begin{aligned}
(2.11) \quad |\hat{\sigma}_k(\xi, \eta)| &= \left| \int_{\beta^k}^{\beta^{k+1}} e^{-2\pi i \langle \Gamma(r), \eta \rangle} h(r) \tau(B_r \xi) dr/r \right| \\
&\leq \left( \int_{\beta^k}^{\beta^{k+1}} |h(r)|^s dr/r \right)^{1/s} \left( \int_{\beta^k}^{\beta^{k+1}} |\tau(B_r \xi)|^{s'} dr/r \right)^{1/s'} \\
&\leq C(\log \beta)^{1/s} \|h\|_{\Delta_s} \|\Omega\|_1^{(s'-2)/s'} \left( \int_{\beta^k}^{\beta^{k+1}} |\tau(B_r \xi)|^2 dr/r \right)^{1/s'},
\end{aligned}$$

where we have used the estimate  $|\tau(\xi)| \leq \|\Omega\|_1$  to get the last inequality. By (2.11) and Lemma 1 we have (2.3). The estimate (2.4) can be proved similarly.

Let  $B_{qs} = (1 - \beta^{-\theta \epsilon_0 / (q' s')})^{-1}$ , where  $\beta \geq 2$ ,  $\theta \in (0, 1)$  and  $\epsilon_0$  is as in (2.3) and (2.4). To prove Theorems 1 and 3, we use the following:

**Proposition 1.** *Suppose that  $\Omega \in L^q$ ,  $q \in (1, 2]$  and  $h \in \Delta_s$ ,  $s \in (1, 2]$ . Let  $|1/p - 1/2| < (1 - \theta)/(s'(1 + \theta))$ . Then, we have*

$$\|Tf\|_p \leq C(\log \beta) \|h\|_{\Delta_s} \|\Omega\|_q B_{qs} B_{q2}^{1/p - 1/p'} \|f\|_p,$$

where  $C$  is a constant independent of  $\Omega$ ,  $h$ ,  $q$ ,  $s$  and  $\beta$ .

**Proposition 2.** *Suppose that  $\Gamma \equiv 0$ . Let  $\Omega \in L^q$ ,  $h \in \Delta_s$ ,  $q, s \in (1, 2]$ . Then, for  $p \in (1 + \theta, (1 + \theta)/\theta)$  we have*

$$\|Tf\|_p \leq C(\log \beta) \|\Omega\|_q \|h\|_{\Delta_s} B_{qs}^{1 + |1/p - 1/p'|} \|f\|_p,$$

where  $C$  is a constant independent of  $\Omega$ ,  $h$ ,  $q$ ,  $s$  and  $\beta$ .

To prove Propositions 1 and 2, we need the following:

**Proposition 3.** *Let  $\mu^*(f)(x) = \sup_k |\mu_k * f(x)|$ . Let  $\Omega \in L^q$ ,  $q \in (1, 2]$ .*

(1) *If  $h \in \Delta_\infty$ , for  $p > 1 + \theta$  we have*

$$\|\mu^*(f)\|_p \leq C(\log \beta) \|\Omega\|_q \|h\|_{\Delta_\infty} B_{q2}^{2/p} \|f\|_p,$$

where  $C$  is a constant independent of  $\Omega$ ,  $h$ ,  $q$  and  $\beta$ .

(2) *Suppose that  $\Gamma \equiv 0$ . Let  $h \in \Delta_s$ ,  $s \in (1, 2]$ . Then, we have*

$$\|\mu^*(f)\|_p \leq C(\log \beta) \|\Omega\|_q \|h\|_{\Delta_s} B_{qs}^{2/p} \|f\|_p$$

for  $p > 1 + \theta$ , where  $C$  is independent of  $\Omega$ ,  $q$ ,  $h$ ,  $s$  and  $\beta$ .

*Proof.* Since the estimate  $\|\mu^*(f)\|_\infty \leq C(\log \beta) \|\Omega\|_1 \|h\|_{\Delta_1} \|f\|_\infty$  follows from (2.1), by interpolation, to prove (1) and (2) of Proposition 3 we may assume  $p \in (1 + \theta, 2]$ .

First, we give a proof of part (1). Define measures  $\nu_k$  on  $\mathbb{R}^n \times \mathbb{R}^m$  by

$$\hat{\nu}_k(\xi, \eta) = \hat{\mu}_k(\xi, \eta) - \hat{\Psi}_k(\xi, \eta),$$

where  $\hat{\Psi}_k(\xi, \eta) = \hat{\varphi}_k(\xi) \hat{\mu}_k(0, \eta)$  with  $\varphi_k(x) = \beta^{-k\gamma} \varphi(A_{\beta^{-k}} x)$ ,  $\varphi \in C_0^\infty$ . We assume that  $\varphi$  is supported in  $\{r(x) \leq 1\}$ ,  $\hat{\varphi}(0) = 1$  and  $\varphi \geq 0$ . Then by (2.1), (2.4) and (2.5), for  $q, s \in (1, 2]$ , we have

$$|\hat{\nu}_k(\xi, \eta)| \leq C(\log \beta) \|\Omega\|_q \|h\|_{\Delta_s} \min \left( 1, (\beta^{k+d} s(\xi))^{1/b_1}, (\beta^k s(\xi))^{-\epsilon_0/(q' s')} \right).$$

We may assume that  $\epsilon_0$  is small enough so that  $\epsilon_0/4 \leq 1/b_1$ . Then, we see that

$$(2.12) \quad |\hat{\nu}_k(\xi, \eta)| \leq CA \min \left( 1, (\beta^{k+d} s(\xi))^\alpha, (\beta^k s(\xi))^{-\alpha} \right),$$

where  $A = (\log \beta) \|\Omega\|_q \|h\|_{\Delta_\infty}$  and  $\alpha = \epsilon_0/(2q')$ .

Let

$$g(f)(x, z) = \left( \sum_{k=-\infty}^{\infty} |\nu_k * f(x, z)|^2 \right)^{1/2}.$$

Then  $\mu^*(f) \leq g(f) + \Psi^*(|f|)$ , where  $\Psi^*(f) = \sup_k \|\Psi_k\| * |f|$ . Let

$$Mg(x) = \sup_{t>0} t^{-\gamma} \int_{r(x-y)<t} |g(y)| dy$$

be the Hardy–Littlewood maximal function on  $\mathbb{R}^n$  with respect to the function  $r$ . By the  $L^p$  boundedness of  $M_\Gamma$  and  $M$ , it is easy to see that  $\|\Psi^*(f)\|_p \leq CA\|f\|_p$  for  $p > 1$ . Thus to prove Proposition 3 (1) it suffices to show

$$(2.13) \quad \|g(f)\|_p \leq CAB^{2/p} \|f\|_p \quad (p \in (1 + \theta, 2]),$$

where  $A$  is as above and  $B = B_{q_2}$ . By a well-known property of Rademacher's functions, (2.13) follows from

$$(2.14) \quad \|U_\epsilon(f)\|_p \leq CAB^{2/p} \|f\|_p \quad (p \in (1 + \theta, 2]),$$

where  $U_\epsilon(f)(x, z) = \sum \epsilon_k \nu_k * f(x, z)$  with  $\epsilon = \{\epsilon_k\}$ ,  $\epsilon_k = 1$  or  $-1$  (the inequality is uniform in  $\epsilon$ ).

We define two sequences  $\{r_m\}_1^\infty$  and  $\{p_m\}_1^\infty$  by  $p_1 = 2$  and

$$\frac{1}{r_m} - \frac{1}{2} = \frac{1}{2p_m}, \quad \frac{1}{p_{m+1}} = \frac{\theta}{2} + \frac{1-\theta}{r_m} \quad \text{for } m \geq 1.$$

Then, we have

$$\frac{1}{p_{m+1}} = \frac{1}{2} + \frac{1-\theta}{2p_m} \quad \text{for } m \geq 1.$$

Thus  $1/p_m = (1 - \eta^m)/(1 + \theta)$ , where  $\eta = (1 - \theta)/2$ , so  $\{p_m\}$  is decreasing and converges to  $1 + \theta$ .

For  $j \geq 1$  we prove

$$(2.15) \quad \|U_\epsilon(f)\|_{p_j} \leq C_j AB^{2/p_j} \|f\|_{p_j}.$$

To prove (2.15) we use the Littlewood–Paley theory. Let  $\{\psi_k\}_\infty^\infty$  be a sequence of non-negative functions in  $C^\infty((0, \infty))$  such that

$$\begin{aligned} \text{supp}(\psi_k) &\subset [\beta^{-k-1}, \beta^{-k+1}], \quad \sum_k \psi_k(t)^2 = 1, \\ |(d/dt)^j \psi_k(t)| &\leq c_j/t^j \quad (j = 1, 2, \dots), \end{aligned}$$

where  $c_j$  is independent of  $\beta \geq 2$ . Define  $S_k$  by

$$(S_k(f))^\wedge(\xi, \eta) = \psi_k(s(\xi)) \hat{f}(\xi, \eta).$$

We write  $U_\epsilon(f) = \sum_{j=-\infty}^\infty U_j(f)$ , where  $U_j(f) = \sum_{k=-\infty}^\infty \epsilon_k S_{j+k}(\nu_k * S_{j+k}(f))$ . Then by Plancherel's theorem and (2.12) we have

$$\begin{aligned} (2.16) \quad \|U_j(f)\|_2^2 &\leq \sum_k C \iint_{D(j+k) \times \mathbb{R}^m} |\hat{\nu}_k(\xi, \eta)|^2 |\hat{f}(\xi, \eta)|^2 d\xi d\eta \\ &\leq CA^2 \min\left(1, \beta^{-2(|j|-1-d)\alpha}\right) \sum_k \iint_{D(j+k) \times \mathbb{R}^m} |\hat{f}(\xi, \eta)|^2 d\xi d\eta \\ &\leq CA^2 \min\left(1, \beta^{-2(|j|-1-d)\alpha}\right) \|f\|_2^2, \end{aligned}$$



where  $D(k) = \{\xi \in \mathbb{R}^n : \beta^{-k-1} < s(\xi) \leq \beta^{-k+1}\}$ . By (2.16) we have

$$(2.17) \quad \begin{aligned} \|U_\epsilon(f)\|_2 &\leq \sum_{-\infty}^{\infty} \|U_j(f)\|_2 \leq C \sum_{-\infty}^{\infty} A \min\left(1, \beta^{-(|j|-1-d)\alpha}\right) \|f\|_2 \\ &\leq CA(1 - \beta^{-\alpha})^{-1} \|f\|_2. \end{aligned}$$

If we denote by  $A(m)$  the estimate of (2.15) for  $j = m$ , this proves  $A(1)$ .

Now, we assume  $A(m)$  and derive  $A(m+1)$  from  $A(m)$ . Note that

$$\nu^*(f) \leq \mu^*(|f|) + \Psi^*(|f|) \leq g(|f|)(x) + 2\Psi^*(|f|),$$

where  $\nu^*(f)(x) = \sup_k |\nu_k| * f(x)$ . Since  $\|g(f)\|_{p_m} \leq CAB^{2/p_m} \|f\|_{p_m}$  by  $A(m)$ , we have

$$\|\nu^*(f)\|_{p_m} \leq CAB^{2/p_m} \|f\|_{p_m}.$$

Also,  $\|\nu_k\| \leq CA$  by (2.1). Thus, by the proof of Lemma for Theorem B in [5, p. 544], we have the vector valued inequality:

$$(2.18) \quad \begin{aligned} \left\| \left( \sum |\nu_k * g_k|^2 \right)^{1/2} \right\|_{r_m} &\leq C (AB^{2/p_m} \sup_k \|\nu_k\|)^{1/2} \left\| \left( \sum |g_k|^2 \right)^{1/2} \right\|_{r_m} \\ &\leq CAB^{1/p_m} \left\| \left( \sum |g_k|^2 \right)^{1/2} \right\|_{r_m}. \end{aligned}$$

By (2.18) and the Littlewood–Paley inequality, we have

$$(2.19) \quad \begin{aligned} \|U_j(f)\|_{r_m} &\leq C \left\| \left( \sum_k |\nu_k * S_{j+k}(f)|^2 \right)^{1/2} \right\|_{r_m} \\ &\leq CAB^{1/p_m} \|f\|_{r_m}. \end{aligned}$$

Here we note that the bounds for the Littlewood–Paley inequality are independent of  $\beta \geq 2$ . Interpolating between (2.16) and (2.19), we have

$$\|U_j(f)\|_{p_{m+1}} \leq CAB^{(1-\theta)/p_m} \min\left(1, \beta^{-\theta\alpha(|j|-1-d)}\right) \|f\|_{p_{m+1}}.$$

Thus

$$\begin{aligned} \|U_\epsilon(f)\|_{p_{m+1}} &\leq \sum_j \|U_j(f)\|_{p_{m+1}} \leq CAB^{(1-\theta)/p_m} (1 - \beta^{-\theta\alpha})^{-1} \|f\|_{p_{m+1}} \\ &\leq CAB^{2/p_{m+1}} \|f\|_{p_{m+1}}, \end{aligned}$$

which proves  $A(m+1)$ . By induction, this completes the proof of (2.15).

Now we prove (2.14). Let  $p \in (1 + \theta, 2]$  and let  $\{p_m\}_1^\infty$  be as in (2.15). Then we have  $p_{N+1} < p \leq p_N$  for some  $N$ . By interpolation between the estimates in (2.15) for  $j = N$  and  $j = N+1$  we have (2.14). This completes the proof of Proposition 3 (1).

Part (2) of Proposition 3 can be proved in the same way. We take  $A = (\log \beta) \|\Omega\|_q \|h\|_{\Delta_s}$  and  $\alpha = \epsilon_0/(q's')$  in (2.12). Then, since

$$\|\Psi^*(f)\|_p \leq C(\log \beta) \|\Omega\|_1 \|h\|_{\Delta_1} \|f\|_p \quad \text{for } p > 1$$

if  $\Gamma \equiv 0$ , the proof of part (1) can be used to get (2.13) with  $A = (\log \beta) \|\Omega\|_q \|h\|_{\Delta_s}$  as above and  $B = B_{qs}$ , and the conclusion of part (2) follows from (2.13).  $\square$

Proof of Proposition 1. To prove Proposition 1 we may assume  $1 < s < 2$ . As in [1], here we apply an idea in the proof of [6, Theorem 7.5]. We consider measures  $\tau_k$  defined by

$$\hat{\tau}_k(\xi, \eta) = \int_{E_k} e^{-2\pi i \langle y, \xi \rangle} e^{-2\pi i \langle \Gamma(r(y)), \eta \rangle} |h(r(y))|^{2-s} |\Omega(y')| r(y)^{-\gamma} dy.$$

Then, the Schwarz inequality implies

$$(2.20) \quad |\sigma_k * f|^2 \leq C(\log \beta) \|h\|_{\Delta_s}^s \|\Omega\|_1 \tau_k * |f|^2.$$

Define measures  $\lambda_k$  by

$$\hat{\lambda}_k(\xi, \eta) = \int_{E_k} e^{-2\pi i \langle y, \xi \rangle} e^{-2\pi i \langle \Gamma(r(y)), \eta \rangle} |\Omega(y')| r(y)^{-\gamma} dy.$$

Since  $|h|^{2-s} \in \Delta_{s/(2-s)}$  and  $\| |h|^{2-s} \|_{\Delta_{s/(2-s)}} = \|h\|_{\Delta_s}^{2-s}$ , if  $u = s/(2-s)$  by Hölder's inequality we have

$$|\tau_k * f| \leq C(\log \beta)^{1/u} \|h\|_{\Delta_s}^{2-s} \|\Omega\|_1^{1/u} (\lambda_k * |f|^{u'})^{1/u'}.$$

Therefore, if  $1 + \theta < r/u' = 2r(s-1)/s$ , by applying (1) of Proposition 3 to  $\{\lambda_k\}$  we see that

$$(2.21) \quad \|\tau^*(f)\|_r \leq C(\log \beta) \|h\|_{\Delta_s}^{2-s} \|\Omega\|_q B_{q^2}^{2/r} \|f\|_r,$$

where  $\tau^*(f) = \sup_k |\tau_k * f|$ . Thus, if  $|1/v - 1/2| = 1/(2r) < 1/(s'(1+\theta))$ , using (2.20), (2.21) and arguing as in the proof of Lemma for Theorem B in [5, p. 544], we see that

$$(2.22) \quad \left\| \left( \sum |\sigma_k * g_k|^2 \right)^{1/2} \right\|_v \leq C(\log \beta) \|h\|_{\Delta_s} \|\Omega\|_q B_{q^2}^{1/r} \left\| \left( \sum |g_k|^2 \right)^{1/2} \right\|_v.$$

We decompose  $Tf = \sum_{j=-\infty}^{\infty} V_j f$ , where  $V_j f = \sum_{k=-\infty}^{\infty} S_{j+k} (\sigma_k * S_{j+k}(f))$ . Then, using (2.22) and the Littlewood–Paley theory, we see that

$$(2.23) \quad \|V_j f\|_v \leq C(\log \beta) \|h\|_{\Delta_s} \|\Omega\|_q B_{q^2}^{1/r} \|f\|_v,$$

where  $|1/v - 1/2| = 1/(2r) < 1/(s'(1+\theta))$ . On the other hand, by (2.1)–(2.3) we have

$$|\hat{\sigma}_k(\xi, \eta)| \leq C(\log \beta) \|\Omega\|_q \|h\|_{\Delta_s} \min(1, (\beta^{k+d} s(\xi))^\kappa, (\beta^k s(\xi))^{-\kappa}),$$

where  $\kappa = \epsilon_0/(q's')$ , and hence, similarly to the proof of (2.16), we can show that

$$(2.24) \quad \|V_j f\|_2 \leq C(\log \beta) \|h\|_{\Delta_s} \|\Omega\|_q \min(1, \beta^{-(|j|-1-d)\kappa}) \|f\|_2.$$

If  $|1/p - 1/2| < (1-\theta)/(s'(1+\theta))$ , then we can find numbers  $v$  and  $r$  such that  $|1/v - 1/2| = 1/(2r) < 1/(s'(1+\theta))$  and  $1/p = \theta/2 + (1-\theta)/v$ . Thus, interpolating between (2.23) and (2.24), we have

$$\|V_j f\|_p \leq C(\log \beta) \|h\|_{\Delta_s} \|\Omega\|_q B_{q^2}^{(1-\theta)/r} \min(1, \beta^{-\theta(|j|-1-d)\kappa}) \|f\|_p.$$

Therefore

$$(2.25) \quad \|Tf\|_p \leq \sum_j \|V_j f\|_p \leq C(\log \beta) \|h\|_{\Delta_s} \|\Omega\|_q B_{q^2}^{(1-\theta)/r} B_{qs} \|f\|_p.$$

This completes the proof of Proposition 1, since  $(1-\theta)/r = |1/p - 1/p'|$ .

Proof of Proposition 2. The  $L^2$  estimates follow from Proposition 1, so on account of duality and interpolation we may assume that  $1+\theta < p \leq 4/(3-\theta)$ . For  $p_0 \in (1+\theta, 4/(3-\theta)]$  we can find  $r \in (1+\theta, 2]$  such that  $1/p_0 = 1/2 + (1-\theta)/(2r)$ .

If  $\Gamma \equiv 0$ , by (2) of Proposition 3 and (2.1), arguing as in (2.18), we have (2.22) with  $B_{q_2}$  replaced by  $B_{q_s}$  for the number  $v$  satisfying  $1/v - 1/2 = 1/(2r)$  (note that  $1/p_0 = \theta/2 + (1 - \theta)/v$ ). Thus, arguing as in the proof of Proposition 1, we have (2.25) with  $p = p_0$  and  $B_{q_s}$  in place of  $B_{q_2}$ . This completes the proof of Proposition 2.

Now we can give proofs of Theorems 1 and 3. To prove Theorem 1, we may assume that  $1 < s \leq 2$ . Let  $\beta = 2^{q'}$  in Proposition 1. Then, since  $\theta$  is an arbitrary number in  $(0, 1)$ , we have Theorem 1 for  $s \in (1, 2]$ .

Next, take  $\beta = 2^{q's'}$  in Proposition 2. Then, we have

$$\|Tf\|_p \leq C(q-1)^{-1}(s-1)^{-1}\|\Omega\|_q\|h\|_{\Delta_s}\|f\|_p$$

for  $p \in (1, \infty)$ , since  $(1 + \theta, (1 + \theta)/\theta) \rightarrow (1, \infty)$  as  $\theta \rightarrow 0$ . From this the result for  $S$  in Theorem 3 follows if we take functions of the form  $f(x, z) = k(x)g(z)$ .

### 3. EXTRAPOLATION

We can prove Theorems 2 and 4 by an extrapolation method similar to the one used in [14]. We give a proof of Theorem 4 for the sake of completeness (Theorem 2 can be proved in the same way). We fix  $p \in (1, \infty)$  and  $f$  with  $\|f\|_p \leq 1$ . Let  $S$  be as in (1.2). We also write  $Sf = S_{h,\Omega}(f)$ . Put  $U(h, \Omega) = \|S_{h,\Omega}(f)\|_p$ . Then we see that

$$(3.1) \quad \begin{aligned} U(h, \Omega_1 + \Omega_2) &\leq U(h, \Omega_1) + U(h, \Omega_2), \\ U(h_1 + h_2, \Omega) &\leq U(h_1, \Omega) + U(h_2, \Omega), \end{aligned}$$

for appropriate functions  $\Omega, h, \Omega_1, \Omega_2, h_1$  and  $h_2$ . Set

$$\begin{aligned} E_1 &= \{r \in \mathbb{R}_+ : |h(r)| \leq 2\}, \\ E_m &= \{r \in \mathbb{R}_+ : 2^{m-1} < |h(r)| \leq 2^m\} \quad \text{for } m \geq 2. \end{aligned}$$

Then  $h = \sum_{m=1}^{\infty} h\chi_{E_m}$ . Put  $e_m = \sigma(F_m)$  for  $m \geq 1$ , where

$$\begin{aligned} F_m &= \{\theta \in \Sigma : 2^{m-1} < |\Omega(\theta)| \leq 2^m\} \quad \text{for } m \geq 2, \\ F_1 &= \{\theta \in \Sigma : |\Omega(\theta)| \leq 2\}. \end{aligned}$$

Let  $\Omega_m = \Omega\chi_{F_m} - \sigma(\Sigma)^{-1} \int_{F_m} \Omega d\sigma$ . Then  $\Omega = \sum_{m=1}^{\infty} \Omega_m$ . Note that  $\int_{\Sigma} \Omega_m d\sigma = 0$ . Applying Theorem 3, we see that

$$(3.2) \quad U(h\chi_{E_m}, \Omega_j) \leq C(q-1)^{-1}(s-1)^{-1}\|h\chi_{E_m}\|_{\Delta_s}\|\Omega_j\|_q$$

for all  $s, q \in (1, 2]$ .

Now we follow the extrapolation argument of A. Zygmund [18, Chap. XII, pp. 119–120]. For  $k \in \mathbb{Z}$ , put

$$\begin{aligned} E(k, m) &= \{r \in (2^k, 2^{k+1}] : 2^{m-1} < |h(r)| \leq 2^m\} \quad \text{for } m \geq 2, \\ E(k, 1) &= \{r \in (2^k, 2^{k+1}] : 0 < |h(r)| \leq 2\}. \end{aligned}$$

Then

$$\begin{aligned} \int_{E(k, m)} |h(r)|^{(m+1)/m} dr/r &\leq Cm^{-a} \int_{E(k, m)} |h(r)| (\log(2 + |h(r)|))^a dr/r \\ &\leq Cm^{-a} L_a(h), \end{aligned}$$

and hence

$$(3.3) \quad \|h\chi_{E_m}\|_{\Delta_{1+1/m}} \leq Cm^{-am/(m+1)} L_a(h)^{m/(m+1)}$$

for  $m \geq 1$ . Also we have

$$(3.4) \quad \|\Omega_j\|_{1+1/j} \leq C 2^j e_j^{j/(j+1)}.$$

From (3.1)–(3.4) we see that

$$\begin{aligned} U(h, \Omega) &\leq \sum_{m \geq 1} \sum_{j \geq 1} U(h \chi_{E_m}, \Omega_j) \leq C \sum_{m \geq 1} \sum_{j \geq 1} j m \|h \chi_{E_m}\|_{\Delta_{1+1/m}} \|\Omega_j\|_{1+1/j} \\ &\leq C(1 + L_a(h)) \sum_{m \geq 1} \sum_{j \geq 1} m^{1-am/(m+1)} j 2^j e_j^{j/(j+1)} \\ &= C(1 + L_a(h)) \left( \sum_{m \geq 1} m^{1-am/(m+1)} \right) \left( \sum_{j \geq 1} j 2^j e_j^{j/(j+1)} \right). \end{aligned}$$

When  $a > 2$ , it is easy to see that  $\sum_{m \geq 1} m^{1-am/(m+1)} < \infty$ . Also, we have

$$\begin{aligned} \sum_{j \geq 1} j 2^j e_j^{j/(j+1)} &= \sum_{e_j < 3^{-j}} + \sum_{e_j \geq 3^{-j}} \\ &\leq \sum_{j \geq 1} j 2^j 3^{-j^2/(j+1)} + \sum_{j \geq 1} j 2^j e_j 3^{j/(j+1)} \\ &\leq C + C \int_{\Sigma} |\Omega(\theta)| \log(2 + |\Omega(\theta)|) d\sigma(\theta). \end{aligned}$$

Collecting the results, we conclude the proof of Theorem 4.

*Remark.* For a positive number  $a$  and a function  $h$  on  $\mathbb{R}_+$ , let

$$N_a(h) = \sum_{m \geq 1} m^a 2^m d_m(h),$$

where  $d_m(h) = \sup_{k \in \mathbb{Z}} 2^{-k} |E(k, m)|$  ( $E(k, m)$  is as above). We define a class  $\mathcal{N}_a$  to be the space of all measurable functions  $h$  on  $\mathbb{R}_+$  which satisfy  $N_a(h) < \infty$ . Then, it can be shown that if  $h \in \mathcal{L}_a$  for some  $a > 2$ , then  $h \in \mathcal{N}_1$ . By a method similar to that used in this section, we can show the  $L^p$  boundedness of  $S$  in Theorem 4 under a less restrictive condition that  $h \in \mathcal{N}_1$  and  $\Omega \in L \log L$  (see [14]).

#### 4. AN ESTIMATE FOR A TRIGONOMETRIC INTEGRAL

Let  $A$  be an  $n \times n$  real matrix and

$$\phi_A(t) = (t - \gamma_1)^{m_1} (t - \gamma_2)^{m_2} \dots (t - \gamma_k)^{m_k}$$

be the minimal polynomial of  $A$ , where  $\gamma_i \neq \gamma_j$  if  $i \neq j$ . Let  $a_i(t) = (t - \gamma_i)^{m_i}$  for  $i = 1, 2, \dots, k$ . Then, we can find polynomials  $b_i(t)$  ( $i = 1, 2, \dots, k$ ) such that

$$\frac{1}{\phi_A(t)} = \sum_{i=1}^k \frac{b_i(t)}{a_i(t)}.$$

For each  $i$ ,  $1 \leq i \leq k$ , let  $P_i$  be the polynomial defined by

$$P_i(t) = \frac{b_i(t)}{a_i(t)} \phi_A(t).$$

We consider the  $n \times n$  matrices  $P_i(A)$ , which are defined as usual (see [8]).

Let

$$V_i = \{z \in \mathbb{C}^n : (A - \gamma_i E)^{m_i} z = 0\} \quad (i = 1, 2, \dots, k),$$

where  $E$  denotes the unit matrix. Then, the vector space  $\mathbb{C}^n$  can be decomposed into a direct sum as

$$\mathbb{C}^n = V_1 \oplus V_2 \oplus \cdots \oplus V_k.$$

Each of the matrices  $P_i(A)$  is the projection onto  $V_i$ ; indeed, we have the following (see [8]):  $P_i(A)z \in V_i$  for all  $z \in \mathbb{C}^n$ , for  $i = 1, 2, \dots, k$ , and

$$\begin{aligned} P_1(A) + P_2(A) + \cdots + P_k(A) &= E, \\ P_i^2(A) &= P_i(A), \quad P_i(A)P_j(A) = 0 \quad \text{if } i \neq j \quad (1 \leq i, j \leq k). \end{aligned}$$

For  $z = (z_i)$  and  $w = (w_i)$  in  $\mathbb{C}^n$ , we write  $\langle z, w \rangle = \sum_{i=1}^n z_i w_i$ . Let

$$(4.1) \quad J(A, \eta, \zeta) = \sum_{i=1}^k \sum_{j=0}^{m_i-1} | \langle (A - \gamma_i E)^j P_i(A) \eta, A^* \zeta \rangle |$$

for  $\eta, \zeta \in \mathbb{R}^n$ . In this section, we prove the following:

**Theorem 5.** *Let  $\eta, \zeta \in \mathbb{R}^n \setminus \{0\}$  and  $0 < a < b$ . Suppose that  $J(A, \eta, \zeta) \neq 0$  and the numbers  $a, b$  are in a fixed compact subinterval of  $(0, \infty)$ . Then, we have*

$$\left| \int_a^b \exp(i \langle t^A \eta, \zeta \rangle) dt \right| \leq C J(A, \eta, \zeta)^{-1/N},$$

where  $N = \deg \phi_A = m_1 + m_2 + \cdots + m_k$  and the constant  $C$  is independent of  $\eta, \zeta, a$  and  $b$ .

Since  $\sum_{i=1}^k P_i(A) = E$ , using the triangle inequality, we see that

$$|\langle \eta, A^* \zeta \rangle| \leq \sum_{i=1}^k |\langle P_i(A) \eta, A^* \zeta \rangle| \leq J(A, \eta, \zeta).$$

Therefore, Theorem 5 implies the following:

**Corollary.** *Let  $\eta, \zeta, a, b$  and  $N$  be as in Theorem 5. Then, we have*

$$\left| \int_a^b \exp(i \langle t^A \eta, \zeta \rangle) dt \right| \leq C |\langle A \eta, \zeta \rangle|^{-1/N}$$

when  $\langle A \eta, \zeta \rangle \neq 0$ .

This is used to prove Lemma 2 in Section 2.

We define the curve  $X(t) = t^A \eta$  for a fixed  $\eta \in \mathbb{R}^n \setminus \{0\}$ . Then, E. M. Stein and S. Wainger [17] proved the following (see [11, 16] for related results):

**Theorem A.** *Suppose that the curve  $X$  does not lie in an affine hyperplane. Then*

$$\left| \int_a^b \exp(i \langle X(t), \zeta \rangle) dt \right| \leq C |\zeta|^{-1/n},$$

where  $C$  is independent of  $\zeta \in \mathbb{R}^n \setminus \{0\}$ ; furthermore, if  $a$  and  $b$  are in a fixed compact subinterval of  $(0, \infty)$ , the constant  $C$  is also independent of  $a$  and  $b$ .

Now, we see that Theorem 5 implies Theorem A. Since  $P_i(A)z \in V_i$  ( $z \in \mathbb{C}^n$ ), we have  $(A - \gamma_i E)^m P_i(A) = 0$  if  $m \geq m_i$  ( $i = 1, 2, \dots, k$ ). Therefore

$$\begin{aligned} \exp((\log t)A)P_i(A) &= \exp((\log t)\gamma_i E) \exp((\log t)(A - \gamma_i E))P_i(A) \\ &= t^{\gamma_i} \sum_{j=0}^{m_i-1} \frac{(\log t)^j}{j!} (A - \gamma_i E)^j P_i(A). \end{aligned}$$

Thus, using  $\sum_{i=1}^k P_i(A) = E$ , we see that

$$(4.2) \quad t^A = \sum_{i=1}^k t^{\gamma_i} \left[ \sum_{j=0}^{m_i-1} \frac{(\log t)^j}{j!} (A - \gamma_i E)^j \right] P_i(A).$$

The assumption on  $X$  of Theorem A can be rephrased as follows: the function  $\psi(t) = \langle t^A \eta, \zeta \rangle$  is not a constant function on  $(0, \infty)$  for every  $\zeta \in \mathbb{R}^n \setminus \{0\}$ . If  $\psi(t)$  is not a constant function, then  $\psi'(t)$  is not identically 0. Thus, since  $t(d/dt)\psi(t) = \langle t^A \eta, A^* \zeta \rangle$ , by (4.2) we have  $J(A, \eta, \zeta) > 0$ , where  $J(A, \eta, \zeta)$  is as in (4.1). Let  $C_0 = \min_{|\zeta|=1} J(A, \eta, \zeta)$  and note that  $C_0 > 0$ . Then, from Theorem 5, it follows that

$$\left| \int_a^b \exp(i\langle X(t), \zeta \rangle) dt \right| \leq C C_0^{-1/N} |\zeta|^{-1/N}.$$

This implies Theorem A, since  $N \leq n$  (in fact, it is not difficult to see that  $N = n$  if  $X$  satisfies the assumption of Theorem A).

In the following, we give a proof of Theorem 5. Let  $I = [\alpha, \beta]$  be a compact interval in  $\mathbb{R}$ . Consider the differential equation

$$(4.3) \quad y^{(k)} + a_1 y^{(k-1)} + a_2 y^{(k-2)} + \dots + a_k y = 0 \quad \text{on } I,$$

where  $a_1, a_2, \dots, a_k$  are complex constants. Let  $\{\varphi_1, \varphi_2, \dots, \varphi_k\}$  be a basis for the space  $S$  of all solutions of (4.3). Then, we use the following to prove Theorem 5.

**Proposition 4.** *Let  $\varphi$  be a real valued function such that  $\varphi' \in S$ . Suppose that  $\varphi' = d_1 \varphi_1 + d_2 \varphi_2 + \dots + d_k \varphi_k$ , where  $d_1, d_2, \dots, d_k$  are complex constants, which are uniquely determined by  $\varphi'$ . Then, we have*

$$\left| \int_{\alpha}^{\beta} e^{i\varphi(t)} dt \right| \leq C (|d_1| + |d_2| + \dots + |d_k|)^{-1/k},$$

where  $C$  is independent of  $\varphi$ ; also the constant  $C$  is independent of  $\alpha, \beta$  if they are within a fixed finite interval of  $\mathbb{R}$ .

To prove Proposition 4 we use the following two lemmas. Both of them are well-known.

**Lemma 3.** *Let  $\varphi$  be a solution of (4.3). Suppose that  $\varphi$  is not identically 0. Then, there exists a positive integer  $K$  independent of  $\varphi$  such that  $\varphi$  has at most  $K$  zeros in  $I$ .*

**Lemma 4** (van der Corput). *Let  $f : [c, d] \rightarrow \mathbb{R}$  and  $f \in C^j([c, d])$  for some positive integer  $j$ , where  $[c, d]$  is an arbitrary compact interval in  $\mathbb{R}$ . Suppose that  $\inf_{u \in [c, d]} |(d/du)^j f(u)| \geq \lambda > 0$ . When  $j = 1$ , we further assume that  $f'$  is monotone on  $[c, d]$ . Then*

$$\left| \int_c^d e^{if(u)} du \right| \leq C_j \lambda^{-1/j},$$

where  $C_j$  is a positive constant depending only on  $j$ . (See [17, 18]).

We now give a proof of Proposition 4. We consider linear combinations  $c_1\varphi_1 + c_2\varphi_2 + \cdots + c_k\varphi_k$ , where  $c_1, c_2, \dots, c_k \in \mathbb{C}$ . We write  $\psi = c_1\varphi_1 + c_2\varphi_2 + \cdots + c_k\varphi_k$  and define

$$N_1(\psi) = |c_1| + |c_2| + \cdots + |c_k|,$$

$$N_2(\psi) = \min_{t \in I} \left( |\psi(t)| + |\psi'(t)| + \cdots + |\psi^{(k-1)}(t)| \right).$$

Let  $U = \{(c_1, c_2, \dots, c_k) \in \mathbb{C}^k : |c_1| + |c_2| + \cdots + |c_k| = 1\}$ . We consider a function  $F$  on  $I \times U$  defined by

$$F(t, c_1, c_2, \dots, c_k) = |\psi(t)| + |\psi'(t)| + \cdots + |\psi^{(k-1)}(t)|.$$

Then, the function  $F$  is continuous and positive on  $I \times U$  (see [4]). Thus, if we put

$$C_0 = \min_{(t, c_1, c_2, \dots, c_k) \in I \times U} F(t, c_1, c_2, \dots, c_k),$$

then we see that  $C_0 > 0$  and  $N_2(\psi) \geq C_0 N_1(\psi)$ .

Therefore, if  $\varphi$  is as in Proposition 4, we have

$$(4.4) \quad \min_{t \in I} \left( |\varphi'(t)| + |\varphi''(t)| + \cdots + |\varphi^{(k)}(t)| \right) \geq C_0 N_1(\varphi').$$

By (4.4), for any  $t \in I$ , there exists  $\ell \in \{1, 2, \dots, k\}$  such that

$$|(d/dt)^\ell \varphi(t)| \geq C N_1(\varphi'), \quad C > 0.$$

Applying Lemma 3 suitably, we can decompose  $I = \cup_{m=1}^H I_m$ , where  $H$  is a positive integer independent of  $\varphi$  and  $\{I_m\}$  is a family of non-overlapping subintervals of  $I$  such that for any interval  $I_m$  there is  $\ell_m \in \{1, 2, \dots, k\}$  satisfying  $|(d/dt)^{\ell_m} \varphi(t)| \geq |(d/dt)^j \varphi(t)|$  on  $I_m$  for all  $j \in \{1, 2, \dots, k\}$ , so  $|(d/dt)^{\ell_m} \varphi(t)| \geq C N_1(\varphi')$  on  $I_m$ , and such that  $\varphi'$  is monotone on each  $I_m$ . Therefore, by Lemma 4 we have

$$\left| \int_{\alpha}^{\beta} e^{i\varphi(t)} dt \right| = \left| \sum_{m=1}^H \int_{I_m} e^{i\varphi(t)} dt \right| \leq C \sum_{m=1}^H \min \left( |I_m|, N_1(\varphi')^{-1/\ell_m} \right)$$

$$\leq C N_1(\varphi')^{-1/k}.$$

Since  $N_1(\varphi') = |d_1| + |d_2| + \cdots + |d_k|$ , this completes the proof of Proposition 4.

Proof of Theorem 5. By the change of variables  $t = e^s$  and an integration by parts argument, we can see that to prove Theorem 5 it suffices to show

$$(4.5) \quad \left| \int_{\alpha}^{\beta} \exp(i \langle e^{tA} \eta, \zeta \rangle) dt \right| \leq C J(A, \eta, \zeta)^{-1/N}$$

for an appropriate constant  $C > 0$ , where  $[\alpha, \beta]$  is an arbitrary compact interval in  $\mathbb{R}$ . Let  $\psi(t) = \langle e^{tA} \eta, \zeta \rangle$ . Then,  $\psi'(t) = \langle e^{tA} \eta, A^* \zeta \rangle$ , and hence, by (4.2) we have

$$\psi'(t) = \sum_{i=1}^k \sum_{j=0}^{m_i-1} c_{ij}(\eta, \zeta) t^j e^{\gamma_i t},$$

where

$$c_{ij}(\eta, \zeta) = \frac{1}{j!} \langle (A - \gamma_i E)^j P_i(A) \eta, A^* \zeta \rangle.$$

It is known that  $N$  functions  $t^j e^{\gamma_i t}$  ( $0 \leq j \leq m_i - 1$ ,  $1 \leq i \leq k$ ) form a basis for the space of solutions for the ordinary differential equation of order  $N$  with

characteristic polynomial  $\phi_A$  (see [4]). Thus, the estimate (4.5) immediately follows from Proposition 4, since  $\sum_{i=1}^k \sum_{j=0}^{m_i-1} |c_{ij}(\eta, \zeta)| \approx J(A, \eta, \zeta)$ .

## REFERENCES

- [1] A. Al-Salman and Y. Pan, *Singular integrals with rough kernels in  $L \log L(S^{n-1})$* , J. London Math. Soc. (2) **66** (2002), 153–174.
- [2] A. P. Calderón and A. Torchinsky, *Parabolic maximal functions associated with a distribution*, Advances in Math. **16** (1975), 1–64.
- [3] A. P. Calderón and A. Zygmund, *On singular integrals*, Amer. J. Math. **78** (1956), 289–309.
- [4] E. A. Coddington, *An Introduction to Ordinary Differential Equations*, Prentice-Hall, Inc, Englewood Cliffs, N. J., 1961.
- [5] J. Duoandikoetxea and J. L. Rubio de Francia, *Maximal and singular integral operators via Fourier transform estimates*, Invent. Math. **84** (1986), 541–561.
- [6] D. Fan and Y. Pan, *Singular integral operators with rough kernels supported by subvarieties*, Amer. J. Math. **119** (1997), 799–839.
- [7] R. Fefferman, *A note on singular integrals*, Proc. Amer. Math. Soc. **74** (1979), 266–270.
- [8] S. Friedberg, A. Insel and L. Spence, *Linear Algebra*, Prentice-Hall, Inc, Englewood Cliffs, N. J., 1979.
- [9] S. Hofmann, *Weighted norm inequalities and vector valued inequalities for certain rough operators*, Indiana Univ. Math. J. **42** (1993), 1–14.
- [10] W. Kim, S. Wainger, J. Wright and S. Ziesler, *Singular integrals and maximal functions associated to surfaces of revolution*, Bull. London Math. Soc. **28** (1996), 291–296.
- [11] A. Nagel and S. Wainger,  *$L^2$  boundedness of Hilbert transforms along surfaces and convolution operators homogeneous with respect to a multiple parameter group*, Amer. J. Math. **99** (1977), 761–785.
- [12] J. Namazi, *On a singular integral*, Proc. Amer. Math. Soc. **96** (1986), 421–424.
- [13] N. Rivière, *Singular integrals and multiplier operators*, Ark. Mat. **9** (1971), 243–278.
- [14] S. Sato, *Estimates for singular integrals and extrapolation*, Studia Math. **192** (2009), 219–233.
- [15] E. M. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality and Oscillatory Integrals*, Princeton University Press Princeton, NJ, 1993.
- [16] E. M. Stein and S. Wainger, *The estimation of an integral arising in multiplier transformations*, Studia Math. **35** (1970), 101–104.
- [17] E. M. Stein and S. Wainger, *Problems in harmonic analysis related to curvature*, Bull. Amer. Math. Soc. **84** (1978), 1239–1295.
- [18] A. Zygmund, *Trigonometric Series*, 2nd ed., Cambridge Univ. Press, Cambridge, London, New York and Melbourne, 1977.

DEPARTMENT OF MATHEMATICS, FACULTY OF EDUCATION, KANAZAWA UNIVERSITY, KANAZAWA 920-1192, JAPAN

*E-mail address*: shuichi@kenroku.kanazawa-u.ac.jp