

# ウイルスおよびアルコールに起因する肝硬変からの 肝癌発癌機構の比較研究

著者	服部 信
著者別表示	Hattori Nobu
雑誌名	昭和62(1987)年度 科学研究費補助金 海外学術研究 研究成果報告書概要
巻	1987
ページ	2p.
発行年	1989-03-29
URL	<a href="http://doi.org/10.24517/00067901">http://doi.org/10.24517/00067901</a>



# 1987 Fiscal Year Final Research Report Summary

## Comparative studies on the mechanism of the development of liver cancer associated with viral and/or alcoholic liver cirrhosis

Research Project

### Project/Area Number

62041804

### Research Category

Grant-in-Aid for Overseas Scientific Research

### Allocation Type

Single-year Grants

### Section

Special Cancer Research

### Research Institution

Kanazawa University

### Principal Investigator

**HATTORI Nobu** Kanazawa University, Dept. of Medicine, professor, 医学部, 教授 (30019640)

### Co-Investigator(Kenkyū-buntansha)

クローディオ ティリベリ トリエステ大学, 医学部, 教授

レナルド ギアレリ トリエステ大学, 医学部, 教授

KOJIRO Masamichi Kurume University, Dept. of Medicine, professor, 医学部, 教授 (90080580)

OKUDA Kunio Chiba University, Dept. of Medicine, professor, 医学部, 教授 (40009519)

KOBAYASHI Kenichi Kanazawa University, Dept. of Medicine, Associate professor, 医学部, 助教授 (70019933)

TIRIBELLI Claudio Trieste University, Dept. of Medicine, professor

GIARELRI Leonard Trieste University, Dept. of Medicine, professor

### Project Period (FY)

1987

### Keywords

Alcoholic liver cirrhosis / Posthepatic liver cirrhosis / Liver cancer / Multinodular type / Nodules-in-nodules / Dysplasia / Corrected age / Mantel-Haenszel

### Research Abstract

Mainly based on autopsy materials in Trieste University including those in Bologna University, over half of the cases with liver cirrhosis were classified to micronodular alcoholic liver cirrhosis. The major type of liver cancer was multi-nodular type composed of several tumor nodules. There was no lesion such as nodules- in-nodules sometimes seen in Japan and there were few cases showed dysplasia in non-cancerous region. Approximately 70% of the cases with cirrhosis were posthepatic type of cirrhosis in Japan. Namely, there were several differences between the findings of liver cancer and underlying cirrhosis of both countries.

Epidemiologically there was a prominent difference between ages of the association of cancer in both countries, i.e., the peak age in Trieste was approximately 70 years old. In contrast the peak age in Japan was approximately 50 years old. This difference was significant using corrected age by the examination method of Mantel-Haenszel. Moreover, the cases cancer sometimes died of liver failure itself.

From these results, it is suggested that alcohol per se may have a weak role in producing liver cancer compared with hepatitis virus itself.

## Research Products (8 results)

	All	Other
	All	Publications (8 results)
[Publications] Okuda K, et al: Cancer Research. V.47 No.9. (1987)		▼
[Publications] Ohnishi K, Okuda K, et al: Hepatology. V.7 No.6. (1987)		▼
[Publications] Okuda, Ishak eds: "Neoplasms of the Liver" Springer—Verlag, Tokyo, (1987)		▼
[Publications] Arias et al eds.; Okuda K, et al: "The Liver Annual 6/1986, Hepatic tumors" Elsevier, Amsterdam, (1987)		▼
[Publications] Tygstrup N et al eds.; Okuda K et al: "Cirrhosis of the Liver, Hepatocarcinogenesis associated with liver cirrhosis" Elsevier, Amsterdam, (1987)		▼
[Publications] Okuda K: "Changing incidence of hepatocellular carcinoma in Japan" Cancer Research. 47. 4967-4972 (1987)		▼
[Publications] Ohnishi K: "Prognosis of hepatocellular carcinoma smaller than 5 cm in relation to treatment" 7. 1285-1290 (1987)		▼
[Publications] Arias A: Elsevier, Amsterdam. The Liver Annual 6/1986, 652 (1987)		▼

**URL:** [https://kaken.nii.ac.jp/report/KAKENHI-PROJECT-62041804/620418041987kenkyu\\_seika\\_hokoku\\_](https://kaken.nii.ac.jp/report/KAKENHI-PROJECT-62041804/620418041987kenkyu_seika_hokoku_)

Published: 1989-03-29