

Setting the criterion for fall risk screening for healthy community-dwelling elderly

著者	Demura Shinichi, Sato Susumu, Shin Sohee, Uchiyama Masanobu
journal or publication title	Archives of Gerontology and Geriatrics
volume	54
number	2
page range	370-373
year	2012-03-01
URL	http://hdl.handle.net/2297/27781

doi: 10.1016/j.archger.2011.04.010

1 Original Article

2

3 Title:

4 Setting the criterion for fall risk screening for healthy community-dwelling elderly

5

6 Running title: Screening for high fall risk among the elderly

7

8

9 Authors

10 Shinichi Demura Graduate school of Natural Science and Technology,
11 Kanazawa University, Kakuma, Kanazawa, Ishikawa,
12 920-1192, Japan

13 Susumu Sato Life-long Sports Core, Kanazawa Institute of Technology,
14 Ohgigaoka 7-1, Nonoichi, Ishikawa, 921-8501, Japan

15 Sohee Shin Center for innovation, Kanazawa University, Kakuma,
16 Kanazawa, Ishikawa, 920-1192, Japan

17 Masanobu Uchiyama Akita Prefectural University,
18 Akita, Akita, 010-0195, Japan

19

20 **Correspondence address:** Susumu Sato,

21 Life-long Sports Core, Kanazawa Institute of Technology,

22 Ohgigaoka 7-1, Nonoichi, Ishikawa, 921-8501, Japan.

23 Phone: +81 76-248-1100(ext.2386), Fax: +81 76-294-6704.

24 e-mail: sssato@neptune.kanazawa-it.ac.jp

25

26

1 Abstract

2 This study aimed to develop a criterion for screening high risk elderly using
3 Demura's fall risk assessment chart (DFRA), compared with the Tokyo metropolitan
4 Institute of gerontology fall risk assessment chart (TMIG). Participants included 1122
5 healthy elderly individuals aged 60 years and over (380 males and 742 females) 15.8%
6 of whom had experienced a fall. We assessed fall risk of the elderly by DFRA and TMIG.
7 To develop a criterion for screening high fall risk subjects among community-dwelling
8 elderly, receiver-operating-characteristic (ROC) analysis was conducted using fall
9 experience (separated into the categories of faller and non-faller) and the following fall
10 risk scale scores: 1) TMIG score, 2) DFRA score, and 3) potential for falling score
11 according to the DFRA (summing the scores of three items). In ROC analyses, the area
12 under the ROC curve (AUC) for evaluating the potential for falling gave a value of .797
13 (95%CI: .759 to .834) which proved better than the evaluation of the overall TMIG
14 (.654, 95%CI: .602 to .706) and DFRA scores (.680, 95%CI: .633 to .727). Assessment of
15 the potential for falling and fall experience are of benefit in screening for elderly
16 persons deemed to be at a high fall risk. Further examinations based on the
17 prospective data setting will be required.

18

19 Key-words: ROC analysis, cross-sectional study, prevention of falls, risk profiles

20

1 **Introduction**

2 Prevention of falls for the elderly is an extremely important social issue
3 (American geriatrics Society, 2001; Perell et al., 2001; Chan et al., 2006; Russell et al., 2009).
4 Various approaches to prevent these falls have been examined, one of which was fall
5 risk assessment. The main objective of fall risk assessment is to connect the outcomes
6 these assessments to prevent falls in the future. Thus, fall risk assessment should
7 provide information concerning the prediction of the possibility of falling in the future
8 and the determination of problems that lead to falls for individuals.

9 In the many cases, before a fall occurs, the “precursors” that a fall is about to
10 happen appear as a stumble, slip, stagger ect.. However, because the causes of a fall
11 are infinite in variety it is difficult to screen for high-fall risk subjects among the
12 elderly population using only a composite index which summarizes the assessments
13 regarding each fall risk factor. Furthermore, in the previous study it was reported that
14 there is a limitation in the ability to predict fall experiences from an overall score
15 consisting of several risk factors because of the diversity pattern of fall causes among
16 individuals (Demura and Sato, 2010b). It may be recommended that the possibility of
17 future falls (screening the high-fall risk elderly) be checked by the assessment of
18 potential for a fall, and, next, a risk profile assessment is conducted for multi-factorial
19 risk domains to determine problems that lead to falls for individuals. Based on these
20 processes, the prevention measures for falls can be developed for the individual.

21 Several fall risk assessments have been reported which have been based on
22 questionnaires and performance tests (Gates et al., 2008; Tiedemann et al., 2008;
23 Suzuki, 2000; Tinetti et al., 1988). Fall risk assessments that are questionnaire-based
24 are an inexpensive and simple method and are widely used for the general population.
25 In Japan, the fall risk assessment chart developed by the Tokyo Metropolitan Institute

1 of Gerontology (TMIG) is widely used for the community-dwelling elderly population
2 (Suzuki, 2000). However, it has been suggested that this chart is unclear with respect
3 to the selection process of the assessment items as well as the basis for criteria
4 calculation for the screening of high risk elderly. Furthermore, it is difficult to
5 determine a risk profile for specific individuals (Demura et al., 2010ab). Considering
6 these problems, we aim to develop a new fall risk assessment chart. We have examined
7 a selection of useful assessment items (Demura et al, 2010a), and have examined
8 useful risk factor to predict fall experience (Demura et al., 2010b). However, there is no
9 criterion for the screening of high fall risk elderly based on objective evidence.

10 This study aims to develop a criterion for screening high-risk elderly with
11 respect to Demura's fall risk assessment chart and, subsequently, to compare these
12 criteria with the TMIG fall risk chart.

13

14 **Method**

15 **Subjects and data collection**

16 The subjects participating in this study were healthy community-dwelling
17 elderly individuals aged 60 and over, living in the Akita, Kanagawa, Ishikawa, Fukui,
18 Nagano, Gifu, Aichi, Tottori and Fukuoka prefectures in Japan. Mail or field surveys
19 were sent to 1927 elderly subjects from which there were 1464 respondents. Among
20 these, 1122 elderly (70.3 +/- 7.1yr) showing missing values of less than 10 percent were
21 used for data analysis in this study. This pool of subject was composed of 380 males
22 (70.5 +/- 7.0 yr) and 742 females (70.4 +/- 7.2yr) with 177 of them (15.8%) having had a
23 fall experience in the last twelve months.

24

25 **Fall risk assessment**

1 Demura’s fall risk assessment chart (DFRA) is composed of previous fall
2 experience and 50 other fall risk assessment items representing the five risk factors
3 regarding the “potential for falling,” “physical function,” “disease and physical
4 symptoms,” “environment,” and “behavior and character” (Demura et al., 2010). The
5 “potential for falling” that a fall is currently happening and is a concept regarding the
6 occurrence of precursors that are related to falls, such as the act of stumbling. We
7 assessed the potential for falling by asking the patients to answer the following three
8 questions: “Have you often stumbled?” “In the past year, have you felt like you might
9 fall down?” and “Have you ever been told that you look like you might fall down?”
10 Physical function was assessed using 22 items selected from three categories
11 (fundamental function, advanced function, and gait) and eight elements (muscular
12 strength, lower limb strength, balancing ability, walking ability, going and down stairs,
13 changing and holding posture, upper limb function, and gait). Diseases and physical
14 symptoms were assessed using thirteen items selected from six categories (dizziness
15 and instances of blackout, medication, sight/hearing and cognitive disorder, cerebral
16 vascular, arthritic and bone disease, and circulatory disease). The environment was
17 assessed using four items selected from two categories (surrounding environment, and
18 clothing). The behavior and character was assessed using eight items selected from
19 four categories (inactivity, frequent urination, fear of falling, and risk behavior). All
20 questions were responded to on a dichotomous scale (yes or no), and with 1 point being
21 assigned to each response falling into the “high risk” category”.

22 In addition, we also used the TMIG fall risk assessment chart. The TMIG
23 assessment chart is composed of 15 items with each item assessed using a dichotomous
24 scale (yes or no). The subject with an overall score of 5 or higher or with fall experience
25 is considered to be at a high risk for a fall.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

Analyses

To develop a criterion for screening high fall risk subjects among the community-dwelling elderly, receiver-operating-characteristic (ROC) analysis was conducted using previous fall experience (faller or non-faller) and the followed fall risk scale scores; 1) TMIG score, 2) DFRA score, and 3) potential for falling score for the DFRA. We performed the ROC analysis on all of the trial models and determine the area under the ROC curve (AUC). Next, we calculated the positive likelihood ratio with a 95% confidence interval and set cut-off points in order to maximize the sensitivity and specificity for each score.

1) ROC analyses based on TMIG score

The TMIG score (TMIG-15) was calculated by summing all 15 items in the TMIG scale. As mentioned above, in the TMIG fall risk scale, a cut-off point for screening high fall risk subjects is recommended to be a score of 5 points without statistical procedures (Suzuki, 2000). To confirm the cut-off point of the TMIG for screening high fall risk person, we conducted ROC analysis using the TMIG-15 as a dependent variable.

The TMIG scale includes previous fall experience. However, we must use fall experience as a dependent variable in this study based on cross-sectional data. Therefore, we confirmed the accuracy of predictions made regarding the TMIG when excluding the influence of the previous fall experience. Thus, we calculated the TMIG score which summed over 14 TMIG item scores, excluding the “previous fall experience” (TMIG-14). Then the ROC analysis was conducted using the TMIIG-14 score as a dependent variable.

2) ROC analyses based on DFRA score

1 The DFRA score was calculated by summing over 50 fall risk item scores. This
2 study conducted ROC analysis using the DFRA score as a dependent variable.

3 3) ROC analyses based on the score of the potential for falling in the DFRA scale

4 The potential for falling in the DFRA scale was calculated by summing over
5 the scores for three items (PF-3). Next, ROC analyses were conducted using this score
6 to confirm the accuracy of predictions regarding these precursors. In our previous
7 study, we confirmed that the relationship between previous fall experiences and the
8 potential for falling score was comparable to those with overall DFRA score. If the
9 degree of fall risk in elderly subjects could be predicted from the score of potential for
10 falling, simplifying as well as improving fall risk screening.

11 Furthermore, for comparison with the TMIG scale, a similar ROC analysis was
12 also conducted using the scores of four items concerning previous fall experience
13 combined with the three potential for falling (PF-4).

14

15 **Results**

16 **1. ROC curve in TMIG**

17 In ROC analysis using the TMIG-14 score (excluding fall experience) (Figure
18 1a), the area under the curve (AUC) was .654 (95%CI: .602 to .706). A cut-off point was
19 set at 3 points and the sensitivity and specificity were .425 and .169, respectively.
20 Figure 1b shows the ROC curve using the TMIG-15 score (including fall experience).
21 The AUC, cut-off point, sensitivity and specificity were .786 (95%CI: .747-.825),
22 4-points, .594, and .831, respectively.

23

24 **2. ROC curve in DFRA**

25 In ROC analysis based on an overall score of DFRA (Figure 2), the AUC

1 was .680 (95%CI: .633 to .727). The cut-off point was set at 22 points, and the
2 sensitivity and specificity were .306 and .072.

3

4 **3. ROC curve in potential for falling DFRA score**

5 In the ROC analysis using the PF-3 score (Figure 3a), the AUC was .797
6 (95%CI: .759 to .834). The cut-off point was set at 1 point, and the sensitivity and
7 specificity were .869 and .657. When using the PF-4 score (including previous fall
8 experiences) (Figure 3b), the AUC was .946 (95%CI: .931 to .960). The cut-off point was
9 set at 2 points, and the sensitivity and specificity were .869 and .906. These results
10 show effectiveness of fall risk prediction using the potential for falling.

11

12 **Discussion**

13 This study examined a criterion for screening high fall risk elderly based on
14 the ROC analysis. The TMIG fall risk scale, which is widely used in Japan,
15 recommends a score of 5-points as a criterion for high fall risk in elderly persons.
16 However, there is no report regarding an objective basis for the calculation of this
17 criterion. In fact, in the examination of the validity of the criterion in the TMIG based
18 on our study sample, cut-off points for screening fallers (participants who had previous
19 experienced episodes of falling) was different from the recommended value. This result
20 indicates that the importance of this statistical demonstration in the development of a
21 criterion for screening.

22 Our previous study has reported that risk factor of the potential for falling are
23 closely related to previous fall experience, compared with other fall risk factors of
24 “physical function,” “disease and physical symptoms,” “environment,” and “character
25 and behavior” (Demura et al., 2010b). Therefore, we examined the screening of high

1 fall risk by potential for falling score, and proposed the criterion in this study.

2 In ROC analysis, the AUC evaluates the diagnostic accuracy of the test
3 because the area is equal to the provability of accurately discriminating between a
4 randomly chosen person with the outcome and a randomly chosen person without the
5 outcome (Eisenmann et al., 2010; Wray et al., 2010). It has been suggested that the
6 AUC be interpreted according to the following guidelines: non-informative/test equal
7 to chance ($AUC = 0.5$), less accurate ($0.5 < AUC < 0.7$), moderately accurate ($0.7 < AUC$
8 < 0.9), highly accurate ($0.9 < AUC < 1.0$), and perfect discriminatory test ($AUC = 1.0$)
9 (Swets, 1988; Eisenmann et al., 2010). An AUC of 0.8 has been stated to represent a
10 reasonably powerful model. In this study, the AUC for evaluating the potential for
11 falling score (three items) gave a value of 0.80 and it was better than for evaluating the
12 overall scores of the TMIG (15 items) and the DFRA (50 items). Furthermore, this
13 value was better than those reported in previous studies examining the validity of
14 performance tests for the screening of high fall risk (Muir et al., 2008). It indicates the
15 availability of screening by the potential for falling.

16 The potential for recurrent falls or multiple falls is high, and “previous fall
17 experience” is one of the important assessment items in a fall risk assessment
18 (American Geriatrics Society, 2001). Therefore, although this study examined cut-off
19 points using the potential for falling score, a fall risk assessment which takes into
20 account previous fall experience in the three items in the potential for falling may
21 prove effective in improving the accuracy of predicting future instances of falling.

22 On the other hand, the criterion proposed in this study has a limitation. Fall
23 risk is defined as the possibility of a fall occurring in the future. Therefore, essentially,
24 it is preferable that validity of a criterion for screening high fall risk is examined by
25 falls in the future based on the prospective study setting. However, because this study

1 is based on a cross-sectional data setting, we have to analyze our results using
2 previous fall experiences. In further examinations, the accuracy of predictions
3 regarding future instances of falling should be examined based on the prospective
4 study.

5 According to the results in this study, the assessment of the potential for
6 falling may be useful to screen high fall risk subjects, but it cannot propose
7 information concerning the specific risk profile for individuals. Comprehensive
8 assessment based on several risk factors is essential for taking measures to prevent
9 falls in the future. Fall risk assessment is not an end in itself, and the outcomes will be
10 incorporated into the prevention of falls. Therefore, it is very important to determine
11 problems for specific individuals in addition to comprehensive screening for patients
12 who are at a high risk for falling. The results of this study support that idea that the
13 potential for falling and previous fall experience provide useful information for the
14 screening of high fall risk subjects. However, we do not deny the significance of the
15 assessment of other risk factors. Further research will be required to develop an
16 assessment of the fall risk profile for individuals based on multiple risk factors.

17

18 **Summary**

19 This study examined a criterion for screening high fall risk elderly subjects
20 and proposed a cut-off point based on the potential for falling score. In addition, in
21 examinations based on our study sample, a cut-off point for screening using the TMIG
22 fall risk scale differed from the previously recommended cut-off value for screening
23 high fall risk elderly. Assessment of the potential for falling and previous fall
24 experience is beneficial for screening high fall risk elderly. In addition, further

1 research examining the accuracy of predictions regarding future instances of falling
2 will be required based on the prospective data setting.

3

4 **Acknowledgment**

5 This work was supported by A Grant-in-Aid for Science Research, the Japan Ministry of
6 Education, Science, Sports and Culture [grant number 21240064].


7

8

1 **References**


- 2 American Geriatrics Society, British Geriatrics Society, and American Academy of Orthopaedic
3 Surgeons Panel on falls prevention., 2001. Guideline for the prevention of falls in older
4 persons. *J. Am. Geriatr. Soc.* 49, 664-72.
- 5 Chan, B. K. S., Marshall, L. M., Winters, K. M., Faulkner, K. A., Schwartz, A.V., Orwoll, E.S.,
6 2006. Incident fall risk and physical activity and physical performance among older men. *Am.*
7 *J. Epidemiol.* 165, 696-703.
- 8 Demura, S., Sato, S., Yokoya, T., Sato, T., 2010a. Examination of useful items in the
9 assessment of fall risk in the community-dwelling elderly Japanese population.
10 *Environ. Health. Prev. Med.* 15, 169-179.
- 11 Demura, S., Sato, S., Yamaji, S., Kasuga, K., Nagasawa, Y., 2010b. Examination of
12 validity of fall risk assessment items for screening high fall risk elderly among the
13 healthy community-dwelling Japanese population. *Archives of Geriatric and*
14 *Gerontology*, in press [Epub a head of print].
- 15 Eisenmann, J. C., Laurson, K. R., DuBose, K. D., Smith, B. K., Donnelly, J. E., 2010.
16 Construct validity of a continuous metabolic syndrome score in children.
17 *Diabetology and Metabolic Syndrome*, 2-8,
18 <http://www.dmsjournal.com/content/2/1/8>.
- 19 Gates, S., Smith, L. A., Fisher, J. D., Lamb, S. E., 2008. Systematic review of accuracy
20 of screening instruments for predicting fall risk among independently living older
21 adults. *Journal of Rehabilitation Research and Development*, 45, 1105-1116.
- 22 Muir, S.W., Berg, K., Chesworth, B., Speechley, M., 2008. Use of the Berg balance scale
23 for predicting multiple falls in community-dwelling elderly people: a prospective
24 study. *Physical Therapy*. 88, 449-459.
- 25 Perell, K. L., Nelson, A., Goldman, R. L., Luther, S. L., Prieto-Lewis, N., Rubenstein, L.

- 1 Z., 2001. Fall risk assessment measures: An analytic review. *J. Gerontol.* 56,
2 M761-6.
- 3 Russell, M. A., Hill, K. D., Day, L. M., Blackberry, I., Gurrin, L. C., Dharmage, S. C.,
4 2009. Development of the falls risk for older people in the community (FROP-Com)
5 screening tool. *Age Ageing.* 38, 40-6.
- 6 Suzuki, T., 2000. Questionnaire for falls assessment of elderly people and its
7 application. *Health assessment manual.* Kosei Kagaku Kenkyusho, Tokyo, pp.
8 142-163 (In Japanese).
- 9 Swets, J, A., 1988. Measuring the accuracy of diagnostic systems. *Science*, 240,
10 1285-1293.
- 11 Tiedemann, A., Shimada, H., Sherrington, C., Murray, S., Lord, S., 2008. The
12 comparative ability of eight functional mobility tests for predicting falls in
13 community-dwelling older people. *Age and Ageing*, 37, 430-435.
- 14 Tinetti, M.E., Speechley, M., Ginter, S.F., 1988. Risk factors for falls among elderly
15 persons living in the community. *N. Engl. J. Med.* 319, 1701-07.
- 16 Wray, N. R., Yang, J., Goddard, M. E., Visscher, P. M., 2010. The genetic interpretation
17 of area under the ROC curve in genomic profiling. *PLoS Genet*, Feb 26,
18 6-2, :e1000864.
- 19



AUC	AUC (95%CI)	Sensitivity	Specificity
0.654	0.602- 0.706	0.425	0.169

Cut-off value
3



AUC	AUC (95%CI)	Sensitivity	Specificity
0.786	0.747-0.825	0.594	0.831

Cut-off value
4

Figure 1. The result of ROC analysis based on the TMIG score

Note) a: ROC curve when using the TMIG-14 score, b: ROC curve when using the TMIG-15 score