

Corrigendum to "characterization of the chemical diversity of glycosylated mycosporine-like amino acids in the terrestrial cyanobacterium Nostoc commune" [Journal of Photochemistry and Photobiology B: Biology 142 (2015) 154-168]

|       |   |
|-------|---|
| メタデータ | 言語: eng<br>出版者:<br>公開日: 2017-10-03<br>キーワード (Ja):<br>キーワード (En):<br>作成者:<br>メールアドレス:<br>所属: |
| URL   | <a href="https://doi.org/10.24517/00010770">https://doi.org/10.24517/00010770</a>           |

This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 International License.



1 **Table 4**  
 2 Summary of the NMR analysis of the 273-Da MAA.

| Position       | <sup>1</sup> H<br>(ppm) | Number<br>of H | Multiplet | <i>J</i> -value<br>(Hz) | <sup>13</sup> C<br>(ppm) | COSY<br>correlation | HMBC<br>correlation |
|----------------|-------------------------|----------------|-----------|-------------------------|--------------------------|---------------------|---------------------|
| 1              |                         |                |           |                         | 185.7                    |                     |                     |
| 2              |                         |                |           |                         | 131.1                    |                     |                     |
| 3              |                         |                |           |                         | 159.8                    |                     |                     |
| 4 <sup>a</sup> | 2.74                    | 1H             | d         | -17.4                   | 33.7                     | H4 <sup>b</sup>     | C3                  |
| 4 <sup>b</sup> | 2.86                    | 1H             | d         | -17.4                   |                          | H4 <sup>a</sup>     |                     |
| 5              |                         |                |           |                         | 73.1                     |                     |                     |
| 6 <sup>a</sup> | 2.36                    | 1H             | d         | -17.4                   | 44.0                     | H6 <sup>b</sup>     | C1, C5              |
| 6 <sup>b</sup> | 2.64                    | 1H             | d         | -17.4                   |                          | H6 <sup>a</sup>     |                     |
| 7              | 3.51                    | 2H             | s         |                         | 68.7                     |                     | C4, C5, C6          |
| 8              | 3.55                    | 3H             | s         |                         | 60.0                     |                     | C2                  |
| 9              | 3.39                    | 2H             | t         | 6.7                     | 43.0                     | H10                 | C3                  |
| 10             | 1.90                    | 2H             | tt        | 6.7, 7.1                | 26.2                     | H9, H11             | C9, C11, C12        |
| 11             | 2.44                    | 2H             | t         | 7.1                     | 32.1                     | H10                 | C9, C12             |
| 12             |                         |                |           |                         | 178.9                    |                     |                     |

3 NMR spectra were recorded with a JEOL ECS400 spectrometer in 30% methanol-d4 in D<sub>2</sub>O  
 4 as a solvent.

5 <sup>a</sup> Higher field signals

6 <sup>b</sup> Lower field signals

7