A cell-autonomous role for JSAP1 and JLP in mouse cerebellar Purkinje cell survival

メタデータ	言語: eng
	出版者:
	公開日: 2017-10-03
	キーワード (Ja):
	キーワード (En):
	作成者: 石川, 桃絵, 佐藤, 時春, 善岡, 克次
	メールアドレス:
	所属:
URL	https://doi.org/10.24517/00011108

This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 International License.



A cell-autonomous role for JSAP1 and JLP in mouse cerebellar Purkinje cell survival

Momoe ISHIKAWA^{1,*}, Tokiharu SATO¹, Hiroshi KIYONARI^{2,3}, and Katsuji YOSHIOKA^{1,*}

¹ Division of Molecular Cell Signaling, Cancer Research Institute, Kanazawa University, Kakuma-machi, Kanazawa 920–1192, Japan

² Animal Resource Development Unit and ³Genetic Engineering Team, RIKEN Center for Life Science Technologies,

2–2–3 Minatojima Minami-machi, Chuou-ku, Kobe 650–0047, Japan.

(Received November 26, 2015 and accepted in revised form December 21, 2015)

Abstract We previously reported that the structurally related JNK scaffolding proteins JSAP1 and JLP play functionally redundant and key roles in maintaining mouse cerebellar Purkinje cell (PC) survival, which may be related to their regulation of axonal transport. The JSAP1 and JLP proteins are also widely expressed in other cell types throughout the brain. Notably, in the cerebellum, JSAP1 is abundantly expressed in the pinceau, a cluster of basket cell (BC) axons and termini that surround the PC axon initial segment. Thus, it is possible that BC-expressed JSAP1 plays a role in PC survival. To investigate this possibility, we generated and analyzed mice containing a PC/BC-specific double knockout (DKO) of *Jsap1* and *Jlp (Jsap1:Jlp* cDKO mice). These mice exhibited PC axonal dystrophy, followed by progressive PC loss, consistent with the phenotypes previously reported for PC-specific *Jsap1* and *Jlp* DKO mice. Furthermore, we found that the phenotypes of *Jsap1:Jlp* cDKO PCs were rescued by PC-specific transgenic expression of JSAP1. Taken together, these results indicate that BC-expressed JSAP1 plays little or no role in PC survival, and that PC-expressed JSAP1 and JLP play cell-autonomous roles in preventing PC degeneration.

Keywords: Axonal swelling; Knockout mice; Neuronal degeneration

1 Introduction

The mitogen-activated protein kinase (MAPK) intracellular signaling pathway, consisting of MAPK, MAPK kinase (MAPKK), and MAPKK kinase, is well conserved from yeast to humans (Cano and Mahadevan, 1995; Herskowitz, 1995; Waskiewicz

^{*}Corresponding authors: momonomi@stu.kanazawa-u.ac.jp and katsuji@staff.kanazawa-u.ac.jp

and Cooper, 1995). Scaffolding proteins associated with MAPK modules contribute, at least in part, to the specificity and efficiency of these signaling cascades (Morrison and Davis, 2003; Yoshioka, 2004; Dhanasekaran et al., 2007). JNK/stress-activated protein kinase-associated protein 1 (JSAP1, also known as JIP3) and JNK-associated leucine zipper protein (JLP, also known as JIP4) are structurally related scaffolding proteins associated with mammalian JNK and p38 MAPK signaling modules (Ito et al., 1999; Kelkar et al., 2000; Lee et al., 2002; Kelkar et al, 2005) that can also interact with kinesin-1 light and heavy chains (KLC and KHC, respectively) through specific binding domains. JSAP1 and JLP are reported to function as adaptor proteins that link cell cargoes to kinesin-1 (Verhey et al., 2001; Nguyen et al., 2005), and regulate kinesin-1 activity and motility (Sun et al., 2011; Watt et al., 2015). Studies of JSAP1 loss-of-function mice indicate that JSAP1 is required during early development of the corpus callosum (Kelkar et al., 2003; Ha et al., 2005), and Jsap1 knockout (KO) mice die shortly after birth due to neuronal defects (Iwanaga et al., 2007). In contrast, Jlp KO mice are viable and grow normally (Iwanaga et al., 2008). Recently, however, JSAP1 and JLP were reported to have crucial and functionally redundant roles in the developing mouse brain and in the survival of Purkinje cells (PCs), both of which are associated with their regulation of kinesin-1-dependent axonal transport (Sato et al, 2015a and 2015b).

JSAP1 and JLP are widely expressed in the mouse brain. Notably, in the cerebellum, JSAP1 is highly expressed in the pinceau, a cluster of basket cell (BC) axons and termini that surround the PC axon initial segment (Miura et al., 2006). Thus, although we concluded from our previous study that PC-expressed JSAP1 and JLP play key, functionally redundant roles in PC survival (Sato et al, 2015b), it remains possible that BC-expressed JSAP1 and JLP are also involved in maintaining PC survival. To investigate this possibility, we generated and analyzed mice containing a PC/BC-specific double KO (DKO) in *Jsap1* and *Jlp* (*Jsap1:Jlp* cDKO), and then evaluated the effect of transgenic rescue with PC-expressed JSAP1.

2 Materials and Methods

2.1 Animals

All experimental procedures involving animals were approved by the Institutional Animal Care and Use Committees of Kanazawa University and RIKEN Kobe, respectively. *GluR* δ 2-*Cre* mice (Yamasaki et al., 2011), in which *Cre* is driven by *GluR* δ 2 promoter and thus is expressed specifically in BC and PC, were provided by Dr. Kenji Sakimura (Niigata University), and were crossed with *Jsap1^{f/f}* (Iwanaga et al., 2007) and *Jlp^{f/f}* (Sato et al., 2015a) mice. The green fluorescent protein (GFP) coding region in pBS-L7-GFP, a gift from Dr. Hirokazu Hirai (Gunma University), and a derivative of the pL7 Δ AUG vector containing the PS-specific *L*7 promoter but disrupting the translational start site (ATG) of the *L*7 gene (Oberdick et al, 1990; Tomomura et al., 2001), was replaced with the entire coding region of the mouse wild-type *Jsap1* gene. The *L*7-*Jsap1* transgene was then injected into fertilized eggs to generate *L*7-*Jsap1* transgenic mice (Accession No. CDB0513T: http://www2.clst.riken.jp/arg /TG%20mutant%20mice%20list.html). Genotyping of *L*7-*Jsap1* transgenic mice was performed using the following two primer sets:

5'-TGTTTGGAGGCACTTCTGACTTGC-3' and

5'-CTCGTCCATCTGGATCTCCATCAT-3' (primer set 1) and

5'-CGGAGAGGATGATGAAACTGAGGA-3' and

5'-CACTCAACTCTTTGTTGCTAGTGCC-3' (primer set 2) giving PCR products of 156-bp and 261-bp, respectively.

2.2 Antibodies (Abs)

The following primary Abs were used in this study: anti-JSAP1 (Miura et al, 2006), anti-calbindin D-28K (C9848, Sigma-Aldrich or AB1778, Millipore), anti-synapto-tagmin 1 (SYT1) (105002, Synaptic Systems), anti-SNAP25 (S5187, Sigma-Aldrich), anti-glial fibrillary acidic protein (GFAP) (G3893, Sigma-Aldrich), anti-phospho-JNK (9251, Cell Signaling), and anti- α -tubulin (T5168, Sigma-Aldrich) Abs. The following secondary Abs were used: Alexa 488- and 568-conjugated anti-mouse or anti-rabbit IgG (Invitrogen) and horseradish peroxidase (HRP)-conjugated anti-mouse or anti-rabbit IgG (GE Healthcare) Abs.

2.3 Western blotting (WB), immunohistochemistry (IHC), and PC quantification

WB, IHC, and PC quantification were performed as described previously (Sato et al., 2008 and 2015b). IHC images were acquired with a confocal laser scanning microscope (LSM510 META, Carl Zeiss) using a 20x or 40x objective lens.

2.4 Reverse transcription (RT)-polymerase chain reaction (PCR)

RT-PCR was carried out as previously described (Sato et al, 2004), using the following primer pairs: *L7-Jsap1*, 5'-CGTGTAACAGTTAATTCCCTGCCTG-3' (*L7*) and 5'-CTCGTCCATCTGGATCTCCATCAT-3' (*Jsap1*); *Gapdh*, 5'-AAATGGTGAAGGTCGGTGTG-3' and 5'-TGAAGGGGGTCGTTGATGG-3'.

2.5 Statistical analysis

Significance was determined using the two-tailed unpaired Student's *t-test*. P values < 0.05 were considered to be statistically significant.

3 Results and Discussion

3.1 PC/BC-specific deletions in *Jsap1* and *Jlp* cause PC axonal swelling and progressive cell loss

To investigate the roles of JSAP1 and JLP in BCs, we generated mice with PC/BCtargeted conditional deletions in *Jsap1*, *Jlp*, or both, using loxP-flanked (floxed) alleles of these genes (Iwanaga et al., 2007; Sato et al., 2015a), in combination with the *GluR* δ 2-*Cre* transgene (Yamasaki et al., 2011). We first performed IHC of the midsagittal brain sections of 24-week-old control (*Jsap1^{f/f}:Jlp^{f/f}*), single *Jsap1* or *Jlp* cKO (*Jsap1^{f/f}:Jlp^{f/+}:GluR* δ 2-*Cre* or *Jsap1^{f/+}:Jlp^{f/f}:GluR* δ 2-*Cre*), and *Jsap1:Jlp* cDKO (*Jsap1^{f/f}:Jlp^{f/f}:GluR* δ 2-*Cre*) mice using an Ab against calbindin, a PC marker. The results showed the presence of swollen PC axons associated with PC loss in the *Jsap1:Jlp* cDKO mice, but not in the single *Jsap1* or *Jlp* cKO mice (Fig. 1A). We quantified the PCs by Niss1 staining of the sections and found that the PC numbers in the cDKO mice were significantly reduced compared to those in the cKO or control mice (Fig. 1B). Taken together, these results indicated that JSAP1 and JLP play functionally redundant roles in PC survival, which may be related to the regulation of axonal transport in PCs.

Next, we used anti-calbindin Ab IHC to examine the PC axons of *Jsap1:Jlp* cDKO mice at different ages (4, 8, 16, and 24 weeks of age; Fig. 2A). Many swollen axons were observed in regions near the PC bodies of the cDKO mice at each of the ages examined. We also quantified the PCs in *Jsap1:Jlp* cDKO mice at those ages, and at 40 weeks of age. The PC numbers were comparable between the control and *Jsap1:Jlp* cDKO mice at 4 and 8 weeks of age (Fig. 2B). However, the cDKO mice at16 weeks of age showed significantly decreased PC numbers compared to control mice, and the degree of PC loss increased with age (Fig. 2B). In addition, intense immunosignals for GFAP, a hallmark of reactive gliosis, were detected in association with PC death in the *Jsap1:Jlp* cDKO mice at 24 weeks of age (Fig. 2C). These results are consistent with those previously obtained using PC-specific *Jsap1:Jlp* DKO mice (Sato et al., 2015b), suggesting that JSAP1 and JLP expressed in BCs have little or no effect on PC survival.

3.2 Selective accumulation of the kinesin-1 cargo SNAP25 in the swollen axons of JSAP1- and JLP-deficient PCs

Axonal dystrophy is thought to reflect defective axonal transport. Since JSAP1 and JLP are known to interact with kinesin-1, a motor protein involved in axonal transport, the axonal dystrophy phenotype of *Jsap1:Jlp* cDKO may be due to defects in kinesin-1-dependent axonal transport. To examine this possibility, we analyzed the swollen axons of PCs in 16-week-old *Jsap1:Jlp* cDKO mice by double-immunostaining, using





Figure 1. PC/BC-specific deletion of Jsap1 and Jlp causes PC axonal swelling and cell loss. (A) IHC analysis of $Jsap1^{f/f}$: $Jlp^{f/f}$ (control), single Jsap1 or Jlp cKO, and Jsap1:Jlp cDKO mice at 24 weeks of age. Midsagittal sections from the indicated mice were stained with an anti-calbindin Ab. PC axonal swelling and loss were seen in the Jsap1:Jlp cDKO mice, but not in the single Jsap1 or Jlp cKO mice. (B) PC quantification. Midsagittal sections from the indicated mice at 24 weeks of age were stained with Nissl, and the PCs per μ m in lobules II to IX were counted (n = 3 mice per group). PC numbers in the Jsap1:Jlp cDKO mice were significantly decreased compared to those in the control mice. Scale bar, 50 μ m. ***P < 0.001; n.s., not significant.

Abs against calbindin and either SNAP25, a kinesin-1 cargo (Hirokawa et al., 2009), or SYT1, a kinesin-3 cargo (Hirokawa et al., 2009). Strong SNAP25 immunosignals were detected throughout almost all of the swollen PC axons, whereas the SYT1 immunosignals were very low and limited in distribution (Fig. 3A). These results, together with the earlier finding of progressive PC loss in these mice, suggested that JSAP1 and JLP play important roles in the kinesin-1-dependent axonal transport in PCs, and that their ablation causes PC axonal dystrophy, leading to neuronal loss.

In addition, we analyzed JNK activation in the axons using a phospho-JNK Ab. The results showed substantial JNK activation in the swollen axons of PCs in the *Jsap1:Jlp* cDKO mice (Fig. 3B), consistent with that observed in the PC-specific *Jsap1:Jlp* DKO



Figure 2. PC axonal dystrophy, progressive PC loss, and reactive gliosis in *Jsap1:Jlp* cDKO mice. (A) Midsagittal sections of control and *Jsap1:Jlp* cDKO mice at 4, 8, 16, and 24 weeks of age were subjected to IHC with an anti-calbindin Ab. Higher magnifications of the boxed areas are shown in the right-most panels. (B) Mid-sagittal sections from control and *Jsap1:Jlp* cDKO mice at the indicated ages were stained with Nissl, and the PCs were counted as in Fig. 1B (n = 4 mice per group). PC numbers were significantly decreased in *Jsap1:Jlp* cDKO mice at 16 weeks of age, compared to control mice, and the degree of PC loss increased with age. (C) Sections of the PC layer in control and *Jsap1:Jlp* cDKO mice at 24 weeks of age were examined by IHC with an anti-GFAP Ab and 4,6-diamidino-2-phenylindole (DAPI). Scale bars, 50 μ m. ***P < 0.001; n.s., not significant.



mice (Sato et al., 2015b). At present, the precise mechanism underlying the PC death induced by the ablation of JSAP1 and JLP remains unknown. However, these results, together with the results of our previous study using *Jsap1:Jlp* DKO primary cultured neurons (Sato et al., 2015a), suggest that JNK activation may be involved in mediating the neuronal death.

bars, $50 \,\mu m$.

3.3 Transgenic expression of JSAP1 in PCs rescues the PC degeneration in *Jsap1:Jlp* cDKO mice

To examine whether the JSAP1 expressed in BCs affects PC survival, we generated transgenic mice expressing the PC *L*7 promoter-driven *Jsap1* transgene (*L*7-*Jsap1*, Fig. 4A), and performed rescue experiments by crossing *Jsap1:Jlp* cDKO mice with the transgenic mice. We first confirmed that the exogenous *Jsap1* mRNA was specifically expressed in the cerebellum, and not the cerebral cortex (Fig. 4B). We then analyzed the compound mice (*Jsap1:Jlp* cDKO/*L*7-*Jsap1*) by WB with an anti-JSAP1



Figure 4. Rescue of PC degeneration in Jsap1:Jlp cDKO mice by the PC-specific transgenic expression of JSAP1. (A) Schematic illustration of DNA construct used to generate transgenic mice that express L7 promoter-driven Jsap1. Arrows indicate primers for RT-PCR in B. (B) Total RNAs were extracted from the cerebellum (Ce) and cerebral cortex (Cx) of 8-week-old control and L7-Jsap1 transgenic mice, as indicated, and RT-PCR was performed using specific primers (indicated as arrows in A) for the L7-Jsap1 transgene, with (lanes 1-4), or without (lanes 5-8) reverse transcriptase (RT). Gapdh RT-PCR was used as a control. (C) WB analysis of JSAP1 in the cerebellum of the indicated mice at 8 weeks of age. α -tubulin was used as a loading control. (D) Midsagittal sections from the indicated mice at 8 weeks of age were examined by double-immunostaining for JSAP1 and calbindin. Asterisks and arrowheads denote JSAP1-labeled pinceaus and swollen PC axons, respectively. Higher magnifications of the boxed regions are shown in the panels at right. (E) Rescue of PC loss in *Jsap1:Jlp* cDKO mice by the transgenic expression of JSAP1. PC numbers in the indicated mice at 24 weeks of age were quantified as in Fig. 1B (n = 4 mice per group). Scale bars, 25 μ m. ***P < 0.001.

Ab (Fig. 4C), and by IHC using Abs against calbindin and JSAP1 (Fig. 4D). The expression levels of JSAP1 in the cerebellum were comparable between the control and *Jsap1:Jlp* cDKO/*L7-Jsap1* mice (Fig. 4C). JSAP1 immunosignals were detected in the pinceau of the control and transgenic mice, but not in that of the *Jsap1:Jlp* cDKO and *Jsap1:Jlp* cDKO/*L7-Jsap1* mice (Fig. 4D), indicating that the exogenous JSAP1 was expressed specifically in the PCs. Although swollen PC axons were observed in the *Jsap1:Jlp* cDKO mice at 8 weeks of age (Fig. 4D lower left panels, see also Fig. 2A), no axonal dystrophy was detected in the *Jsap1:Jlp* cDKO/*L7-Jsap1* mice at the same age (Fig. 4D lower right panels). Furthermore, the decreased number of PCs in the *Jsap1:Jlp* cDKO mice at 24 weeks of age was rescued in the *Jsap1:Jlp* cDKO/*L7-Jsap1* mice of the same age (Fig. 4E). Taken together, these results indicated that the JSAP1 and JLP play functionally redundant roles in PC survival.

In conclusion, our molecular genetic and immunohistological analyses highlighted the key roles of JSAP1 and JLP in mouse PC survival, and strongly suggested that JSAP1 and JLP function cell autonomously to prevent PC degeneration. At present, the role of JSAP1 in the pinceau remains unknown. Further studies are required to clarify this issue.

Acknowledgements

We thank Drs. Kenji Sakimura (Niigata University) and Hirokazu Hirai (Gunma University) for providing the $GluR\delta^2$ -Cre mice and pBS-L7-GFP vector, respectively.

References

- Cano E, Mahadevan LC. (1995) Parallel signal processing among mammalian MAPKs. *Trends Biochem Sci.* 20: 117–122.
- Dhanasekaran DN, Kashef K, Lee CM, Xu H, Reddy EP. (2007) Scaffold proteins of MAP-kinase modules. Oncogene. 26: 3185–3202.
- Ha HY, Cho IH, Lee KW, Lee KW, Song JY, Kim KS, Yu YM, Lee JK, Song JS, Yang SD, Shin HS, Han PL. (2005) The axon guidance defect of the telencephalic commissures of the JSAP1-deficient brain was partially rescued by the transgenic expression of JIP1. *Dev Biol.* 277: 184–199.
- Herskowitz I. (1995) MAP kinase pathways in yeast: for mating and more. Cell. 80: 187-197.
- Hirokawa N, Noda Y, Tanaka Y, Niwa S. (2009) Kinesin superfamily motor proteins and intracellular transport. Nat Rev Mol Cell Biol. 10: 682–696.
- Ito M, Yoshioka K, Akechi M, Yamashita S, Takamatsu N, Sugiyama K, Hibi M, Nakabeppu Y, Shiba T, Yamamoto KI. (1999) JSAP1, a novel Jun N-terminal protein kinase (JNK) that functions as scaffold factor in the JNK signaling pathway. *Mol Cell Biol.* 19: 7539–7548.
- Iwanaga A, Sato T, Sugihara K, Hirao A, Takakura N, Okamoto H, Asano M, Yoshioka K. (2007) Neuralspecific ablation of the scaffold protein JSAP1 in mice causes neonatal death. *Neurosci Lett.* 429: 43–48.
- Iwanaga A, Wang G, Gantulga D, Sato T, Baljinnyam T, Shimizu K, Takumi K, Hayashi M, Akashi T, Fuse H, Sugihara K, Asano M, Yoshioka K. (2008) Ablation of the scaffold protein JLP causes

60 A cell-autonomous role for JSAP1 and JLP in mouse cerebellar Purkinje cell survival

reduced fertility in male mice. Transgenic Res. 17: 1045-1058.

- Kelkar N, Gupta S, Dickens M, Davis RJ. (2000) Interaction of a mitogen-activated protein kinase signaling module with the neuronal protein JIP3. *Mol Cell Biol*. 20: 1030–1043.
- Kelkar N, Delmotte MH, Weston CR, Barrett T, Sheppard BJ, Flavell RA, Davis RJ. (2003) Morphogenesis of the telencephalic commissure requires scaffold protein JNK-interacting protein 3 (JIP3). Proc Natl Acad Sci USA. 100: 9843–9848.
- Kelkar N, Standen CL, Davis RJ. (2005) Role of the JIP4 scaffold protein in the regulation of mitogenactivated protein kinase signaling pathways. *Mol Cell Biol.* 25: 2733–2743.
- Lee CM, Onésime D, Reddy CD, Dhanasekaran N, Reddy EP. (2002) JLP: A scaffolding protein that tethers JNK/p38MAPK signaling modules and transcription factors. *Proc Natl Acad Sci USA*. 99: 14189–14194.
- Miura E, Fukaya M, Sato T, Sugihara K, Asano M, Yoshioka K, Watanabe M. (2006) Expression and distribution of JNK/SAPK-associated scaffold protein JSAP1 in developing and adult mouse brain. J Neurochem. 97: 1431–1446.
- Morrison DK, Davis RJ. (2003) Regulation of MAP kinase signaling modules by scaffold proteins in mammals. *Annu Rev Cell Dev Biol.* 19: 91–118.
- Nguyen Q, Lee CM, Le A, Reddy EP. (2005) JLP associates with kinesin light chain 1 through a novel leucine zipper-like domain. *J Biol Chem.* 280: 30185–30191.
- Oberdick J, Smeyne RJ, Mann JR, Zackson S, Morgan JI. (1990) A promoter that drives transgene expression in cerebellar Purkinje and retinal bipolar neurons. *Science*. 248: 223–226.
- Sato S, Ito M, Ito T, Yoshioka K. (2004) Scaffold protein JSAP1 is transported to growth cones of neurites independent of JNK signaling pathways in PC12h cells. *Gene.* 329: 51–60.
- Sato T, Torashima T, Sugihara K, Hirai H, Asano M, Yoshioka K. (2008) The scaffold protein JSAP1 regulates proliferation and differentiation of cerebellar granule cell precursors by modulating JNK signaling. *Mol Cell Neurosci.* 39: 569–578.
- Sato T, Ishikawa M, Mochizuki M, Ohta M, Ohkura M, Nakai J, Takamatsu N, Yoshioka K. (2015a) JSAP1/JIP3 and JLP regulate kinesin-1-dependent axonal transport to prevent neuronal degeneration. *Cell Death Differ*. 22: 1260–1274.
- Sato T, Ishikawa M, Yoshihara T, Nakazato R, Higashida H, Asano M, Yoshioka K. (2015b) Critical role of JSAP1 and JLP in axonal transport in the cerebellar Purkinje cells of mice. *FEBS Lett.* 589: 2805–2811.
- Sun F, Zhu C, Dixit R, Cavalli V. (2011) Sunday Driver/JIP3 binds kinesin heavy chain directly and enhances its motility. *EMBO J.* 30: 3416–3429.
- Tomomura M, Rice DS, Morgan JI, Yuzaki M. (2001) Purification of Purkinje cells by fluorescenceactivated cell sorting from transgenic mice that express green fluorescent protein. *Eur J Neurosci*. 14: 57–63.
- Verhey KJ, Meyer D, Deehan R, Blenis J, Schnapp BJ, Rapoport TA, Margolis B. (2001) Cargo of kinesin identified as JIP scaffolding proteins and associated signaling molecules. J Cell Biol. 152: 959–970.
- Waskiewicz AJ, Cooper JA. (1995) Mitogen and stress response pathways: MAP kinase cascades and phosphatase regulation in mammals and yeast. *Curr Opin Cell Biol.* 7: 798–805.
- Watt D, Dixit R, Cavalli V. (2015) JIP3 Activates Kinesin-1 Motility to Promote Axon Elongation. J Biol Chem. 290: 15512–15525.
- Yamasaki M, Miyazaki T, Azechi H, Abe M, Natsume R, Hagiwara T, Aiba A, Mishina M, Sakimura K, Watanabe M. (2011) Glutamate receptor $\delta 2$ is essential for input pathway-dependent regulation of synaptic AMPAR contents in cerebellar Purkinje cells. *J Neurosci*. 31: 3362–3374.
- Yoshioka K. (2004) Scaffold proteins in mammalian MAP kinase cascades. J Biochem. 135: 657-661.