

A Note on a Definition of (G)-convergence

メタデータ	言語: English 出版者: 公開日: 2017-10-03 キーワード: 作成者: Watanebe, Chikara, 渡辺, 力 メールアドレス: 所属:
URL	https://doi.org/10.24517/00011333

This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 International License.



A Note on a Definition of (G)-convergence

Chikara WATANABE

Department of Mathematics, Faculty of Science, Kanazawa University
 (Received 23 October 1969)

In a Lecture note given by G. Stolzenberg¹⁾, the definition of (G)-convergence is given as follows :

let X be a metric space and $\{S_i\}$ be a sequence of closed subsets of X , then it is said that the sequence $\{S_i\}$ converges to a closed subset S if for any compact set $K \subset X$, $\{S_i \cap K\}$ is a convergent sequence in $\text{Comp}(K)$ and $S = \text{Ulim}_{K \rightarrow \infty} (S_i \cap K)$.

Moreover it is denoted that if X is σ -compact, then a family of closed subsets of X is normal in the above sense. In our former papers²⁾, we used the above property for a family of analytic sets in a domain of C^1 . However, recently, M. Kita³⁾ pointed out that no convergent sequence of points is normal in the above definition. Therefore we amend the definition of (G)-convergence as follows.

Definition. Let (X, ρ) be a metric space and $\{S_i\}$ be a sequence of closed subsets of X . We say that $\{S_i\}$ converges geometrically to a closed subset S of X if

(i) for any point $p \in S$, there is a sequence $\{p_i\}$ of points such that $p_i \in S_i$ and $p_i \rightarrow p$.

(ii) for any compact set K and positive number ϵ , there is a positive integer $\nu_0 = \nu_0(K, \epsilon)$ such that $S_\nu \cap K \subset S^{(\epsilon)} \cap K$ for $\nu \geq \nu_0$, where $S^{(\epsilon)} = \bigcup_{q \in S} \{q' \in X; \rho(q, q') < \epsilon\}$.

Note that from this definition the following properties are obtained immediately.

1. If $S \cap K = \emptyset$, then $S_\nu \cap K = \emptyset$ for sufficiently large ν .
2. If a sequence $\{p_i\}$ of points $p_i \in S_i$ converges to $p \in X$, then $p \in S$.
3. If $\{S_i\}$ converges geometrically to S and T , then $S = T$.

LEMMA 1. *If X is σ -compact, then a family \mathfrak{C} of closed subsets is normal.*

-
- 1) Volumes, limits and extensions of analytic varieties, Lecture note in Math., No. 19, Springer Verlag (1966).
 - 2) On normarity of a family of analytic sets, Sci. Rep. Kanazawa Univ., **12** (1967), 209-213.
 On a family of pure-dimensional analytic sets, *ibid*, **13** (1968), 73-82.
 A remark on the theorem of Bishop, Proc. Japan Acad. **45** (1969), 243-246.
 - 3) Department of Mathematics, Faculty of Science, Tokyo University.

Proof. Let $\{K_\nu\}$ be a sequence of compact subsets of X such that $K_1 \subset K_2 \subset K_3 \dots$, and $\bigcup_{\nu=1}^{\infty} K_\nu = X$. Take any sequence $\{S_i\}$ of \mathcal{C} . Since $\text{Ccmp}(K_1)$ is compact metric space, there is a subsequence $\{S_i^{(1)}\}$ of $\{S_i\}$ such that $S_i^{(1)} \cap K_1$ converges to T_1 in $\text{Ccmp}(K_1)$. Also, since $\text{Comp}(K_2)$ is compact, there is a subsequence $\{S_i^{(2)}\}$ of $\{S_i^{(1)}\}$ such that $S_i^{(2)} \cap K_2$ converges to T_2 in $\text{Comp}(K_2)$. We continue this process. A diagonal sequence $\{S_i^{(j)}\}$ converges to T_j in $\text{Comp}(K_j)$ for any j . Let $T = \bigcup T_\nu$. From the property of the space $\text{Comp}(K)$, we have only to prove that T is closed. Evidently we may assume $T \neq \emptyset$. Let $\{p_j\}$ be a sequence of points in T such that $p_j \rightarrow p \in X$. Put $K = \{p, p_j, j=1, 2, \dots\}$, then for some positive integer j_0 , $K \subseteq \overset{\circ}{K}_{j_0}$. We shall show that $p_j \in T_{j_0}$ for any j . Let $p_j \notin T_{j_0}$. Since $p_j \in T_{j'}$ for some j' , there is a sequence of points $q_\nu^{(j)} \in S_\nu^{(j)}$ such that $q_\nu^{(j)} \rightarrow p_j$ ($\nu \rightarrow \infty$), and since $p \in \overset{\circ}{K}_{j_0}$, we may assume that $q_\nu^{(j)} \in K_{j_0}$.

Let $d = \rho(p_j, T_{j_0})$. Since $S_\nu^{(j)} \cap K_{j_0}$ converges to T_{j_0} , there is a positive integer ν_0 such that $S_\nu^{(j)} \cap K_{j_0} \subset T_{j_0}^{(d/2)} \cap K_{j_0}$ for $\nu \geq \nu_0$. Also we may assume that $\rho(q_\nu^{(j)}, p_j) < \frac{d}{2}$ for $\nu \geq \nu_0$. Since $q_\nu^{(j)} \in S_\nu^{(j)} \cap K_{j_0}$, $q_\nu^{(j)} \in T_{j_0}^{(d/2)}$ for $\nu \geq \nu_0$. On the other hand, $\rho(q_\nu^{(j)}, T_{j_0}) = \min_{t \in T_{j_0}} \rho(q_\nu^{(j)}, t) = \rho(q_\nu^{(j)}, t_0) \geq \rho(p_j, t_0) - \rho(q_\nu^{(j)}, p_j) > d - \frac{d}{2} = \frac{d}{2}$. This is a contradiction and $p_j \in T_{j_0}$ for any j . Since T_{j_0} is closed $p \in T_{j_0} \subset T$. Q.E.D.

LEMMA 2. Let $\{S_i\}$ be a sequence of purely k -dimensional analytic sets in a domain D of C^n . If $\{S_i\}$ converges analytically to a purely k -dimensional analytic set S in D , then $\{S_i\}$ converges geometrically to S .

Proof. This is a direct conclusion of an analytic convergence.

LEMMA 3. Let X be a metric space and $\{S_i\}$ be a sequence of closed subsets of X which converges geometrically to S . If there are positive constant N, M such that for any point $p \in S_i$, $H_d(S_i, rB(p:r)) \geq Nr^d$ where H_d is a d -dimensional Hausdorff measure and $B(p:r)$ is a relatively compact open ball of radius r with center p in X , and that $H_d(S_i) < M$ for all i , then for any compact set K , $H_d(S, rK) \leq M4^d/N$. Moreover if X is σ -compact, then $H_d(S) \leq M4^d/N$.

The proof of this Lemma is the same as that given in the Lecture note of G. Stolzenberg. From Lemma 3, the following Theorem of Bishop⁴⁾ holds.

THEOREM OF BISHOP. Let $\{S_i\}$ be a sequence of purely k -dimensional analytic sets in a domain D of C^n which converges geometrically to a non-empty closed set S . If the volumes of S_i are uniformly bounded, then S is also an analytic set in D .

4) Conditions for the analyticity of certain sets. Mich. Math. J., 11 (1964), 289-304.