

初島住彦* 日本産クロウメモドキ属の一新種

S. HATUSIMA* : A New *Rhamnus* from Japan


タイシヤククロウメモドキ (新称) これは朝鮮のマルバクロウメモドキ (*R. koraiensis* K. C. SCHNEIDER.) に最も近い

が葉はやや全縁状となり、葉の下面は全面でなく脈上のみに毛があり、葉縁の毛が著しいので区別できる。

また、キビノクロウメモドキにも関係深く清水氏はこれをキビノクロウメモドキの多毛の変種とした位である。しかし、キビノクロウメモドキは果梗が長さ 1~2 cm となり、葉縁の鋸歯が著しく、葉の下面の毛は少なくとも脈上以外にもあるので異なる。

Rhamnus chugokuensis
HATUSIMA, sp.
nov.

Rhamnus yoshinoi var. *velutina* T. Shimizu

Fig. 1. *Rhamnus chugokuensis* HATUSIMA (type)

* 琉球大学理工学部生物学教室 Faculty of Sci. & Engin., Ryukyu University

in Act. Phytotax. et Geobot. 18(1960) 163; Journ. Facult. Textile Sci. & Technol. Shinshu Univ. 36, ser. A. Biol. 12 (1963) 49 ?

Frutex ca. 2-3m. altus, inermis, ramuli oppositi, hornotini glabri, rubro-purpurascéntes leviter nitentes ca. 1.2mm. crassi, annotini 2.5mm crassi. Gemmae compressae ca. 2 mm longae, squamis paucis ciliatis obtectae. Stipulae lineares ca 2 mm longae margine ciliatae mox caducae. Folia opposita vel subopposita, decidua, matura crasse membranacea, oblongo-obovata raro oblanceolata, plerumque 4-5 cm longa, 1.5~2.0 cm lata, apice breviter acuminate vel obtusa saepe ad summum obtusa vel retusa, basi attenuata vel acuta, margine subintegra vel minutissime calloso-serrulata, nervis lateralibus utrinque 3-4 (5), supra tota pageine marginaque dense hirtella, subtus tantum ad nervos dense albo-hirtella et barbellata. Petioli 3-9 mm longi dense hirtelli. Fructus 1-4, axillares, ut videtur globosus purpureo-niger, ca 5 mm crassus, semina 2, brunnescens, obovato-elliptica, ca 4 mm longa et 2.5 mm lata, circa basin pore aperta. Pedicelli ca. 5 mm longi, glabri.

Type : Taisyaku-kyo, Pref. Hiroshima, S. HATUSIMA 26490, Oct. 19, 1931
(in Herb. Facult. Agric. Kagoshima Univ.)

This is most closely related to *R. koraiensis* SCHNEID. from which it differs by its nearly entire obovate-oblong leaves being hirtellous not on all over but only on nerves on the beneath. *R. yoshinoi* MAK. differs from the present species by its distinct crenulate-serrate leaves with acuminate apices and less denser indumentum on the beneath and its longer pedicels attaining about 1-2 cm long.
