

Chromosome numbers of 41 cultivated taxa of *Prunus* subg. *Cerasus* in Japan

メタデータ	言語: eng 出版者: 公開日: 2017-10-03 キーワード (Ja): キーワード (En): 作成者: メールアドレス: 所属:
URL	http://hdl.handle.net/2297/48615

Yoshikane Iwatsubo¹, Tetsuya Kawasaki² and Naohiro Naruhashi¹: Chromosome numbers of 41 cultivated taxa of *Prunus* subg. *Cerasus* in Japan

¹Department of Biology, Faculty of Science, Toyama University, Gofuku 3190, Toyama 930-8555, Japan;
²Takasago 4-16-5, Saitama City, Saitama 336-0011, Japan

Hundreds of cultivated taxa of flowering cherries are currently known in Japan (Nihon Hana no Kai 1982; Kawasaki 1994). Chromosome studies on the flowering cherries have been done by many workers (Ishikawa 1916; Okabe 1927, 1928; Sax 1931; Tanaka and Oginuma 1975; Oginuma and Tanaka 1976; Nishikawa 1985; Oginuma 1987 a,b, 1988; Somego 2000). Recently, the authors reported on the chromosome numbers of 193 taxa of flowering cherries in Japan (Iwatsubo et al. 2002). In some cultivated taxa, the chromosome counts presented by the authors, however, are different from the counts reported by previous researchers. Along with the taxa whose chromosome numbers are unknown, those taxa will require further cytological study, in understanding the cause of disagreement of the chromosome numbers. This second paper of a series, that deals with chromosome numbers of flowering cherries in Japan, comprises 41 taxa of *Prunus* subg. *Cerasus*.

Materials and methods

Meristems of leaf buds of 41 taxa of *Prunus* subg. *Cerasus* collected from the plants preserved in the Yuki Experimental Station of the Flower Association of Japan (Nihon Hana no Kai) were used in the study. After being collected, the leaf buds were readily pretreated in a 2 mM 8-hydroxyquinoline solution for a few hours at room temperature and kept at ca. 5°C for about 15 h, and then fixed and kept in Newcomer's fluid at room temperature until used. After immersion in 1 N HCl for a few hours, the leaf buds were macerated in 1 N HCl at 60°C for 10 minutes, and then washed in tap water. They

were stained in a drop of 1.5% lacto-propionic orcein on the slide glass and ordinary squash technique was applied in preparation.

Results and discussion

The taxa studied, the number of plants examined, and their code numbers in the Yuki Experimental Station of the Flower Association of Japan, along with the voucher specimens of some taxa kept in TNS, are presented in Table 1. The chromosome numbers determined in the study and reported previously are also listed in Table 1. All the taxa studied showed 2n=16 or 2n=24 chromosomes. As seen in Table 1, chromosome counts are reported here for the first time for 19 taxa: 17 are diploid taxa, and two are triploid taxa.

The present result supports our previous study (Iwatsubo et al. 2002) on the chromosome numbers of *P. ×introrsa* 'Introrsa' and *P. lannesiana* 'Sirayuki' being both triploid plants with 2n =24; *P. ×subhirtella* 'Autumnalis' and *P. ×subhirtella* 'Subhirtella' being both diploid plants with 2n=16. These plants were initially reported as follows: *P. ×introrsa* 'Introrsa' as a diploid plant with 2n=16 (Somego 2000), *P. lannesiana* 'Sirayuki' as a diploid plant with 2n=16 (Okabe 1927), *P. ×subhirtella* 'Autumnalis' as a triploid plant with 2n=24 (Oginuma and Tanaka 1976; Oginuma 1987), and *P. ×subhirtella* 'Subhirtella' as a triploid plant with 2n=24 (Oginuma and Tanaka 1976; Oginuma 1987). The disagreement in chromosome numbers of the four cultivars suggests that each is composed of two chromosome strains.

Table 1. List of studied taxa of *Prunus* subg. *Cerasus*, Japanese name, number of individuals examined, individual code number (voucher specimen number), and present and previous chromosome counts

Taxon	Japanese name	Present count (2n)	No. of plants examined	Individual code numbers in *YES report (*Voucher number)	Previous report	Reference
<i>P. apetala</i> (Siebold et Zucc.) Franch. et Sav.	Chojizakura (丁字桜)	16	1	434	16	Okabe 1927 (as <i>P. crassipes</i>) ; Oginuma and Tanaka 1976; Oginuma 1987
<i>P. incisa</i> Thunb.	Mame-zakura (豆桜)	16	1		16	Okabe 1927, 1928; Sax 1931; Oginuma and Tanaka 1976; Oginuma 1987
<i>P. incisa</i> Thunb. × <i>P. lannesiana</i> (Carrière) E.H.Wilson var. <i>speciosa</i> (Koidz.) Makino 'Umineko'	Umineko (海猫)	16	1	411 (Kawasaki 5257)	16	Iwatsubo et al. 2002
<i>P. ×introrsa</i> Yagi ex Ohwi 'Introrsa'	Tsubaki-kanzakura (椿寒桜)	24	2	70 (Kawasaki 5213), 83	16	Somego 2000
					24	Iwatsubo et al. 2002
<i>P. jamasakura</i> Siebold ex Koidz. 'Floridula'	Seigan-zakura (晝顛桜)	16	1	267		
'Kamatari-zakura'	Kamatari-zakura (鎌足桜)	16	1	9809		
'Sendaiya'	Sendaiya (仙台屋)	16	2	1318, 1558		
'Shizuka-noi'	Shizuka-noi (静香匂)	16	1	1560		
<i>P. ×kobukuzakura</i> Ohwi	Kobuku-zakura (子福桜)	16	1	86 (Kawasaki 841)	16	Iwatsubo et al. 2002
<i>P. lannesiana</i> (Carrière) E.H.Wilson 'Beni-tamanishiki'	Beni-tamanishiki (紅玉躑躅)	16	1	1474		
'Gosiozakura'	Gosho-zakura (五所桜)	16	1	9704		
'Iyo-kumagai'	Iyo-kumagai (伊予熊谷)	16	1	9705		
'Matsumae-hayazaki'	Matsumae-hayazaki (松前早崎)	24	1	939	24	Iwatsubo et al. 2002
'Matsumae-ohsio'	Matsumae-ohsio (松前大潮)	16	1	1691		
'Mitsukabizakura'	Mitsukabizakura (三ヶ日桜)	16	1	9624	16	Iwatsubo et al. 2002
'Ohsawazzakura'	Ōsawa-zakura (大沢桜)	16	1	9706		
'Senriko'	Senrikō (千里香)	24	1	738	24	Okabe 1927 (as <i>P. serrulata</i> f. <i>picta</i>)
'Shiogama'	Shiogama-zakura (鹽釜桜)	16	1	9521		
'Shizuka'	Shizuka (静香)	16	1	9536	16	Iwatsubo et al. 2002
'Sirayuki'	Shirayuki (白雪)	24	3	664, 655, 794	16	Okabe 1927 (as <i>P. serrulata</i> 'Shirayuki')
'Sirotae'	Sirotae (白妙)	24	1	699	24	Iwatsubo et al. 2002
					24	Okabe 1927 (as <i>P. serrulata</i> f. <i>albida</i>) ; Oginuma 1987; Iwatsubo et al. 2002

<i>Spiralis'</i>						
'Taihaku'						
'Wasinowo'						
	cv					
<i>P. speciosa</i> (Koidz.) Makino 'Akami-ohshima'						
<i>P. maximowiczii</i> Ruprecht						
<i>P. ×media</i> Miyoshi 'Media'						
<i>P. ×miyoshii</i> Ohwi 'Ambigua'						
<i>P. ×parvifolia</i> (Matsum.) Koehne 'Parvifolia'						
<i>P. sargentii</i> Rehder						
	cv					
<i>P. ×sasaga-oshidori</i> H.Kubota 'Sasaga-oshidori'						
<i>P. ×sieboldii</i> (Carrière) Wittm. 'Caespitosa'						
<i>P. ×subhirtella</i> Miq. 'Autumnalis'						
<i>P. verecunda</i> (Koidz.) Koehne 'Antiqua'						
<i>Prunus</i> sp.						
<i>Prunus</i> sp.						
<i>Prunus</i> sp.						
<i>Prunus</i> sp.						

*YES : Yuki Experimental Station of the Flower Association of Japan (Nihon Hana no Kai), Ibaraki Prefecture.

**Voucher specimens kept in TNS.

Acknowledgements

The authors are grateful to Mr. Seiji Saito (Yuki Experimental Station of the Flower Association of Japan) and Mr. Hideaki Tanaka (Flower and Green Institute of the Flower Association of Japan), who had provided help and encouragement during this work. We should like to thank the authority of the Yuki Experimental Station of the Flower Association of Japan for permission to collect plant materials from the reserves.

References

- Ishikawa, M. 1916. A list of the number of chromosomes. Bot. Mag. Tokyo **30**: 404–448.
- Iwatsubo, Y., Kawasaki, T. and Naruhashi, N. 2002. Chromosome numbers of 193 cultivated taxa of *Prunus* subg. *Cerasus* in Japan. J. Phytogeogr. Taxon. **50**: 21–34.
- Kawasaki, T. 1994. Nihon no Sakura (Flowering cherries of Japan). 383 pp. Yama-kei Publishers, Tokyo. (in Japanese)
- Nihon Hana no Kai. 1982. Sakura no hinshu ni kansuru chosakenkyu hokoku. 448 pp. Nihon Hana no Kai, Tokyo. (in Japanese)
- Nishikawa, T. 1985. Chromosome counts of flowering plants of Hokkaido (8). Journ. Hokkaido Univ. Educ., Sect. 2 B, **35**: 31–42. (in Japanese)
- Oginuma, K. 1987 a. Karyomorphological studies on *Prunus* in Japan. J. Sci. Hiroshima Univ., Ser. B, Div. 2, Bot. **21**: 1–66.
- Oginuma, K. 1987 b. Karyomorphological studies on some species of Japanese *Prunus*. Plant Chromosome Research 1987: 131–134.
- Oginuma, K. 1988. A cytological study of the tetraploid *Prunus pseudo-cerasus* ($2n=32$). La Kromosomo II–**51–52**: 1710–1714.
- Oginuma, K. and Tanaka, R. 1976. Karyomorphological studies on some cherry trees in Japan. J. Jpn. Bot. **51**: 104–109.
- Okabe, S. 1927. Cytological studies on *Prunus*. Bot. Mag. Tokyo **41**: 398–404. (in Japanese with English summary)
- Okabe, S. 1928. Zür Cytologie der Gattung *Prunus*. Sci. Repts. Tohoku Univ., Ser. IV, Biol. **3**: 733–743.
- Sax, K. 1931. The origin and relationships of the Pomoideae. J. Arn. Arb. **12**: 3–22.
- Somego, M. 2000. Cytogenetic studies of the *Prunus yedoensis* Matsum. (Somei-yoshino). Sakura Science **7**: 14–22. (in Japanese with English summary)
- Tanaka, R. and Oginuma, K. 1975. Karyological study in *Prunus* \times *yedoensis*. Chromosome Information Service **19**: 13–15.
- (Received November 7, 2003; accepted December 15, 2003)
- 岩坪美兼¹・川崎哲也²・鳴橋直弘¹：サクラ属 *Cerasus* 亜属 41 分類群の染色体数**
- サクラ属 *Cerasus* 亜属 41 分類群(野生種を含む)について、染色体数の観察を行った。観察の結果は以下のとおりである。これらのうち 19 分類群の染色体数は初めての報告である。
- 二倍体 ($2n=16$) のサクラ：
- アカミオオシマ, イヨクマガイ, ウズザクラ, ウミネコ, オオサワザクラ, オオヤマザクラ, カバザクラ, カマタリザクラ, クラマウズ, ゴショザクラ, コヒガン, コブクザクラ, サクナミキクザクラ, ササガオシドリ, シオガマザクラ, シズカ, シズカニオイ, ジュウガツザクラ, スマウラフゲンゾウ, セイガンザクラ, センダイヤ, タイザンフクン, チョウジザクラ, トウキヨウザクラ, ナラノヤエザクラ, ベニタマニシキ, マツマエオオシオ, マメザクラ, ミッカビザクラ, ミヤマザクラ。
- 三倍体 ($2n=24$) のサクラ：
- アイヅウスズミ, イトウザクラ, シラユキ, シロタエ, センリコウ, タイハク, タカサゴ, ツバキカンザクラ, フユザクラ, マツマエハヤザキ, ワシノオ。
- 以上の 41 分類群のなかで、シラユキとツバキカンザクラには二倍体が報告されており、コヒガンとジュウガツザクラには三倍体の報告がある。これらについては今回も調査を行ったが、前回 (Iwatsubo et al. 2002) と同様、シラユキとツバキカンザクラは三倍体、コヒガンとジュウガツザクラは二倍体であった。
- (¹〒930-8555 富山市五福3190 富山大学理学部生物学科; ²〒336-0011 さいたま市高砂4-16-5)