

Multiple-subgenotype infections of *Giardia intestinalis* detected in Palestinian clinical cases using a subcloning approach

| | |
|-------|---|
| メタデータ | 言語: eng 出版者: 公開日: 2017-10-03 キーワード (Ja): キーワード (En): 作成者: メールアドレス: 所属: |
| URL | http://hdl.handle.net/2297/18152 |

1 **Multiple-subgenotype infections of *Giardia intestinalis* detected in Palestinian**
2 **clinical cases using a subcloning approach**

3

4 Amjad I. A. Hussein^{a,b}, Tomohiro Yamaguchi^a, Kentaro Nakamoto^a, Motohiro Iseki^a,
5 Masaharu Tokoro^{a,*}

6 *^aDepartment of Parasitology, Graduate School of Medical Science, Kanazawa*
7 *University, Ishikawa 920-8640, Japan*

8 *^bChemical, Biological and Drug Analysis Center, An-Najah National University,*
9 *Nablus, West Bank, Palestine*

10

11 Abbreviations: 18S rRNA, 18S small-subunit ribosomal RNA; GDH, glutamate
12 dehydrogenase; SNPs, single-nucleotide polymorphisms; WHO, World Health
13 Organization; PBS, phosphate-buffered saline; PCR, Polymerase chain reaction; dNTP,
14 deoxynucleotide triphosphate; DMSO, dimethyl sulfoxide; DDBJ, DNA Data Bank of
15 Japan; NJ, neighbor-joining; EMBL, European Molecular Biology Laboratory

16

17 *Corresponding author. Tel.: +81-76-265-2821; fax: +81-76-234-4242. *E-mail address:*
18 *tokoro@med.kanazawa-u.ac.jp* (M. Tokoro)

19

1 **Abstract**

2 To evaluate the geographic distribution of *G. intestinalis* genotypes in Nablus,
3 West Bank, Palestine, a genotyping study was performed using clinical fecal samples.
4 Microscopic examination confirmed that 8 of 69 (11.6%) samples were *G. intestinalis*
5 positive, and subsequent genotyping analyses targeting the small-subunit ribosomal
6 RNA (18S rRNA) and glutamate dehydrogenase (GDH) genes revealed the *G.*
7 *intestinalis* genotypes within the 8 samples. Of these 8 samples, 6 were clustered with
8 assemblage A-II and the remaining 2 samples were clustered with assemblage B by 18S
9 rRNA gene analysis; however, direct sequencing of the GDH gene segments from the
10 latter 2 samples showed a mixed infection profile. To assess those samples, we
11 employed a subcloning approach and successfully isolated 6 independent assemblage B
12 subgenotypes. These partial GDH gene sequences (393 bp) had 15 single-nucleotide
13 polymorphisms, all of which were synonymous transition substitutions at the third
14 nucleotide position of codons. From the results, we concluded that the highly
15 polymorphic gene loci such as GDH gene locus might provide us an opportunity to
16 obtain a detailed molecular data even from the samples with multiple-subgenotype
17 mixed infections. Therefore, subcloning approach is recommended in genotyping
18 studies, especially in those conducted in giardiasis-endemic areas, where the repeated
19 and cumulative infections could be commonly expected.

20 *Key words:* *Giardia intestinalis*; Intraspecific diversity; Mixed infection; Subcloning;

1 Palestine

2 **1. Introduction**

3 *Giardia intestinalis* (syn. *G. duodenalis* and *G. lamblia*) inhabits the small
4 intestine, causing a wide range of symptoms and conditions, from asymptomatic to
5 severe diarrhea with or without malabsorption and weight loss. In developing countries,
6 the prevalence of giardiasis in patients with diarrhea has been reported to be
7 approximately 20% (range, 5%–43%) [1]. According to the World Health Organization
8 (WHO) estimates, the prevalence of symptomatic giardiasis is about 200 million cases
9 worldwide with some 500,000 new infections each year [2]. Giardiasis is thought to be
10 the most common community-derived disease leading to significant morbidity and
11 mortality worldwide [3,4]. Person-to-person, zoonotic, water-borne, and food-borne
12 transmissions can occur through the fecal-oral route after direct or indirect contact with
13 the infective-stage cysts of the organism [3,5,6].

14 The presence of morphologically indistinguishable characteristics such as host
15 specificity of *G. intestinalis* isolated from humans and various animals has led to the
16 advanced use of genetic markers to analyze diversity at the genomic DNA level [7-10].
17 These molecular studies have efficiently discriminated the assemblages and revealed the
18 complex genotypic structures within *G. intestinalis* populations [10,11]. At least 7
19 major assemblages, A–G [10], are considered valid in this species. The major genotypes
20 of *G. intestinalis* isolated from humans have been reported as assemblages A and B,

1 with 2 well-known subgenotypes in each, namely, subgenotypes A-I and A-II in
2 assemblage A and subgenotypes B-III and B-IV in assemblage B [9,10,12]. These
3 subgenotypes are used as references in many studies, and their detection in animal
4 sources suggests the alarming possibility of zoonotic transmission [11,13]. Furthermore,
5 detection of viable *G. intestinalis* cysts in fly has suggested that intense transmission
6 could also occur indirectly [14]. The remaining genetic assemblages (C, D, E, F, and G)
7 appear to be host-restricted to animals [4,10].

8 At present, the occurrence of mixed infections of *G. intestinalis* [15,16], the
9 role of different assemblages and subgenotypes [15], genetic diversity [17], frequency
10 of transmission [11,18], evolution [10], and clinical significance [19,20] remain
11 debatable in genotyping studies. To address these issues, molecular epidemiological
12 studies are required especially in endemic areas [4]; however, most molecular
13 epidemiological studies have been conducted in developed countries [11,12,21,22].

14 In the present study, we collected *G. intestinalis* samples from Nablus, West
15 Bank, Palestine, and performed subcloning analysis by using genomic DNA directly
16 extracted from human fecal specimens to evaluate the geographic distribution of the *G.*
17 *intestinalis* genotypes and to better understand the occurrence of mixed-genotype
18 infection in an endemic area.

19

20 **2. Materials and Methods**

1 2.1 Fecal sample collection and microscopic examination

2 Sixty-nine fecal samples were obtained from patients who sought medical
3 treatment for abdominal complaints during February and March 2006 in Nablus, West
4 Bank, Palestine. All the samples were preserved in 2.5% (w/v) potassium dichromate at
5 4°C and subsequently processed for cyst purification by using the sucrose centrifugal
6 flotation method as described previously [23]. The presence of pathogenic intestinal
7 protozoan parasites and the number of cysts were microscopically assessed with 600×
8 magnification using some of the purified samples. The cyst count was
9 semi-quantitatively showed as high (+++; multiple cysts in 1 view field), moderate (++;
10 single cyst in 1 view field), and low (+; single cyst in multiple view fields). The purified
11 cysts were stored at –20°C in phosphate-buffered saline (PBS; pH 7.2) until further
12 analysis. Each clinical fecal sample was collected after obtaining informed consent from
13 the patients and providing them with a unique identification number on site in order to
14 protect their personal information.

15

16 2.2 DNA extraction

17 After 3 cycles of freezing at –80°C and thawing at 95°C, genomic DNA was
18 extracted from the cysts in PBS solution by using the QIAamp DNA Mini Kit (Qiagen
19 Sciences, Maryland, USA) according to the manufacturer's instructions. The extracted
20 genomic DNA was concentrated by ethanol precipitation and preserved as an aqueous

1 solution at -20°C until use.

2

3 2.3 Polymerase chain reaction (PCR) targeting the 18S rRNA gene locus

4 A partial DNA fragment of the 18S rRNA gene of *G. intestinalis* was amplified
5 using previously described primers [10] (G18S2: 5'-TCCGGTYGATTCTGCC-3' and
6 G18S3: 5'-CTGGAATTACCGCGGCTGCT-3') in 0.2-ml thin-walled PCR tubes on
7 MyCycler (BioRad Laboratories, California, USA). PCR was carried out in a 20- μl
8 reaction mixture containing 1 μl of the extracted *Giardia* DNA template, 1 \times GC buffer I,
9 0.4 μM of each primer, 0.5 mM of each deoxynucleotide triphosphate (dNTP), and 1 U
10 of LA Taq polymerase (TaKaRa Bio Inc, Shiga, Japan), with 5% dimethyl sulfoxide
11 (DMSO; Wako Pure Chemical Industries, Osaka, Japan) as an additive. The cycling
12 parameters were as follows: initial denaturation at 94°C for 1 min; followed by 30
13 cycles at 94°C for 30 s, 60°C for 30 s, and 72°C for 50 s; and a final extension at 72°C
14 for 5 min. The PCR products were electrophoresed on 2% LO3 agarose (TaKaRa). The
15 purified products were stained with ethidium bromide and then visualized on a UV
16 transilluminator. The target fragments were then purified from the agarose gel by using
17 the Quantum PrepTM Freeze 'N Squeeze DNA Gel Extraction Spin Columns (BioRad
18 Laboratories) according to the manufacturer's instructions.

19

20 2.4 PCR targeting the GDH gene locus

1 A partial DNA fragment of the GDH gene of *G. intestinalis* was amplified using
2 previously described seminested PCR primers [24] (GDHeF:
3 5'-TCAACGTYAAYCGYGGYTTCCGT-3', GDHiF:
4 5'-CAGTACAACTCYGCTCTCGG-3', and GDHiR:
5 5'-GTTRTCCTTGACATCTCC-3') in 0.2-ml thin-walled PCR tubes on MyCycler
6 (BioRad Laboratories). Primary PCR was carried out in a 20- μ l reaction mixture
7 containing 1 μ l of the extracted *Giardia* DNA template, 1 \times KOD-Plus buffer, 0.4 μ M of
8 each primer, 0.2 mM of each dNTP, 1 mM of MgSO₄, and 1 U of KOD-Plus DNA
9 Polymerase (TOYOBO, Osaka, Japan), with 5% DMSO as an additive. The cycling
10 parameters were as follows: initial denaturation at 94°C for 30 s; followed by 30 cycles
11 at 94°C for 20 s, 63.4°C for 30 s, 72°C for 45 s; and a final extension at 72°C for 3 min.
12 The reaction mixture and cycle parameters for secondary PCR were the same as those
13 used for the primary PCR except for the following 2 steps: (1) the initial denaturation
14 step, which was performed for 30 s instead of 20 s and (2) the annealing step, which
15 was performed at 62°C instead of 63.4°C. Electrophoresis and visualization of the
16 amplified products and purification of the target fragments were performed following
17 the same procedures as used for the 18S rRNA gene locus.

18

19 2.5 DNA sequence analysis

20 The purified PCR products containing the 18S rRNA gene locus were directly

1 sequenced with appropriate primers in both directions on ABI Prism 310 Genetic
2 Analyzer by using the ABI Prism BigDye Terminator Cycle Sequencing Ready
3 Reaction Kit (PE Applied Biosystems, California, USA). The purified PCR products
4 containing the GDH gene locus were also sequenced directly as described above.
5 However, some of the PCR products, i.e., those containing the fragments of the GDH
6 gene isolated from 3 samples (PalH4, PalH6, and PalH8) were subcloned into the
7 *EcoRV* site of pBluescript II SK(+) (Stratagene, California, USA), and their full-length
8 sequences were determined using T3 and T7 primers with appropriate internal
9 sequencing primers. A subcloning strategy was adopted to determine the sequences
10 from the abovementioned samples, because mixed-genotype infection profiles were
11 observed in 2 samples (PalH4 and PalH8). The sample PalH6 was also analyzed by
12 subcloning as a control sample of non-mixed infection. At least 2 independent PCR
13 products were used in this process to confirm the sequencing results. DNA sequences
14 were analyzed using the DNAsis-Mac v3.6 (Hitachi, Yokohama, Japan).

15

16 2.6 Sequence alignment and phylogenetic analysis

17 All reference sequences of the 18S rRNA and GDH genes of *G. intestinalis*
18 used in this study were obtained from the DNA Data Bank of Japan (DDBJ) by using
19 the *blastn* algorithm (<http://blast.ddbj.nig.ac.jp/top-e.html>). Alignments and
20 phylogenetic analysis were performed by running the ClustalW v1.83 program on the

1 DDBJ homepage (<http://clustalw.ddbj.nig.ac.jp/top-e.html>). An unrooted
2 neighbor-joining (NJ) tree composed of the nucleotide sequences obtained in this study
3 and 17 sequences of the GDH gene from various assemblages of *G. intestinalis* was
4 reconstructed using TreeView v1.6.6 with *G. ardeae* as the outgroup. Branch lengths
5 and bootstrap values (1000 replicates) were derived from the NJ analysis.

6

7 2.7 Nucleotide sequence accession numbers

8 The partial sequences of the GDH gene of *G. intestinalis* reported in the
9 present study were submitted to the DDBJ/European Molecular Biology Laboratory
10 (EMBL)/GenBank nucleotide sequence databases under accession numbers from
11 AB295649 to AB295654.

12

13 3. Results

14 3.1 Microscopic examination and patient background

15 A summary of the microscopic examination results and detailed backgrounds
16 of the patients who excreted *Giardia* cysts in their fecal samples are shown in Table 1.
17 Of the 69 fecal samples collected, 8 were found to be cyst positive on direct
18 microscopic examination after purification by the sucrose centrifugal flotation method
19 (Table 1). The patients comprised 6 males, 1 female, and 1 person of unknown sex, all
20 in the age range of 1 to 36 years. All the patients complained of abdominal pain as the

1 main symptom; however, no cases of watery diarrhea were observed. The cyst
2 abundances in samples were revealed as a high cyst count (+++) in the PalH4 and
3 PalH8 samples, and a moderate (++) to low (+) cyst count in all the other samples.

4

5 3.2 18S rRNA gene analysis

6 18S rRNA gene segments (375 bp) were successfully amplified from all the
7 samples by PCR. All the 18S rRNA gene sequences obtained in this study showed
8 100% identity to the reference sequences (Table 2). Two identical sequences from the
9 PalH4 and PalH8 samples were homologous to a sequence of assemblage B
10 (DDBJ/GenBank/EMBL accession number, **AF199447**) [25], and all the sequences
11 from the other samples were identical to a sequence of assemblage A (**AF199446**) [25].
12 As observed in previous works [10,12], the results showed a comparatively low
13 resolution due to the conserved nature of the 18S rRNA gene sequence, and could not
14 be used to differentiate the sequences at the subgenotype level (data not shown).
15 However, all the 18S rRNA gene analysis results were consistent with the GDH gene
16 analysis results described below, and no discrepancy was observed in the determined
17 genotypes (Table 1).

18

19 3.3 GDH gene analysis

20 GDH gene segments (393 bp) were successfully amplified by PCR and their

1 full-length sequences were determined as described in “Materials and Methods” from
2 all the samples except PalH1 and PalH3. The GDH gene sequences from the PalH2,
3 PalH5, PalH6, and PalH7 samples showed 100% homology to a sequence of
4 assemblage A-II (**L40510**, Table 2). The PalH4 and PalH8 samples, whose sequences
5 were determined using the subcloning technique as described in “Materials and
6 Methods,” revealed complex mixed subgenotypes. Each sample contained at least 3
7 independent isolates (PalH4: PalH4-1, PalH4-2, and PalH4-3; PalH8: PalH8-1, PalH8-2,
8 and PalH8-3), which were all clustered into assemblage B (Table 2, Fig. 1). There was
9 no overlapping of subgenotype sequences among those samples. While the sequences of
10 6 clones of the control PalH6 sample showed complete identity to the direct sequencing
11 result mentioned above, thus the possibility of PCR mutagenesis was denied in this
12 methodology.

13

14 3.4 Analyses of the samples with mixed subgenotypes

15 Of the 34 clones isolated from the PalH4 sample, 44.1%, 20.6%, and 35.3%
16 clones showed the PalH4-1, PalH4-2, and PalH4-3 subgenotypes, respectively, while of
17 the 16 clones isolated from the PalH8 sample, 31.3%, 37.4%, and 31.3% clones showed
18 the PalH8-1, PalH8-2, and PalH8-3 subgenotypes, respectively. A
19 DDBJ/GenBank/EMBL database search revealed 2 homologues; the sequence of
20 subgenotype PalH4-3 was identical to the unpublished sequence gi-hum1 (**DQ840541**)

1 reported from Poland and that of subgenotype PalH8-3 was identical to GH-135
2 (**AB195224**) reported from Japan. The sequences of none of the other subgenotypes
3 showed homology with any sequence in the database (Table 2, Fig. 1). Alignment of all
4 the 6 confirmed nucleotide sequences of the clones revealed 12 single-nucleotide
5 polymorphisms (SNPs) within the PalH4 subgenotypes and 10 SNPs within the PalH8
6 subgenotypes. As a result, a total of 15 SNPs were confirmed among the 6
7 subgenotypes (Table 3). Translation of all the GDH gene sequences of the subgenotypes
8 revealed a conserved characteristic; the sequences of all the assemblage B subgenotypes
9 were translated to a certain conserved amino acid sequence, since all the substitutions
10 were silent and positioned at the third nucleotide of codons (Table 3).

11 Phylogenetically (Fig. 1), the GDH gene sequences of the subgenotypes of the
12 PalH2, PalH5, PalH6, and PalH7 samples, all of which were completely homologous
13 with a reference sequence of assemblage A-II (Ad-2), formed a cluster together with the
14 reference sequence of assemblage A-I (Ad-1) with a high bootstrap value (99.9%). This
15 cluster was clearly differentiated from an assemblage B cluster, which included all the
16 mixed-subgenotype isolates and reference sequences from various organisms. Although
17 the assemblage B cluster seemed widespread and highly divergent, the local bootstrap
18 value (100%) was statistically significant and supported the formation.

19

20 **4. Discussion**

1 In the present study, we identified 2 cases of intra-assemblage mixed infections
2 in a single fecal sample. The mixed infection may be because the patients had been
3 repeatedly and cumulatively exposed to the pathogen due to poor sanitation in the study
4 areas in Nablus, West bank, Palestine (Table 1). The only previous report on intestinal
5 protozoan parasites in the area [26] showed the prevalence of intestinal parasites such as
6 *Entamoeba histolytica* (22.9%), *G. intestinalis* (7.3%), and *Ascaris lumbricoides* (5.7%),
7 thus supporting our speculation. In addition, the fact that a case of mixed infection is
8 rarely reported in developed countries [12,17-19,27,28].

9 In contrast to a comparatively high prevalence of inter-assemblage mixed
10 infections, which were detected using common methods such as restriction fragment
11 length polymorphism [12,18,24,27,29] or PCR with assemblage-specific primer sets
12 [8,12,24,30], the prevalence of intra-assemblage mixed infections, which were detected
13 by their mixed nucleotide sequence profiles obtained by direct sequencing, has been
14 rare [15,16,18]. Considering these facts together with the subcloning results obtained in
15 this study, it appears that the incidence of intra-assemblage mixed infections is
16 underestimated. Lalle and colleagues [18] stated that they reported only
17 inter-assemblage mixed infections because of the technical difficulties encountered in
18 the determination of intra-assemblage mixed infections. In other words, previous studies
19 apparently failed to detect intra-assemblage mixed infections possibly due to the
20 isolation process-related selection bias to in vitro culture of *G. intestinalis* and also due

1 to the technical limitations of the above mentioned conventional methods. However, we
2 analyzed the genomic DNA directly extracted from the fecal specimens and revealed
3 the original DNA population structures in the samples by using a subcloning technique.

4 In addition, as shown by our results, 18S rRNA gene analysis could not detect
5 the mixed-infection profiles, suggesting that highly polymorphic target gene loci such
6 as the GDH gene locus are needed for precise evaluation of mixed infections. It is also
7 suggested that the high levels of diversity observed within assemblage B could provide
8 a suitable target for evaluating intra-assemblage mixed infections, while assemblage A
9 does not seem to be suitable for the purpose because of its conservative features.

10 Similar to *Giardia* genotyping reports from other geographic areas, e.g., USA
11 [17], UK [27], Netherlands [12,19], Italy [18,28], India [16,17], China [30], Mexico
12 [31,32], and Australia [9,10,33], the samples from Palestine also showed only
13 assemblage A and B and none of the remaining genotypes (C–G), which have mainly
14 been reported in animals [4,10,16]. All the determined sequences of assemblage A
15 samples from Palestine were completely identical to the sequence of reference
16 assemblage A-II, which has been reported worldwide, e.g., in UK [27], Italy [28], India
17 [16], Mexico [31,32], and Australia [10].

18 Although the phylogenetic analysis of the GDH gene sequences indicated a
19 high degree of genetic polymorphism within the assemblage B cluster, alignment
20 analysis of the polymorphisms within the GDH gene locus revealed that all nucleotide

1 substitutions observed among the assemblage B subgenotypes were positioned at the
2 third nucleotide of the codons and were transition mutations, i.e., they were
3 purine-purine or pyrimidine-pyrimidine substitutions (Table 3). Due to the restricted
4 substitutions, all the mutations on the nucleotide level were synonymous, resulting in no
5 change in the deduced amino acid sequence. In other words, all the nucleotide
6 substitutions converged on 1 amino acid sequence (Table 3). On the other hand, in
7 assemblage A subgenotypes, all the reference sequences including those of assemblage
8 A-II subgenotypes of the GDH gene appeared to converge on an another amino acid
9 sequence (data not shown).

10 These findings reasonably suggest that genetic diversity in *G. intestinalis* could
11 be analyzed at 2 independent levels: (1) inter-assemblage amino acid diversity, which
12 appears completely conserved within each assemblage and may reflect unrevealed
13 ancient divergence and (2) intra-assemblage nucleotide diversity, which may reflect an
14 ongoing process of nucleotide mutations under the restriction of the former divergence.
15 Although the driving force and maintenance mechanism of these 2 levels of diversity
16 remain unclear, the concept could contribute to a better understanding of the evolution
17 of *G. intestinalis*.

18 In conclusion, the concept of 2 levels of diversity could be useful for
19 elucidating the evolutionary patterns and present population structure of *G. intestinalis*.
20 Furthermore, the comparatively high prevalence of intra-assemblage mixed infections

1 of *G. intestinalis* observed in this study indicates the common occurrence of repeated
2 and cumulative infections by the pathogen, especially in endemic areas. Therefore, a
3 subcloning approach targeting highly polymorphic gene loci is recommended to obtain
4 precise and detailed molecular epidemiological data, which could directly contribute to
5 a better understanding of the intraspecific diversity of this unique pathogen.

6

7 **Acknowledgements**

8

9 This work was supported by a Grant-in-Aid for Scientific Research to MT
10 (18406007, 20406007) from the Ministry of Education, Culture, Sports, Science and
11 Technology of Japan and a grant for research on emerging and reemerging infectious
12 diseases to MT (17211101) from the Ministry of Health, Labour and Welfare of Japan.

13

14 **References**

15 [1] Rabbani GH, Islam A. Giardiasis in humans: populations most at risk and prospects
16 for control. *Giardia intestinalis: from molecules to disease*. CAB International,
17 Wallingford, UK. : Thompson RCA, Reynoldson JA, and Lymbery AJ (ed.),
18 1994:217-49.

19 [2] World Health Organization. Infections and parasitic diseases. Section in Global
20 health situation and projections-estimates, WHO, Switzerland, 1992:45-50.

- 1 [3] Caccio SM, Thompson RCA, McLaughlin J, Smith HV. Unravelling
2 *Cryptosporidium* and *Giardia* epidemiology. Trends Parasitol 2005;21:430-37.
- 3 [4] Thompson RCA, Monis PT. Variation in *Giardia*: implications for taxonomy and
4 epidemiology. Adv Parasitol 2004;58:69-137.
- 5 [5] Hunter PR, Thompson RCA. The zoonotic transmission of *Giardia* and
6 *Cryptosporidium*. Int J Parasitol 2005;35:1181-90.
- 7 [6] Smith HV, Caccio SM, Tait A, McLaughlin J, Thompson RCA. Tools for
8 investigating the environmental transmission of *Cryptosporidium* and *Giardia*
9 infections in humans. Trends Parasitol 2006;22:160-67.
- 10 [7] Bertrand I, Albertini L, Schwartzbrod J. Comparison of two target genes for
11 detection and genotyping of *Giardia lamblia* in human feces by PCR and
12 PCR-restriction fragment length polymorphism. J Clin Microbiol 2005;43:5940-5944.
- 13 [8] Guy RA, Xiao C, Horgen PA. Real-time PCR assay for detection and genotype
14 differentiation of *Giardia lamblia* in stool specimens. J Clin Microbiol
15 2004;42:3317-20.
- 16 [9] Hopkins RM, Meloni BP, Groth DM, Wetherall JD, Reynoldson JA, Thompson
17 RCA. Ribosomal RNA sequencing reveals differences between the genotypes of
18 *Giardia* isolates recovered from humans and dogs living in the same locality. J Parasitol
19 1997;83:44-51.
- 20 [10] Monis PT, Andrews RH, Mayrhofer G, Ey PL. Molecular systematics of the

1 parasitic protozoan *Giardia intestinalis*. Mol Biol Evol 1999;16:1135-44.

2 [11] van Keulen H, Macechko PT, Wade S, Schaaf S, Wallis PM, Erlandsen SL.

3 Presence of human *Giardia* in domestic, farm and wild animals, and environmental

4 samples suggests a zoonotic potential for giardiasis. Vet Parasitol 2002;108:97-107.

5 [12] van der Giessen JWB, de Vries A, Roos M, Wielinga P, Kortbeek LM, Mank TG.

6 Genotyping of *Giardia* in Dutch patients and animals: a phylogenetic analysis of human

7 and animal isolates. Int J Parasitol 2006;36:849-58.

8 [13] Aloisio F, Filippini G, Antenucci P, Lepri E, Pezzotti G, Caccio SM, Pozio E.

9 Severe weight loss in lambs infected with *Giardia duodenalis* assemblage B. Vet

10 Parasitol 2006;142:154-58.

11 [14] Szostakowska B, Kruminis-Lozowska W, Racewicz M, Knight R, Tamang L,

12 Myjak P, Graczyk TK. *Cryptosporidium parvum* and *Giardia lamblia* recovered from

13 feral filth flies. Appl Environ Microbiol 2004;70:3742-44.

14 [15] Robertson LJ, Hermansen L, Gjerde BK, Strand E, Alvsvag JO, Langeland N.

15 Application of genotyping during an extensive outbreak of waterborne giardiasis in

16 Bergen, Norway, during autumn and winter 2004. Appl Environ Microbiol

17 2006;72:2212-17.

18 [16] Traub RJ, Monis PT, Robertson I, Irwin P, Mencke N, Thompson RCA.

19 Epidemiological and molecular evidence supports the zoonotic transmission of *Giardia*

20 among humans and dogs living in the same community. Parasitology 2004;128:253-62.

- 1 [17] Sulaiman IM, Fayer R, Bern C, Gilman RH, Trout JM, Schantz M, Das P, Lal AA,
2 Xiao L. Triosephosphate isomerase gene characterization and potential zoonotic
3 transmission of *Giardia duodenalis*. Emerg Infect Dis 2003;9:1444-52.
- 4 [18] Lalle M, Pozio E, Capelli G, Bruschi F, Crotti D, Caccio SM. Genetic
5 heterogeneity at the beta-giardin locus among human and animal isolates of *Giardia*
6 *duodenalis* and identification of potentially zoonotic genotypes. Int J Parasitol
7 2005;35:207-13.
- 8 [19] Homan WL, Mank TG. Human giardiasis: genotype linked differences in clinical
9 symptomatology. Int J Parasitol 2001;31:822-26.
- 10 [20] Read C, Walters J, Robertson ID, Thompson RCA. Correlation between genotype
11 of *Giardia duodenalis* and diarrhoea. Int J Parasitol 2002;32:229-31.
- 12 [21] Abe N, Kimata I, Tokoro M. Genotyping of *Giardia* isolates from humans in Japan
13 using the small subunit ribosomal RNA and glutamate dehydrogenase gene sequences.
14 Jpn J Infect Dis 2005;58:57-58.
- 15 [22] Sulaiman IM, Jiang J, Singh A, Xiao L. Distribution of *Giardia duodenalis*
16 genotypes and subgenotypes in raw urban wastewater in Milwaukee, Wisconsin. Appl
17 Environ Microbiol 2004;70:3776-80.
- 18 [23] Abe N, Kimata I, Iseki M. Identification of genotypes of *Cryptosporidium parvum*
19 isolates from a patient and a dog in Japan. J Vet Med Sci 2002;64:165-68.
- 20 [24] Read CM, Monis PT, Thompson RC. Discrimination of all genotypes of *Giardia*

- 1 *duodenalis* at the glutamate dehydrogenase locus using PCR-RFLP. Infect Genet Evol
2 2004;4:125-30.
- 3 [25] Thompson RCA, Hopkins RM, Homan WL. Nomenclature and genetic groupings
4 of *Giardia* infecting mammals. Parasitol Today 2000;16:210-13.
- 5 [26] Ali-Shtayeh MS, Hamdan AH, Shaheen SF, Abu-Zeid I, Faidy YR. Prevalence and
6 seasonal fluctuations of intestinal parasitic infections in the Nablus area, West Bank of
7 Jordan. Ann Trop Med Parasitol 1989;83:67-72.
- 8 [27] Amar CFL, Dear PH, Pedraza-Diaz S, Looker N, Linnane E, McLauchlin J.
9 Sensitive PCR-restriction fragment length polymorphism assay for detection and
10 genotyping of *Giardia duodenalis* in human feces. J Clin Microbiol 2002;40:446-52.
- 11 [28] Caccio SM, de Giacomo M, Pozio E. Sequence analysis of the beta-giardin gene
12 and development of a polymerase chain reaction-restriction fragment length
13 polymorphism assay to genotype *Giardia duodenalis* cysts from human fecal samples.
14 Int J Parasitol 2002;32:1023-30.
- 15 [29] Monis PT, Mayrhofer G, Andrews RH, Homan WL, Limper L, Ey PL. Molecular
16 genetic analysis of *Giardia intestinalis* isolates at the glutamate dehydrogenase locus.
17 Parasitology 1996;112:1-12.
- 18 [30] Lu SQ, Baruch AC, Adam RD. Molecular comparison of *Giardia lamblia* isolates.
19 Int J Parasitol 1998;28:1341-45.
- 20 [31] Lalle M, Jimenez-Cardosa E, Caccio SM, Pozio E. Genotyping of *Giardia*

1 *duodenalis* from humans and dogs from Mexico using a beta-giardin nested polymerase
2 chain reaction assay. J Parasitol 2005;91:203-5.

3 [32] Ponce-Macotela M, Martinez-Gordillo MN, Bermudez-Cruz RM,
4 Salazar-Schettino PM, Ortega-Pierres G, Ey PL. Unusual prevalence of the *Giardia*
5 *intestinalis* A-II subtype amongst isolates from humans and domestic animals in
6 Mexico. Int J Parasitol 2002;32:1201-2.

7 [33] Monis PT, Andrews RH, Mayrhofer G, Kulda J, Isaac-Renton JL, Ey PL. Novel
8 lineages of *Giardia intestinalis* identified by genetic analysis of organisms isolated from
9 dogs in Australia. Parasitology 1998;116:7-19.

10

1 **Tables and figure legends**

2

3 Table 1

4 Genotyping results assessed in this study

5 Genotypes of the PalH1 and PalH3 samples, which were identical to an assemblage A

6 reference (AF199446) and those of the PalH4 and PalH8 samples, which were identical7 to an assemblage B reference (AF199447), were determined by 18S rRNA gene

8 analysis. Assemblage A-II and multiple-subgenotype infections of assemblage B were

9 determined by GDH gene analysis.

| Sample name | Genotype |
|-------------|---------------------------|
| PalH1 | Assemblage A |
| PalH2 | Assemblage A-II |
| PalH3 | Assemblage A |
| PalH4 | Assemblage B ^a |
| PalH5 | Assemblage A-II |
| PalH6 | Assemblage A-II |
| PalH7 | Assemblage A-II |
| PalH8 | Assemblage B ^a |

10 ^aMultiple-subgenotype infection of assemblage B

11

1 Table 2

2 GDH gene sequences used in this study

| Isolate name ^a | Accession number | Assemblage ^b | Place of origin | Reference |
|---------------------------|---------------------------------|-------------------------|-----------------|-------------------|
| Ad-1 | <u>AY178735</u> | A-I | Australia | [10, 32] |
| Ad-2 | <u>L40510</u> | A-II | Australia | [10, 32] |
| BAH-12 | <u>AF069059</u> | B-III | Australia | [10, 32] |
| Ad-7 | <u>L40508</u> | B-IV | Australia | [10, 32] |
| Ad-136 | <u>U60982</u> | C (Dog) | Australia | [33] |
| Ad-148 | <u>U60986</u> | D (Dog) | Australia | [33] |
| P-15 | <u>U47632</u> | E (Livestock) | Australia | [10] |
| Ad-23 | <u>AF069057</u> | F (Cat) | Australia | [10] |
| Ad-157 | <u>AF069058</u> | G (Rat) | Australia | [10] |
| GH-135 | <u>AB195224</u> | B | Japan | [21] |
| Ad-156 | <u>AY178752</u> | B2 (Marmoset) | Australia | Direct submission |
| Ad-158 | <u>AY178753</u> | B2 (Marmoset) | Australia | Direct submission |
| gi-hum1 | <u>DQ840541</u> | B | Poland | Direct submission |
| gi-cat1 | <u>DQ840542</u> | B (Cat) | Poland | Direct submission |
| NLH25 | <u>AY826193</u> | B | Netherlands | [12] |
| NLH35 | <u>AY826197</u> | B | Netherlands | [12] |
| gd-ber4 | <u>DQ090535</u> | B | Norway | [15] |
| PalH4-1 | <u>AB295649</u> | B | Palestine | This study |
| PalH4-2 | <u>AB295650</u> | B | Palestine | This study |
| PalH4-3 | <u>AB295651</u> | B | Palestine | This study |
| PalH8-1 | <u>AB295652</u> | B | Palestine | This study |
| PalH8-2 | <u>AB295653</u> | B | Palestine | This study |
| PalH8-3 | <u>AB295654</u> | B | Palestine | This study |
| - | <u>AF069065</u> | <i>Giardia ardeae</i> | - | [10] |

3 ^aIsolate names and genotypes are derived from the DDBJ/GenBank/EMBL database and
4 reference information. ^bInformation in parentheses indicates the source of those
5 samples that were not obtained from human subjects.

6

1 Table 3
 2 Alignment of the GDH nucleotide sequences by using assemblage B subgenotypes
 3 isolated from the mixed-infection samples
 4 Substituted positions were determined by the alignment of 393-bp GDH gene fragments.
 5 Nucleotide position numbers were determined according to the reference assemblage
 6 B-III (**AF-069059**), and substitutions at the third nucleotides of codons are shown by a
 7 hyphen “-”. Genotypes of isolates from each mixed-infection sample consisted of 3
 8 subgenotypes: PalH4 (PalH4-1, PalH4-2, and PalH4-3) and PalH8 (PalH8-1, PalH8-2,
 9 and PalH8-3).

10

| Reference and Subgenotypes | Deduced amino acid and substituted nucleotide sequences | | | | | | | | | | | | | | |
|----------------------------------|---|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|
| | Asn AA- | Leu CT- | Leu CT- | Pro CC- | Gly GG- | Phe TT- | Cys TG- | Thr AC- | Gly GG- | Ile AT- | Tyr TA- | Phe TT- | Leu CT- | Arg AG- | Gly GG- |
| | 87 | 99 | 147 | 150 | 156 | 219 | 222 | 237 | 258 | 309 | 330 | 351 | 363 | 402 | 417 |
| AF069059 | C | C | T | G | C | T | C | T | G | C | C | C | C | G | G |
| PalH4-1 | C | C | T | G | C | T | C | T | G | T | C | C | T | G | G |
| PalH4-2 | T | C | T | G | C | C | C | C | A | C | T | T | C | A | A |
| PalH4-3 | C | T | C | G | C | C | C | C | G | C | T | C | C | A | G |
| PalH8-1 | C | T | C | A | C | T | T | T | G | C | C | C | C | G | G |
| PalH8-2 | C | T | T | G | T | C | C | C | G | C | T | C | C | A | G |
| PalH8-3 | C | C | T | G | C | T | C | T | G | T | C | C | C | G | G |

1 Fig. 1. NJ tree reconstructed with the GDH gene sequences obtained in this study and
2 references of *G. intestinalis*. Six isolates from 2 mixed-infection samples and the
3 assemblage A-II reference sequence (**L40510**)*, which was identical to the sequences of
4 the subgenotypes from the PalH2, PalH5, PalH6 and PalH7 samples, are shown in bold.
5 All reference sequences are shown by isolate names and their details are listed in Table
6 2. Numbers on nodes represent bootstrap values from 1000 replicates.
7

Fig. 1

