

Quantitative analysis of the Trendelenburg test and invention of a modified method

メタデータ	言語: eng 出版者: 公開日: 2017-12-05 キーワード (Ja): キーワード (En): 作成者: メールアドレス: 所属:
URL	http://hdl.handle.net/2297/46797

TITLE PAGE

Title

Quantitative Analysis of the Trendelenburg Test and Invention of a Modified Method

Author names and final degree

Kenji Fujita, MD, PhD¹ (kkkkkenji76@yahoo.co.jp)

Tamon Kabata, MD, PhD¹ (tamonkabata@yahoo.co.jp)

Yoshitomo Kajino, MD, PhD¹ (yoshitomokajino@gmail.com)

Shintaro Iwai, MD, PhD¹ (shntr.iwai@gmail.com)

Kazunari Kuroda, MD, PhD¹ (kurokazu0108@yahoo.co.jp)

Kazuhiro Hasegawa, MD¹ (kazuhiro63hasegawa@gmail.com)

Katsuo Fujiwara, PhD² (fujikatu@med.m.kanazawa-u.ac.jp)

Hiroyuki Tsuchiya, MD, PhD¹ (tsuchi@med.kanazawa-u.ac.jp)

The affiliation and address

1. Department of Orthopaedic Surgery, Graduate School of Medical Science, Kanazawa University
2. Department of Human Movement and Health, Graduate School of Medical Science, Kanazawa University

Conflict of interest statement

None

Ethical review committee statement

This investigational protocol was conducted with the approval of the Kanazawa University Graduate School of Medicine Ethics Committee. In accordance with the requirements of this review, all subjects were provided informed consent.

Please address all correspondence to:

Tamon Kabata, MD, PhD

Department of Orthopaedic Surgery, Graduate School of Medical Science, Kanazawa University,
13-1 Takaramachi, Kanazawa, Ishikawa, 920-8641, Japan

Phone: +81-76-265-2374

Fax: +81-76-234-4261

E-mail: tamonkabata@yahoo.co.jp

1 **Quantitative Analysis of the Trendelenburg Test and Invention of a Modified Method**

2

3 **Abstract**

4 *Background* While the Trendelenburg test has been used for 120 years to detect hip abductor
5 muscle weakness, the methodology has not been standardised.

6 *Purposes* This study undertook to quantitatively analyze the relation between abductor
7 muscle activity and pelvic tilt angle in the Trendelenburg one-leg stance, examine the pitfalls
8 associated with performing the T-test, and develop a modified method that will produce
9 reliable results.

10 *Methods* A convenience sample of 15 healthy males was asked to assume a one-leg stance in
11 ten different postures, five with mild flexion on the unsupported side, and five with severe
12 flexion. Trunk sway angle, pelvic tilt angle, and the pelvic on femur (POF) angle were
13 measured for each posture. Statistical analysis was used to assess differences in hip abductor
14 activity and pelvic tilt angle between the control posture and the test postures.

15 *Results* With minimum trunk sway, hip abductor muscle activity increases when the pelvis is
16 elevated and decreases when it is dropped. With trunk sway toward the test side, abductor
17 muscle activity decreased when the pelvis was elevated; with trunk sway toward the non-test
18 side, muscle activity stayed approximately constant when the pelvis was dropped.

19 *Conclusions* Based on the results we developed a modified T-test methodology that would
20 improve reliability. This test should be performed with minimum trunk sway and severe
21 flexion on the non-test side. The assessment of muscle weakness is based on whether the
22 patient can keep the single-leg standing posture when forced to elevate the pelvis, not simply
23 on the pelvic drop. In future research, we will perform the modified T-test on patients with a
24 suspected hip abductor deficiency, and assess the usefulness of the modified test.
25

26 **Introduction**

27 Weakness of the abductor muscle is the major cause of claudication resulting from diseases of
28 the hip joint. Therefore, evaluation of hip abductor muscle strength is important in diagnosing
29 and treating such diseases. The Trendelenburg (T) test was first reported by Friedrich
30 Trendelenburg in 1895 as a physical examination method for detecting severe abductor
31 muscle weakness [1]. Generally, if the patient assumes a one-leg stance and the pelvis drops
32 on the non-test (the non-stance) side, the test result is read as positive; in other words, the
33 abductor muscle is weak. The T-test has long been a popular method for physical
34 examinations; however, the details and evaluation method of the procedure are usually
35 described vaguely and have not been standardised. In 1985, Hardcastle et al [2] developed a
36 T-test method which they called the standard Trendelenburg (sT)-test. In this method, the
37 participant is instructed to elevate the pelvis as high as possible on the non-test side, and if
38 sufficient elevation of the pelvis can be maintained for 30 seconds, the test result is
39 considered negative. If insufficient elevation and drop of the pelvis occur, the test result is
40 considered positive. This method is now used worldwide as a method for evaluating hip
41 abductor function after total hip replacement (THR). However, its reliability does not yet
42 clear. In daily medical practice, we often hesitate to judge that the test is positive or negative
43 because the pelvic drop (or insufficient pelvic elevation) is not immediately obvious.
44 Additionally, we wonder whether the hip flexion angle of non-test side and a trunk sway has

45 an effect on the results. The aims of this study were to quantitatively analyze the relationship
46 between abductor muscle activity and the pelvic tilt angle in the one-leg stance posture; to
47 examine the pitfalls associated with performing the **T-test**; and ultimately, to invent a better
48 method, which we call **the modified Trendelenburg (mT)-test**.

49

50 **Materials and Methods**

51 *Participants*

52 A convenience sample of 15 healthy men (30 hips) was recruited. Inclusion criteria were as
53 follows: 1) older than 18 years, 2) no current or previous injury to the lumbar spine, pelvis,
54 or lower extremities within the past 12 months, 3) no previous surgery to the lumbar spine,
55 pelvis, or hip, 4) normal passive and active range of motion of both hips, 5) 5 of 5 scores
56 bilaterally on manual muscle testing of the hip abductor muscle. The mean age was 31.4
57 years (range, 22-55 years) , the mean body weight was 63.9 kg (range, 54.5-86.0 kg) ,
58 and the mean height was 171.5 cm (range, 163.2-180.7 cm).

59

60 **Methods**

61 *Procedures*

62 Participants assumed a one-leg stance in 10 different postures, as described below (Figure

63 1) ; measurements were taken of the trunk sway angle, the pelvic tilt angle, and the pelvic-
64 on-femur (POF) angle (Figure 2) ; and the relationship between hip abductor muscle
65 activity and the pelvic tilt angle was assessed. "Posture 1: control" was defined as a one-leg
66 stance posture in which participants were given no instruction regarding pelvic tilt. In
67 "Posture 2: pelvic drop" and "Posture 3: pelvic elevation" the participants were instructed to
68 drop the pelvis, and to elevate it, respectively. In postures 1-3, the participants were instructed
69 to minimize trunk sway, in order to eliminate any effect which the trunk sway might have on
70 hip abductor muscle activity and pelvic tilt angle. Finally, in order to assess the effect which
71 trunk sway has on hip abductor muscle activity and pelvic tilt angle, in "Posture 4: trunk
72 sway toward test side" and "Posture 5: trunk sway toward non-test side" the participants were
73 instructed to lean the trunk toward the test side and non-test side, respectively. In postures 4
74 and 5, the participants were given no instruction regarding pelvic tilt.

75 For each posture, the participants were asked to perform mild and severe flexion of the non-
76 test side hip in order to assess the effect that the flexion angle of the non-test side hip has on
77 hip abductor muscle activity and pelvic tilt angle. Mild flexion was defined as raising the toe
78 of the non-stance side as high as the medial malleolus of the stance side, for approximately
79 30 degrees of hip flexion. Severe flexion was defined as raising the toe of the non-stance side
80 as high as the knee of the stance side, approximately 80 degrees of hip flexion. Measurements
81 of the one-leg stance postures and the hip abductor muscle activity were performed once, for

82 two seconds per posture, for 10 different postures.

83 **Measurement of the one-leg stance posture (Figure 2)**

84 The Trunk sway angle, pelvic tilt angle, and pelvic on femur (POF) angle were measured for
85 each posture using retroreflective markers fixed by adhesive tape. A total of six retroreflective
86 markers were placed on the bilateral acromion processes of the scapula, the bilateral anterior
87 superior iliac spine (ASIS), and the bilateral second metatarsal head. The postures were
88 measured using a camera (EOS Kiss X3, canon, Japan) and two-dimensional motion analysis
89 software (Move-tr/2D, Library, Japan). The camera was positioned 10 meters from the
90 participants. A first photograph was taken in the bilateral-leg standing posture just after all
91 markers were placed. If the line between two markers of the bilateral acromion processes of
92 the scapula and/or the line between two markers of the bilateral ASIS were oblique relative to
93 the floor, the markers were placed correctly again. We started the measurement only after
94 ensuring these markers were placed almost horizontally to the floor. Once the participant was
95 balanced in the one-leg stance posture, the camera shot continuously every 0.3 seconds for
96 two seconds and the mean value of these angles over the two- second duration was calculated
97 with the analysis software. As illustrated in Fig. 2, the pelvic tilt angle is formed by the line
98 of the bilateral ASIS and the horizontal line. The trunk sway angle is formed by the line of
99 the bilateral acromion processes and the horizontal line. We assigned a positive value to the
100 pelvic elevation of the non-stance side, the trunk sway toward the stance side. The POF angle

101 is formed by the line of the bilateral ASIS and the line connecting the ASIS and the second
102 metatarsal head on the test side.

103 **Measurement of hip abductor muscle activity with electromyography (EMG)**

104 The hip abductor muscle activity in each one-leg posture were measured quantitatively by a
105 surface EMG of the gluteus medius muscle. Surface electrodes were used in a bipolar
106 derivation in order to record the EMG from the gluteus medius muscle, after proper skin
107 preparation to reduce electrode input impedance to below 5k Ω . EMG data were sent to a
108 computer (Dimension 9150, DELL, DELL Japan) via an A/D converter (AD16-64(LPCI)LA,
109 Contec, Japan) at 1,000 Hz with 16-bit resolution. Subsequent analyses were performed using
110 BIMUTAS II software (Kissei Comtec Co. Ltd.). EMG data were 40-Hz high-pass filtered in
111 order to exclude electrocardiographic and movement artifacts, and then were full-wave
112 rectified and integrated for two seconds, which is the same time used to measure postures
113 with the camera. Hip abductor muscle activity was expressed as relative muscle activity of
114 the gluteus medius muscle; relative muscle activity was assessed by the relative ratio of the
115 activity on the EMG recording for two seconds during 100% maximal muscle force in
116 manual muscle testing.

117 *Statistical analysis*

118 Statistical analysis was performed using SPSS (PASW Statistics Base v19; SPSS Inc.,

119 Chicago, Illinois). A two-sided paired t-test was used to assess the differences in hip abductor
120 muscle activity and the pelvic tilt angle between the control posture and each of the four test
121 postures. Data were expressed as mean and standard deviation (SD). The level of significance
122 was set at $p < 0.05$. The study had ethical approval from XXXXX Hospital Ethical Review
123 Board. All subjects gave their consent to participate in this study.

124

125 **Results**

126 Abductor muscle activity, the pelvic tilt angle, the trunk sway angle, and the POF angle on
127 the stance side of each posture are shown in detail in Table 1 (for mild flexion) and Table 2
128 (for severe flexion). For ease of understanding, the results of the abductor muscle activity and
129 measurement of each posture are shown as a pattern diagram in Figure 3. The relationship
130 between abductor muscle activity and the pelvic tilt angle is shown in Figures 4 and 5. In
131 summary, the results show that in relation to hip abductor muscle activity and pelvic tilt in the
132 control posture and under conditions of minimal trunk sway, hip abductor muscle activity
133 increased when the pelvis was elevated, and decreased when the pelvis was dropped
134 compared with that in the control posture. However, with trunk sway toward the test side, hip
135 abductor muscle activity decreased when the pelvis was elevated compared with that in the
136 control posture; with trunk sway toward the non-test side, the hip abductor muscle activity
137 stayed approximately constant when the pelvis was dropped. The difference in hip flexion

138 angle did not have much influence on the relationship between abductor muscle activity and
139 the pelvic tilt angle. Pelvises tended to be more elevated in severe hip flexion postures than
140 in mild hip flexion postures.

141

142 **Discussion**

143 The common causes of claudication resulting from diseases of the hip joint are hip pain, leg
144 length discrepancy, and weakness of the abductor muscle. Therefore, we think that
145 orthopaedic doctors should be able to diagnose the causes of claudication in patients in just
146 under a minute in the consultation room by performing the T-test, determining the presence
147 or absence of pain, and checking for leg length discrepancy on a plain X-ray. The T-test is the
148 simplest practical screening method for deciding whether or not weakness of the abductor
149 muscle is the cause of claudication.

150 The T-test was developed by Friedrich Trendelenburg in 1895, even before the widespread
151 use of radiography [1], and in the almost 120 years since then, it has become a standard
152 physical examination method for identifying weakness in the hip abductor muscles. However,
153 the details and evaluation method of the procedure have never been standardised.

154 A famous text described how to perform the classic T-test as follows: the foot on the non-test
155 side should be lifted by flexing the knee while keeping the thigh extended so that the psoas

156 muscle cannot elevate on that side [3]. Generally, a normal hip will be held stable; if the
157 pelvis drops on the non-stance side during the one-leg stance posture, the test result is
158 considered to be positive, indicating weakness of the hip on which the subject is standing.
159 Baker et al. performed the T-test in the classic way [4], but this procedure is very likely to
160 result in a pelvic drop on the non-test side, leading to false positive results. Because the psoas
161 muscle of the non-stance side and abductor muscle of the stance side seem to act in
162 coordination, it can be difficult even for normal people to elevate the pelvis on the non-stance
163 side without hip flexion of that side.

164 In 1985, Hardcastle et al [2] developed a detailed methodology for the T-test, which they
165 reported as the sT-test. In this method, the participant is instructed to elevate the pelvis as
166 high as possible on the non-test side, and if sufficient elevation can be maintained for 30
167 seconds, the test result is considered negative. Pai et al. [5] used the sT-test to conduct
168 evaluations after THR, and emphasised the significance of the method. The sT-test has also
169 been cited in many other papers [6-10], and it is currently recognised worldwide as a standard
170 method for the postoperative evaluation of the hip abduction function after THR. However,
171 the validity of the sT-test has not yet been verified in quantifiable terms.

172 The reliability of T-test and sT-test does not yet clear. Kendall et al. [11] used
173 ultrasonography, after causing a considerable decrease in hip abductor muscle strength by
174 administering a superior gluteal nerve block, to evaluate the validity of the sT-test. Their

175 results indicated that muscle weakness and pelvic drop were not correlated and they
176 concluded that the sT-test was not useful as a method for diagnosing a decrease in hip
177 abductor muscle strength. Other reports [12, 13] also have stated that the correlation between
178 abductor muscle weakness and pelvic tilt is weak. Therefore, the assessment of the results in
179 the sT-tests described in previous reports [5-10] might have depended more on the examiners'
180 preconceptions than on actual hip abductor muscle function.

181 In our daily medical practice, we also have often hesitated to make a definitive judgment as
182 to whether a particular test is positive or negative. We wanted to develop a T-test
183 methodology that would improve its accuracy in the diagnosis of hip abductor deficiency.
184 Therefore, we quantitatively analyzed the relationship between abductor muscle activity and
185 the pelvic tilt angle in the one-leg stance posture, using the results of that analysis to examine
186 the pitfalls of performing the Trendelenburg test, and to invent a better method, which we call
187 the mT-test.

188 Our measurement results indicate first, that a direct correlation between the pelvic drop (or
189 elevation) and the decrease (or the increase) of hip abductor muscle activity occurs only
190 when there is minimal trunk sway. Therefore, we specify that the mT-test should be
191 performed under the condition of minimal trunk sway.

192 Secondly, it can be stated that, in the control posture, a naturally, artless pelvic elevation
193 occurs while standing on a single leg, even if the patient is not conscious of it. Therefore, to

194 avoid false-negative results during the T-tests to detect pelvic drop due to abductor muscle
195 weakness, patients incapable of achieving a sufficient pelvic elevation will also need to be
196 included among positive cases, along with patients whose pelvis is noticeably lower than the
197 horizontal reference line. Similarly, Hardcastle et al [2] stated that they considered patients
198 with insufficient pelvic elevation as positive cases.

199 Finally, the pelvic drop is likely to occur even in the absence of abductor muscle weakness if
200 the hip abductor muscle is not being worked fully, as the pelvic drop occurs in healthy
201 subjects when they relax their muscles. Therefore, to avoid false-negative results, a forced
202 elevation of the pelvis needs to be performed during the T-test. We examined methods that
203 allow for performing that forced elevation of the pelvis. We noticed that the pelvic drop is
204 accompanied by a lateral movement of the pelvis towards the test side (Figure 6) to maintain
205 balance on one leg. We considered that the lateral movements of the pelvis during testing can
206 be prevented through immobilization if the examiner places a hand on the outer side of the
207 greater trochanter, thus forcing elevation of the pelvis. Furthermore, as noted previously,
208 natural elevation of the pelvis might more likely be achieved through severe flexion of the
209 hip joint on the non-test side.

210 From these considerations, we determined that in performing the T-test we should 1) ask
211 patients to minimize trunk sway, 2) ask them to flex hips severely on non-test side, and 3)
212 make our judgment based on whether or not the patient can elevate the pelvis when forced to

213 do so, not just from the pelvic drop. Thus, we developed a modified Trendelenburg (mT)
214 test as follows (see Figure 7).

215 1. The participant is instructed to adopt a standing posture with feet close together, and the
216 examiner immobilises the participant by holding the hip outside the greater trochanter on the
217 test side. (This is designed to prevent the greater trochanter moving outside, but the
218 examiner must not try to push it inside.)

219 2. The participant is instructed to flex the hip on the non-test side and to lift that knee high,
220 while minimizing trunk sway. If the single-leg standing posture is impossible to maintain, the
221 test result is considered positive. (The participant will find it easier to understand an order
222 to "flex the hip and lift the knee high" than "to flex the hip severely.")

223 The biggest advantage of this method is that it is not necessary to base the diagnosis on a
224 slight pelvic tilt change, because the single-leg standing posture in itself becomes difficult
225 when a hip abductor muscle deficiency exists.

226 *Limitation*

227 ~~Downing ND et al. [4] and Picado CH et al. [11] evaluated hip abductor function using the T-~~
228 ~~test before and after THR and reported a significant decrease in T test positive results.-~~
229 ~~However, it is easy to get false positive T test readings from patients with significant hip-~~
230 ~~pain, since even a patient who has normal hip abductor power in a supine position cannot~~

231 ~~produce hip abductor power in the one leg stance position when hip pain is severe. This is~~
232 ~~because when the patient produces hip abductor power while standing on one leg, the~~
233 ~~resultant force goes up to the hip joint [10], and hip pain becomes acute. Thus, T test~~
234 ~~evaluations before and after THR may only indicate lessening of hip pain rather than an~~
235 ~~improvement of the hip abduction muscle deficiency. We believe that the T test, including the~~
236 ~~mT test, may be not useful at all for assessing hip abductor deficiency before THR and soon~~
237 ~~after THR in patients with strong hip pain.~~

238 This study is for young healthy males. The original subject of the mT test is patients of
239 diseases of hip joint, for instance, osteoarthritis of the hip. However, we think that the
240 relationship between the pelvic tilt and the abductor muscle strength is equivalent in these
241 patients and healthy individuals, unless the patients have a severe hip contracture. From now
242 on, We would like to evaluate the clinical relevance of the new mT test in patients of diseases
243 of hip joint.

244

245 **Conclusions**

246 We quantitatively analyzed the relationship between abductor muscle activity and pelvic tilt
247 angle in the Tredenburg one-leg stance. The results of our analysis indicate that when we
248 perform the T-test, we should 1) ask the patient to minimize trunk sway, and to flex the hip

249 and elevate the knee high on the non-stance side, and 2) evaluate an insufficient pelvic
250 elevation as well as a pelvic drop as positive. However, since the pelvic tilt is not often
251 immediately obvious, we devised a better method, the mT-test, which does not require an
252 assessment of pelvic tilt. In future research, we will perform the mT-test on patients with a
253 suspected hip abductor deficiency, and assess the usefulness of the modified test.

254

255 **Competing interests**

256 The authors declare that they have no competing interests.

257

258 **Authors' contributions**

259 XXXXX conceived and carried out the all experiments, performed the statistical analysis, and
260 drafted the manuscript. XXXXX and XXXXX contributed in developing the study design,
261 collecting patients' data. XXXXX, XXXXX and XXXXX conceived of the study, and
262 participated in its design and coordination and helped to draft the manuscript. All authors
263 read and approved the final manuscript.

264

265 **References**

- 266 1. Trendelenburg F. The classic Trendelenburg's test: 1895 – Friedrich Trendelenburg, 1844–
267 1924 (Reprinted 1966). Clin Orthop and Relat Res. 1998 Oct;355:3–7.
- 268 2. Hardcastle P, Nade S. The significance of the Trendelenburg test. J Bone Joint Surg. 1985
269 Nov;67(5):741–746.
- 270 3. Dietrich Tönnis. Congenital Dysplasia and Dislocation of the Hip in Children and Adults.
271 With Collaboration of Helmut Legal and Reinhard Graf Translated by Terry C. Telger:
272 Springer- Verlag Berlin Heidelberg 1984: 84-85.
- 273 4. Baker AS, Bitounis VC. Abductor function after total hip replacement. An
274 electromyographic and clinical review. J Bone Joint Surg Br. 1989 Jan;71(1):47-50.
- 275 5. Pai VS. Significance of the Trendelenburg test in total hip arthroplasty. Influence of lateral
276 approaches. J Arthroplasty. 1996 Feb;11(2):174-9.
- 277 6. Ramesh M, O'Byrne JM, McCarthy N, Jarvis A, Mahalingham K, Cashman WF. Damage
278 to the superior gluteal nerve after the Hardinge approach to the hip. J Bone Joint Surg Br.
279 1996 Nov;78(6):903-6.
- 280 7. Downing ND, Clark DI, Hutchinson JW, Colclough K, Howard PW. Hip abductor strength
281 following total hip arthroplasty: a prospective comparison of the posterior and lateral
282 approach in 100 patients. Acta Orthop Scand. 2001 Jun;72(3):215-20.

283 8. Asayama I, Chamnongkich S, Simpson KJ, Kinsey TL, Mahoney OM. Reconstructed hip
284 joint position and abductor muscle strength after total hip arthroplasty. *J Arthroplasty*. 2005
285 Jun;20(4):414-20.

286 9. Picado CH, Garcia FL, Marques W Jr. Damage to the superior gluteal nerve after direct
287 lateral approach to the hip. *Clin Orthop Relat Res*. 2007 Feb;455:209-11.

288 10. Kiyama T, Naito M, Shitama H, Maeyama A. Hip abductor strengths after total hip
289 arthroplasty via the lateral and posterolateral approaches. *J Arthroplasty*. 2010 Jan;25(1):76-
290 80.

291 11. Kendall KD, Patel C, Wiley JP, Pohl MB, Emery CA, Ferber R. Steps toward the
292 validation of the Trendelenburg test: the effect of experimentally reduced hip abductor
293 muscle function on frontal plane mechanics. *Clin J Sport Med*. 2013 Jan;23(1):45-51.

294 12. Kendall KD, Schmidt C, Ferber R. The relationship between hip-abductor strength and
295 the magnitude of pelvic drop in patients with low back pain. *J Sport Rehabil*. 2010
296 Nov;19(4):422-35.

297 13. Youdas JW, Madson TJ, Hollman JH. Usefulness of the Trendelenburg test for
298 identification of patients with hip joint osteoarthritis. *Physiother Theory Pract*. 2010
299 Apr;26(3):184-94.

300

301

302 **Figure Legends**

303 **Figure 1:** The ten one-leg stance postures

304 **Figure 2:** The measurement of the one-leg stance posture by trunk sway angle, pelvic tilt
305 angle, and pelvic-on-femur angle

306 **Figure 3:** Pattern diagram of the one-leg stance postures with the hip abductor muscle
307 activity of each posture

308 **Figure 4:** The relationship between pelvic tilt angle and hip abductor muscle activity with
309 mild flexion of the hip on the non-test side

310 **Figure 5:** The relationship between pelvic tilt angle and hip abductor muscle activity with
311 severe flexion of the hip on the non-test side

312 **Figure 6:** Pelvic outside movement accompanied by pelvic drop (a. Pattern diagram, b.
313 Radiograph)

314 **Figure 7:** The modified Trendelenburg test (mT-test) method

315

316 **Table 1:** Measurement values of the one-leg stance postures and the hip abductor muscle
317 activity of each posture with mild flexion of the hip on the non-test side

318 **Table 2:** Measurement values of the one-leg stance postures and the hip abductor muscle
319 activity of each posture with severe flexion of the hip on the non-test side


Figure 1

Figure 1




the mild flexion of the hip
on non-test side

1 control 2 pelvic drop 3 pelvic elevation 4 trunk sway toward test side 5 trunk sway toward non-test side




the severe flexion of the hip
on non-test side




a. the trunk sway angle

The trunk sway angle is formed by the line of the bilateral acromion processes and the horizontal line.



b. the pelvic tilt angle

The pelvic tilt angle is formed by the line of the bilateral ASIS and the horizontal line.





c. the pelvic-on-femur angle

The pelvic-on-femur angle is formed by the line of the bilateral ASIS and the line connecting the ASIS and the second metatarsal head on the stance side.


R(L)-AP: right(left) acromion processes

R(L)-ASIS: right(left) anterior superior iliac spine


L-MH: left second metatarsal head






the pelvic tilt angle (°)



The abductor muscle activity (%)








a. the comparison of the pelvic elevation posture with the severe flexion of the hip and the pelvic drop posture with the mild flexion of the hip

b. 33 year-old man. Radiographs of Bilateral leg stance posture with feet close together (b1), and one leg stance posture of the pelvic elevation (b2) and the pelvic drop (b3). He was instructed not to move a stance foot position.

: The greater trochanter moved outside 4.5cm in the pelvic drop posture, compared with the bilateral leg stance posture and the pelvic elevation posture.

Solid lines show the lateral edge of the greater trochanter. The dashed lines connecting the bilateral femoral heads show pelvic tilt.



negative

1. The participant is instructed to adopt a standing posture with feet close together, and the examiner immobilises the participant by holding the hip outside the greater trochanter on the test side.
2. The participant is instructed to flex the hip on the non-test side and to lift that knee high, while minimizing trunk sway. If the single-leg standing posture is impossible to maintain, the test result is considered positive.

Table 1 Measurement values of the one-leg stance postures and the hip abductor muscle activity of each posture with mild flexion of the hip on the non-test side

posture	1. control	2. pelvic drop	3. pelvic elevation	4. trunk sway toward test side	5. trunk sway toward non-test side
the trunk sway angle(°)	0.2±2.1	-2.1±5.3	2.6±5.7	24.2±7.3	-20.3±6.8
instruction to participants	minimize the trunk sway	minimize the trunk sway	minimize the trunk sway	lean the trunk toward the test side	lean the trunk toward the non-test side
the pelvic tilt angle(°)	1.9±2.6	-4.3±3.2	10.2±4.7	5.6±4.3	-1.9±3.2
instruction to participants	none	drop the pelvis on non-test side	elevate the pelvis on non-test side	none	none
the POF angle (°)	84.0±2.6	77.1±3.6	92.6±4.5	89.8±4.5	77.7±3.2
instruction to participants	none	none	none	none	none
Abductor muscle activity (%)	18.5±9.9	7.6±5.5	23.9±11.8	12.4±6.3	16.8±10.4

Table 2 Measurement values of the one-leg stance postures and the hip abductor muscle activity of each posture with severe flexion of the hip on the non-test side

posture	1. control	2. pelvic drop	3. pelvic elevation	4. trunk sway toward test side	5. trunk sway toward non-test side
the trunk sway angle(°)	0.4±2.5	-3.5±4.1	1.1±4.3	24.4±7.1	-21.7±7.6
instruction to participants	minimize the trunk sway	minimize the trunk sway	minimize the trunk sway	lean the trunk toward the test side	lean the trunk toward the non-test side
the pelvic tilt angle(°)	6.8±4.3	-3.0±3.0	14.3±6.0	12.1±6.5	1.0±4.0
Instruction to participants	none	drop the pelvis on non-test side	elevate the pelvis on non-test side	none	None
the POF angle (°)	89.0±4.7	78.1±3.5	96.8±6.1	96.8±7.1	80.6±4.4
instruction to participants	none	none	none	none	none
Abductor muscle activity (%)	19.0±7.8	9.8±7.4	27.0±12.6	13.6±6.5	20.0±12.3