

Sphingosine-1-phosphate receptor 2 protects against anaphylactic shock through suppression of endothelial nitric oxide synthase in mice

メタデータ	言語: eng 出版者: 公開日: 2017-10-03 キーワード (Ja): キーワード (En): 作成者: メールアドレス: 所属:
URL	http://hdl.handle.net/2297/36259

TITLE PAGE

Original Article

Sphingosine-1-phosphate receptor-2 protects against anaphylactic shock through suppression of eNOS in mice¹

Hong Cui, MD,^{a,2} Yasuo Okamoto, MD, PhD,^a Kazuaki Yoshioka, PhD,^a Wa Du, MD, PhD,^a Noriko Takuwa, MD, PhD,^{a,b} Wei Zhang, MD, PhD,^c Masahide Asano, PhD,^d Toshishige Shibamoto, MD, PhD,^c and Yoh Takuwa, MD, PhD^{a*}

^aDepartment of Physiology, School of Medicine, and ^dDivision of Transgenic Animal Science, Advanced Science Research Center, Kanazawa University; ^bIshikawa Prefectural Nursing University; ^cDepartment of Physiology II, Kanazawa Medical University, Japan

* Corresponding author

Yoh Takuwa, MD, PhD

Department of Physiology

Kanazawa University School of Medicine

13-1 Takara-machi, Kanazawa, 920-8640, Japan

Tel: +81-76-265-2165,

Fax: +81-76-234-4223

E-mail: ytakuwa@med.kanazawa-u.ac.jp

Declaration of all sources of funding: A Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science, a Grant-in-Aid for Scientific Research on Priority Areas from the Ministry of Education, Culture, Sports, Science and Technology in Japan, funds from Takeda Science Foundation, funds from the Kanazawa University Strategic Research Development Program, and the IPNU Research Promotion Program.

Word count: 3771

ABSTRACT

Background: Sphingosine-1-phosphate receptor-2, S1P₂, is expressed in vascular endothelial cells (EC). However, the role of S1P₂ in vascular barrier integrity and anaphylaxis is not well understood. Endothelial nitric oxide (NO) synthase (eNOS) generates NO to mediate vascular leakage, compromising survival in anaphylaxis. We recently observed that endothelial S1P₂ inhibits Akt, an activating kinase of eNOS.

Objective: We tested the hypothesis that endothelial S1P₂ might suppress eNOS, exerting a protective effect against endothelial barrier-disruption and anaphylaxis.

Methods: Mice deficient in S1P₂ and eNOS underwent antigen challenge or platelet-activating factor (PAF) injection. Analyses were performed to examine vascular permeability and the underlying mechanisms.

Results: *S1pr2* deletion augmented vascular leakage and lethality after either antigen challenge or PAF injection. PAF injection induced activation of Akt and eNOS in the aorta and lung of *S1pr2*-null mice, which were augmented compared with wild-type mice. Consistently, PAF-induced increase in cyclic GMP level in the aorta was enhanced in *S1pr*-null mice. Genetic *Nos3* deletion or pharmacological eNOS blockade protected *S1pr2*-null mice from aggravation of barrier-disruption after antigen challenge and PAF injection. EC isolated from *S1pr2*-null mice exhibited greater stimulation of Akt and eNOS with enhanced NO production in response to S1P or PAF, compared with wild-type EC. Moreover, *S1pr2*-deficient EC showed more severe disassembly of adherens junctions with augmented *S*-nitrosylation of β -catenin in response to PAF, which was restored by pharmacological eNOS blockade.

Conclusion: S1P₂ diminishes harmful robust eNOS stimulation and thereby attenuates vascular barrier-disruption, suggesting potential usefulness of S1P₂ agonists as novel

therapeutic agents for anaphylaxis.

Key Messages

- Endothelial S1P₂ plays a protective role against acute barrier disruption during anaphylaxis.
- The barrier-protective effect of S1P₂ is mediated through suppression of eNOS.

Capsule Summary

Endothelial S1P₂ protects against vascular leakage in anaphylaxis through suppressing production of nitric oxide, which disrupts the endothelial barrier, and could be a promising novel target for treating anaphylaxis.

Key words: sphingosine-1-phosphate, S1P₂, vascular permeability, anaphylaxis, endothelial nitric oxide synthase, nitric oxide, nitrosylation, β -catenin, VE-cadherin, adherens junction

Abbreviations used:

AJ: adherens junctions

BSA: bovine serum albumin

cGMP: cyclic GMP

EC: endothelial cell

eNOS: endothelial NO synthase

HCT: hematocrit

L-NAME: N ω -L-nitro-argininemethyl ester

MLEC: mouse lung endothelial cell

NO: nitric oxide

OVA: ovalbumin

PAF: platelet-activating factor

PI3K: phosphatidylinositol 3-kinase

PTEN: phosphatase and tensin homolog deleted from chromosome 10

S1P: sphingosine-1-phosphate

VEGF: vascular endothelial growth factor-A

WT: wild-type

INTRODUCTION

Sphingosine-1-phosphate (S1P) has been identified as an important regulator of vascular and immune functions.^{1,2} S1P is synthesized by sphingosine kinases-1 and -2 in a wide variety of cells.³ Plasma S1P, which is present at the concentration of $\sim 10^{-6}$ M, is derived from erythrocytes, vascular endothelial cells (EC) and other cells.⁴⁻⁶ Most of the biological activities of S1P are mediated by the S1P-specific G protein-coupled receptor family S1P₁~ S1P₅.⁷ Among these, S1P₁, S1P₂ and S1P₃ are widely expressed major receptor subtypes. EC in culture express readily detectable levels of S1P₁ and S1P₃ whereas S1P₂ expression is much lower or undetectable.^{8,9} However, we recently observed by analyzing LacZ-knockin mice at the *S1pr2* locus that EC in a variety of organs express S1P₂ in vivo.¹⁰ S1P₁ and S1P₂ exert distinct effects on cell responses, including migration and proliferation, through the preferential coupling to G_i and G_{12/13}, respectively.¹¹

Platelet-activating factor (PAF), histamine and leukotrienes are edematogenic mediators that disrupt vascular barrier integrity, which is sustained mainly by the endothelial adherens junction (AJ) comprising VE-cadherin and its associated proteins.¹² In anaphylaxis, these edematogenic mediators are released mainly from mast cells through antigen engagement of mast cell-associated IgE and act on vascular endothelium and smooth muscle to cause severe vascular leakage and hypotension.¹³ To establish novel therapies for anaphylaxis, it is important to better understand the pathophysiological mechanisms underlying vascular barrier disruption. Both in vitro and in vivo studies have documented that S1P maintains vascular endothelial barrier function via endothelial S1P₁.^{14,15} A subsequent study showed that plasma S1P is required to prevent vascular leakage in anaphylaxis after antigen challenge or injections of PAF and histamine.¹⁶ Thus, the plasma S1P–endothelial S1P₁ axis plays

a central role in maintaining vascular barrier integrity.¹⁷ In contrast to S1P₁, pharmacological blockade of S1P₂ was reported to reduce histamine-induced vascular leakage.¹⁸ More recently, it has been shown that genetic deletion of *S1pr2* attenuated vascular leakage and hypothermia in anaphylaxis after antigen challenge.¹⁹ *S1pr2* deletion was accompanied by diminished elevation of plasma histamine and cytokines after antigen challenge, suggesting that stimulation of mast cell activation by S1P₂ is a determinant of vascular hyperpermeability and anaphylaxis. Unlike this report, however, another group showed that S1P₂ was rather vital for recovery from histamine-induced hypothermia through regulating vascular tone and histamine clearance, which suggests that S1P₂ exerts a protective effect against anaphylaxis at a site distal to mast cell activation.^{20,21} Thus, the in vivo role of S1P₂ in anaphylaxis is not yet fully understood.

Previous studies demonstrated that endothelial nitric oxide (NO) synthase (eNOS or NOS3)-derived NO is involved in an increase of vascular permeability caused by PAF and vascular endothelial growth factor-A (VEGF).²²⁻²⁴ The activity of the calmodulin-dependent enzyme eNOS is stimulated by an increase in intracellular free Ca²⁺, interaction with regulatory proteins, and phosphorylation mediated largely by Ca²⁺-independent protein kinases including Akt and protein kinase A.²⁵ Phosphorylation of eNOS sensitizes eNOS to Ca²⁺, resulting in augmented activation of eNOS. Most edematogenic mediators, including PAF and histamine, act through G protein-coupled receptors. Importantly, anaphylactic shock has been shown to depend on eNOS activation mediated by endothelial G_{q/11} and phosphatidylinositol 3-kinase (PI3K)-Akt.^{26,27} We recently found that endogenous S1P₂ mediates inhibition of Akt in EC,¹⁰ unlike S1P₁ which stimulates eNOS via Akt.²⁸ These observations raise the possibility that S1P₂ might suppress eNOS through inhibition of Akt, exerting a

rescue effect on vascular hyperpermeability.

In the present study, we tested this hypothesis by determining how genetic *Nos3* deletion and pharmacological eNOS blockade affected S1P₂-mediated effects on vascular hyperpermeability in anaphylaxis in mice. Our results indicate that S1P₂ protects mice from vascular barrier disruption elicited by either antigen challenge or PAF injection and that this protective effect of S1P₂ is mediated through suppression of anaphylaxis-associated eNOS stimulation. These observations provide the basis for targeting S1P₂ as a novel treatment of anaphylaxis.

METHODS

Supplemental information can be found in the Methods section in this article's Online Repository.

Mice

All animal experiments were conducted according to the Fundamental Guidelines for Proper Conduct of Animal Experiment and Related Activities in Academic Research Institutions under the jurisdiction of the Ministry of Education, Culture, Sports, Science and Technology of Japan, and were approved by the Committee on Animal Experimentation of Kanazawa University. *S1pr2*-null (*S1pr2*^{-/-}) mice were previously described.¹⁰ *S1pr2*^{-/-} mice (129Ola;C57BL/6J mixed background) were back-crossed with C57BL/6J mice (#000664, Charles River) once. In some experiments, we employed another *S1pr2*-deleted (*S1pr2*^{LacZ/LacZ}) mouse in which LacZ is knocked in at the *S1pr2* locus.¹⁰ *Nos3*^{-/-} mice (#002684, C57BL/6J background) were obtained from the Jackson Laboratory. *S1pr2*^{-/-}; *Nos3*^{-/-} mice and wild-type (WT) littermates were generated by heterozygous matings.

Statistics

All values are expressed as mean ± SEM. Data were analyzed using unpaired, two-tailed Student's *t*-test, and two-way ANOVA followed by Bonferroni post test to determine statistical significance using GraphPad Prism software. Values of *P* < 0.05 were considered statistically significant.

RESULTS

S1P₂ protects mice from vascular hyperpermeability in anaphylaxis

We studied how S1P₂ affected vascular hyperpermeability and mortality using two models of active systemic anaphylaxis, which predominantly depends on PAF.²⁹ Mice had been sensitized with bovine serum albumin (BSA) or ovalbumin (OVA) and challenged with these antigens. After BSA challenge, *S1pr2*^{-/-} mice displayed greater extravasation of i.v. injected Evans blue in the lung compared with WT mice (Fig 1, A and B). Consistent with this, *S1pr2*^{-/-} mice showed higher hematocrit (HCT) due to augmented plasma leakage compared with WT mice (Fig 1, C). Evans blue leakage under saline administration was similar in *S1pr2*^{-/-} and WT mice, suggesting that basal vascular permeability was not altered in *S1pr2*^{-/-} mice. *S1pr2*^{-/-} mice began to die within 10 min after the antigen challenge and all *S1pr2*^{-/-} mice died by 30 min, whereas WT mice started to die as late as 25 min and 45% of WT mice survived the antigen challenge at 60 min (Fig 1, D). Similarly, in OVA-sensitized *S1pr2*^{-/-} mice, challenge with OVA augmented Evans blue leakage and HCT elevation compared with those in WT mice (Fig E1).

We next studied the vascular permeability responses to i.v. injection of the anaphylactic mediator PAF. *S1pr2*^{-/-} mice showed greater PAF-induced extravasation of Evans blue dye in the lung than did WT mice (Fig E2, A and B). Histological analysis of the lungs showed more pronounced accumulation of perivascular fluid in *S1pr2*^{-/-} mice after PAF injection, compared with WT mice (Fig E2, C). *S1pr2*^{-/-} mice exhibited higher HCT compared with that of WT mice at 10 min after PAF injection (Fig E2, D). Consequently, survival after PAF injection was severely impaired in *S1pr2*^{-/-} mice: 90% of WT mice survived PAF challenge whereas less than 10% of *S1pr2*^{-/-} mice survived (Fig E2, E). We also analyzed the effect of PAF on dermal

vascular permeability in *S1pr2*^{-/-} and WT mice using a Miles assay. Basal vascular permeability was similar in *S1pr2*^{-/-} and WT mice (Fig E2, F). Intradermal injection of PAF induced a dose-dependent increase in leakage of Evans blue dye in both genotypes, with greater leakage of Evans blue in *S1pr2*^{-/-} mice compared with WT mice. *S1pr2*^{-/-} mice also displayed enhanced dermal leakage of Evans blue dye after local histamine injection compared with WT mice (data not shown). To delineate how the genetic background of mice affected the anaphylactic responses, we analyzed *S1pr2*^{LacZ/LacZ} mice¹⁰ that had been back-crossed with C57BL/6 mice 4 times and WT littermates. *S1pr2*^{LacZ/LacZ} mice showed exaggerated responses of Evans blue leakage and HCT elevation after PAF challenge compared with WT mice (Fig E3), which were similar to those in *S1pr2*^{-/-} mice. In these experiments, we anesthetized mice for a very short time (~1 min) with ether. In the non-anesthetized condition, we observed essentially the same exaggerated responses in HCT and death after PAF injection in *S1pr2*^{-/-} mice compared with WT mice (Fig E4). These data indicate that *S1pr2*^{-/-} mice show enhanced vascular permeability with increased mortality in anaphylaxis.

Akt/eNOS activation is augmented in aorta and lung of *S1pr2*^{-/-} mice after PAF challenge

S1pr2^{-/-} and WT mice, which received saline injection, showed similar levels of basal phosphorylation of Akt (Fig 2, A and C) and eNOS (Fig 2, B and D) in both aorta and lung. PAF injection stimulated Akt phosphorylation in aorta from both *S1pr2*^{-/-} and WT mice but with a greater increase in *S1pr2*^{-/-} mice (Fig 2, A). In contrast, PAF-induced activation of ERK or p38MAPK was not different between *S1pr2*^{-/-} and WT mice (data not shown). Notably, PAF injection increased eNOS phosphorylation in aorta from both mouse groups with a greater degree of stimulation in *S1pr2*^{-/-} mice (Fig 2, B).

Similarly to aorta, PAF increased phosphorylation in Akt and eNOS in lung from both *S1pr2*^{-/-} and WT mice with greater degrees of stimulation in *S1pr2*^{-/-} mice (Fig 2, C and D). NO synthesized by eNOS stimulates the formation of cyclic GMP (cGMP) in vascular wall. The basal level of cGMP in aorta was similar between *S1pr2*^{-/-} and WT mice (Fig 2, E). PAF injection induced a larger increase in aortic cGMP level in *S1pr2*^{-/-} mice than in WT mice. These observations raise the possibility that the exaggerated increase in vascular permeability in *S1pr2*^{-/-} mice might be due to augmented activation of the Akt/eNOS signaling pathway and generation of NO in the vasculature.

Pharmacological inhibition of NO synthase or genetic deletion of *Nos3* protect *S1pr2*^{-/-} mice from vascular hyperpermeability in anaphylaxis

Administration of N ω -L-nitro-argininemethyl ester (L-NAME) 3 h before PAF injection prevented the exaggerated HCT elevation after PAF challenge in *S1pr2*^{-/-} mice (Fig E5, A). L-NAME also protected *S1pr2*^{-/-} mice against PAF-induced lethality (Fig E5, B). These observations suggest that NO is involved in PAF-induced vascular leakage and lethality in *S1pr2*^{-/-} mice. To elucidate the role of eNOS in exaggerated PAF-induced vascular leakage in *S1pr2*^{-/-} mice, we crossed *S1pr2*^{-/-} mice with *Nos3*^{-/-} mice to generate *S1pr2*^{-/-};*Nos3*^{-/-} compound mutant mice. The HCT after PAF injection in *S1pr2*^{-/-};*Nos3*^{-/-} mice was lower than that in *S1pr2*^{-/-} mice and was not different from that in WT mice or *Nos3*^{-/-} mice (Fig 3, A). Thus, augmentation of the PAF-induced HCT elevation in *S1pr2*^{-/-} mice was abolished by *Nos3* deficiency. Consistently, *S1pr2*^{-/-};*Nos3*^{-/-} mice were almost completely rescued from PAF-induced lethality (Fig 3, B).

We also studied the effects of *Nos3* deficiency on vascular responses in

anaphylaxis after BSA challenge. The magnitude of the increase in plasma histamine level after antigen challenge was similar in WT, *S1pr2*^{-/-}, *Nos3*^{-/-}, and *S1pr2*^{-/-};*Nos3*^{-/-} mice (Fig E6, A). The HCT at 10 min after antigen challenge in *S1pr2*^{-/-};*Nos3*^{-/-} mice was lower than that in *S1pr2*^{-/-} mice and similar to that in WT mice and *Nos3*^{-/-} mice (Fig 3, C). These observations suggest that eNOS in the vascular endothelium is responsible for *S1pr2* deficiency-induced aggravation of vascular hyperpermeability in anaphylaxis. We studied the hypotensive response to an antigen challenge in these mutant mice. A previous study³⁰ documented that in *S1pr2*^{-/-} mice, the in vivo pressor response and the in vitro vascular contractile response to phenylephrine were diminished, the latter of which was endothelium-dependent. *S1pr2*^{-/-} mice showed significantly lower steady state systolic and diastolic blood pressure than WT mice while both *Nos3*^{-/-} mice and *S1pr2*^{-/-};*Nos3*^{-/-} mice displayed higher systolic blood pressure than WT mice (Table E1). Antigen challenge induced a rapid drop in blood pressure to a level of ~40 mmHg in anesthetized WT mice, and hypotension persisted for more than 40 min (Fig 3, D). In *S1pr2*^{-/-} mice, blood pressure declined more rapidly and became unmeasurably low, leading to death within 25 min. Under the same condition, *S1pr2*^{-/-};*Nos3*^{-/-} mice and *Nos3*^{-/-} mice showed only a mild, transient reduction in blood pressure and recovered by 50 min. These observations indicate that *S1pr2*^{-/-} mice exhibit severe fatal circulatory shock and that deletion of *Nos3* gene protects both *S1pr2*^{-/-} and WT mice from anaphylactic shock. Antigen challenge in WT mice induced a decline in core body temperature (mean change of -1.6°C at 20 min) (Fig E6, B). *S1pr2*^{-/-} mice displayed more severe hypothermia compared with WT mice. *S1pr2*^{-/-};*Nos3*^{-/-} mice and *Nos3*^{-/-} mice exhibited only a mild, transient decrease in body temperature, which was less marked compared with that in WT mice. Consistent with all these data, *S1pr2*^{-/-};*Nos3*^{-/-} mice were protected from antigen

challenge-induced lethality: *S1pr2*^{-/-} single mutant mice started to die at 8 min after antigen challenge with 85% mortality at 60 min, whereas *S1pr2*^{-/-};*Nos3*^{-/-} mice began to die at 45 min and mortality at 60 min was 38% (Fig 3, E). *Nos3*^{-/-} single mutant mice were also protected from antigen challenge-induced lethality as reported previously.²⁶

S1P₂ negatively regulates Akt–eNOS pathway in vascular EC

In lung EC (MLEC) from WT mice, PAF stimulated phosphorylation of both Akt and eNOS (Fig 4, A and B). S1P alone did not affect phosphorylation of Akt or eNOS but rather reduced PAF-induced phosphorylation of eNOS in WT MLEC. In contrast to WT MLEC, S1P by itself stimulated phosphorylation of Akt and eNOS in *S1pr2*^{-/-} MLEC. This is most likely because S1P₁, which is a major S1P receptor in *S1pr2*^{-/-} MLEC, mediated S1P-induced phosphorylation of Akt and eNOS whereas S1P₂ counteracted S1P₁-mediated phosphorylation of Akt and eNOS in WT MLEC. Furthermore, PAF stimulated phosphorylation of Akt and eNOS to a greater degree in *S1pr2*^{-/-} MLEC compared with WT MLEC. Unlike WT MLEC, S1P did not reduce PAF-induced phosphorylation of Akt and eNOS in *S1pr2*^{-/-} MLEC, so that phosphorylation of Akt and eNOS in *S1pr2*^{-/-} MLEC stimulated with S1P plus PAF was substantially greater than in WT MLEC. We examined how S1P₂ affected the production of NO in MLEC. In MLEC isolated from WT mice, either PAF or VEGF but not S1P stimulated the generation of NO (Fig 4, C). In *S1pr2*^{-/-} MLEC, PAF and VEGF stimulated NO production to a greater degree compared with WT EC. Moreover, S1P by itself modestly stimulated NO production in *S1pr2*^{-/-} MLEC, which was consistent with the results of eNOS phosphorylation (Fig 4, B). In addition, S1P slightly enhanced PAF-induced NO production in *S1pr2*^{-/-} MLEC. These observations

suggest that S1P₂ exerts inhibitory effects on NO production in EC through reducing phosphorylation of Akt and eNOS.

S1P₂ contributes to maintaining AJ through inhibiting PAF-induced S-nitrosylation of β -catenin

NO mediates disruption of endothelial AJ, which comprise VE-cadherin and intracellular associated proteins including β -catenin and p120 catenin,¹² and increases vascular permeability.^{22, 23, 25} Recent studies^{31, 34} demonstrated that VEGF stimulation of eNOS induced S-nitrosylation of β -catenin, leading to disassembly of AJ complexes. We examined how S1P₂ influenced PAF-induced disassembly of AJ and S-nitrosylation of β -catenin in a monolayer of MLEC. Immunofluorescent staining with anti-VE-cadherin and anti- β -catenin antibodies showed that VE-cadherin and β -catenin were uniformly distributed at cell-cell contacts in both WT and *S1pr2*^{-/-} MLEC in a non-treated condition, indicating that AJ were formed (Fig 5, A). PAF treatment resulted in a discontinuous, zigzag pattern of staining of both VE-cadherin and β -catenin in both WT and *S1pr2*^{-/-} MLEC, indicating disruption of AJ. PAF-induced disruption of AJ was more severe in *S1pr2*^{-/-} MLEC than in WT MLEC. Pretreatment with L-NAME almost totally prevented disruption of AJ in both WT and *S1pr2*^{-/-} MLEC. We analyzed S-nitrosylation of β -catenin in MLEC treated with PAF and S1P in the presence and absence of L-NAME. In WT EC, PAF plus S1P tended to increase S-nitrosylation of β -catenin but not significantly (Fig 5, B). In contrast, PAF plus S1P substantially increased S-nitrosylation of β -catenin in *S1pr2*^{-/-} MLEC. L-NAME abolished S-nitrosylation of β -catenin induced by PAF plus S1P in both WT and *S1pr2*^{-/-} MLEC. These results suggest that S1P₂ inhibits PAF-induced disassembly of AJ through mechanisms involving inhibition of NO production and thereby

prevention of *S*-nitrosylation of β -catenin.

DISCUSSION

The present study provides evidence supporting that S1P₂ is protective against acute vascular barrier disruption, based on the composite results obtained in two anaphylaxis models of antigen challenge and PAF injection. Our study also revealed the mechanisms underlying the protective effect of S1P₂: S1P₂ suppressed harmful robust activation of eNOS, which plays a key role in vascular leakage in anaphylaxis. Thus, our study identified the role of a novel endothelial signaling pathway comprising S1P₂–Akt inhibition–eNOS inhibition, which plays a protective role against vascular barrier-disruption and may represent a promising new target for treating vascular hyperpermeability.

We studied the effects of *S1pr2* deficiency on acute vascular barrier disruption in active anaphylaxis and PAF injection. In the active anaphylaxis model, our data demonstrated that *S1pr2* deficiency exacerbated vascular leakage, hypotension, hypothermia and lethality without causing an increase in plasma histamine level (Figs 1 and 3, and Fig E6). Moreover, *S1pr2* exacerbated vascular leakage and lethality provoked by injection of the anaphylactic mediator PAF (Fig E2). These results together imply that S1P₂ alleviates acute vascular permeability largely by acting at a site distal to the release of anaphylactic mediators from mast cells; i.e., very likely at the vasculature itself. Our observations are consistent with the studies by Olivera et al.,^{20,21} which showed that histamine injection-induced hypothermia and hypotension were aggravated in *S1pr2*^{-/-} mice, but are not consistent with that by Oskeritzian et al.,¹⁹ which showed that *S1pr2* deficiency alleviated vascular leakage and hypothermia after antigen challenge, contrasting to our study. The findings by Oskeritzian et al.¹⁹ also differ from our observations, in that PAF or histamine injection-induced vascular leakage and hypothermia were not different between

S1pr2^{-/-} and WT mice. The reasons for the discrepancies between their study and the present study are not clear. However, it is possible that a difference in the experimental models, i.e., passive anaphylaxis¹⁹ vs. active anaphylaxis (our study), might have led to the different results. The genetic backgrounds of mice were not exactly identical but very similar (mixed background of C57BL/6 x 129/Sv¹⁹ vs. back-crossing once or twice of C57BL/6 x 129/Ola with C57BL/6 (our study)).

The present study demonstrated that endothelial S1P₂ inhibits eNOS (Fig 4). *S1pr2* deficiency de-inhibits eNOS in the vasculature in vivo (Fig 2). NO generated by eNOS is involved in vascular hyperpermeability in response to pro-inflammatory mediators such as PAF and VEGF.²²⁻²⁴ Moreover, acute hypotension and lethality after antigen challenge and PAF injection entirely depend on the PI3K/Akt/eNOS pathway.²⁶ Our results also showed that inhibition of eNOS action by genetic or pharmacological manipulation in *S1pr2*^{-/-} mice markedly improved vascular leakage, hypotension and survival rate after antigen challenge, thus abrogating the effects of *S1pr2* deficiency (Fig 3 and Fig E5). All these observations together indicate that S1P₂ provides a protective effect against vascular barrier disruption largely by suppressing eNOS.

NO regulates many biological processes including vasorelaxation, through stimulating soluble guanylate cyclase and cGMP production.³² However, the previous study²⁶ showed that a specific inhibitor of soluble guanylate cyclase or genetic deletion of soluble guanylate cyclase- α 1, a major isoform in the vasculature, did not protect mice from PAF-induced lethality, suggesting that NO might mediate shock in a manner independent of cGMP. In addition to guanylate cyclase-cGMP, NO may affect biological processes through causing S-nitrosylation of proteins.³³ AJ control endothelial barrier function.¹² Recent studies showed that VEGF induced

S-nitrosylation of β -catenin, resulting in the dissociation of β -catenin from VE-cadherin and disassembly of AJ complexes.^{31, 34} In pulmonary EC, PAF and S1P induced S-nitrosylation of β -catenin with disassembly of VE-cadherin and β -catenin at cell-cell contacts, which was augmented by *S1pr2* deficiency (Fig 5). Of note, pharmacological blockade of eNOS restored the aggravation of S-nitrosylation of β -catenin and AJ disassembly by *S1pr2* deficiency. Therefore, our data together with these recent observations indicate that S1P₂ exerts a protective effect on barrier functions through mechanisms involving inhibition of S-nitrosylation of β -catenin and consequent disassembly of the AJ.

Akt is a major Ca²⁺-independent eNOS-activating protein kinase.³⁵ *S1pr2* deficiency augmented activation of Akt as well as eNOS in the vasculature (Fig 2). Therefore, it is likely that increased Akt activity at least in part mediates enhanced eNOS phosphorylation in *S1pr2*^{-/-} mice. Endogenously expressed S1P₂ is nearly exclusively coupled to G_{12/13} and downstream Rho.³⁶ S1P₂ was previously shown to stimulate 3'-specific phosphatase of phosphoinositides, "phosphatase and tensin homolog deleted from chromosome 10 (PTEN)", via Rho kinase.³⁷ Phosphatidylinositol-3,4,5-P₃ generated by PI3K mediates Akt stimulation. S1P₂-mediated stimulation of PTEN decreases the cellular level of phosphatidylinositol-3,4,5-P₃, thereby resulting in inhibition of Akt activation.³⁷ The previous gene knockout study by Korhonen et al.²⁷ showed that endothelial G_{q/11}-mediated signaling is crucially involved in NO formation and vascular barrier disruption induced by PAF and other mediators. G_{q/11} mediates activation of the calmodulin-dependent enzyme eNOS through both mechanisms of Ca²⁺ mobilization and Akt-mediated eNOS phosphorylation.³⁵ Hence, the study by Korhonen et al.²⁷ and our study together suggest that the PAF receptor, which is a G_{q/11}-coupled receptor,

mediates an increase in vascular permeability through eNOS stimulation whereas S1P₂ counteracts this G_{q/11}-mediated action through G_{12/13}-mediated suppression of the Akt–eNOS pathway.

Among the S1P receptors, S1P₁ has been shown to potentiate endothelial barrier integrity in vivo and in vitro.^{8, 14, 38} A recent study using sphingosine kinases-null mice indicated that plasma S1P is required for vascular integrity.¹⁶ Furthermore, administration of synthetic S1P₁ agonists, which powerfully induce downregulation of S1P₁, but also that of S1P₁ antagonists induced pulmonary edema.^{15, 17, 39} Therefore, S1P₁ seems to maintain vascular integrity in a tonic manner.¹⁶ S1P₁ and S1P₂ activate distinct signaling pathways: S1P₁ mediates G_i-dependent Rac activation, which is implicated in the vascular barrier integrity-maintaining action of S1P₁,^{8, 14} whereas S1P₂ mediates inhibition of Rac via G_{12/13}.^{10, 36} Moreover, S1P₁ stimulates eNOS phosphorylation through G_i–PI3K–Akt, unlike S1P₂.⁴⁰ However, the stimulatory effect of S1P₁ on phosphorylation of Akt and eNOS seems to be counteracted by S1P₂ in WT EC, as suggested by our observation that S1P increased phosphorylation of Akt and eNOS in *S1pr2*-deficient EC but not WT EC (Fig 4). These distinct signaling properties likely underlie the differential roles of S1P₁ and S1P₂ in the regulation of barrier integrity: *S1pr2* deficiency does not affect baseline vascular permeability but prevents enhanced vascular leakage after antigen challenge or injections of PAF and histamine (Figs 1, and Figs E1 and E2).²⁰ Consistent with this is the previous observation that baseline permeability was not affected by the presence or absence of eNOS.^{22, 23} Different from S1P₂, S1P₁ maintains not only constitutive barrier integrity but protects against barrier disruption caused by anaphylaxis and inflammation.^{15, 16, 39} It is likely that S1P₁ protects the integrity of AJ through a mechanism different from eNOS–NO regulation. Thus, these observations together with the previous

observations²⁷ suggest that receptor signaling mediated by $G_{q/11}$, G_i and $G_{12/13}$ play distinct critical roles in triggering, preventing and limiting vascular leakage.

We have demonstrated that $S1P_2$ suppresses stimulation of eNOS and NO generation in EC through inhibition of Akt during anaphylaxis. As a result, $S1P_2$ protects against the disassembly of endothelial AJ by inhibiting *S*-nitrosylation of β -catenin, and thereby vascular leakage. An $S1P_2$ -selective agonist may have a distinct clinical benefit over an $S1P_1$ -selective agonist as a therapeutic for anaphylaxis.

REFERENCES

1. Rosen H, Gonzalez-Cabrera P, Marsolais D, Cahalan S, Don AS, Sanna MG. Modulating tone: The overture of s1p receptor immunotherapeutics. *Immunol Rev* 2008;223:221-235.
2. Skoura A, Hla T. Lysophospholipid receptors in vertebrate development, physiology, and pathology. *J Lipid Res* 2009;50 Suppl:S293-298.
3. Spiegel S, Milstien S. Functions of the multifaceted family of sphingosine kinases and some close relatives. *J Biol Chem* 2007;282:2125-2129.
4. Pappu R, Schwab SR, Cornelissen I, Pereira JP, Regard JB, Xu Y, Camerer E, Zheng YW, Huang Y, Cyster JG, Coughlin SR. Promotion of lymphocyte egress into blood and lymph by distinct sources of sphingosine-1-phosphate. *Science* 2007;316:295-298.
5. Venkataraman K, Lee YM, Michaud J, Thangada S, Ai Y, Bonkovsky HL, Parikh NS, Habrukowich C, Hla T. Vascular endothelium as a contributor of plasma sphingosine 1-phosphate. *Circ Res* 2008;102:669-676.
6. Yatomi Y, Igarashi Y, Yang L, Hisano N, Qi R, Asazuma N, Satoh K, Ozaki Y, Kume S. Sphingosine 1-phosphate, a bioactive sphingolipid abundantly stored in platelets, is a normal constituent of human plasma and serum. *J Biochem* 1997;121:969-973.
7. Ishii I, Fukushima N, Ye X, Chun J. Lysophospholipid receptors: Signaling and biology. *Annu Rev Biochem* 2004;73:321-354.
8. Lee MJ, Thangada S, Claffey KP, Ancellin N, Liu CH, Kluk M, Volpi M, Sha'afi RI, Hla T. Vascular endothelial cell adherens junction assembly and morphogenesis induced by sphingosine-1-phosphate. *Cell* 1999; 99:301-312.
9. Ryu Y, Takuwa N, Sugimoto N, Sakurada S, Usui S, Okamoto H, Matsui O,

- Takuwa Y. Sphingosine-1-phosphate, a platelet-derived lysophospholipid mediator, negatively regulates cellular rac activity and cell migration in vascular smooth muscle cells. *Circ Res* 2002;90:325-332.
10. Du W, Takuwa N, Yoshioka K, Okamoto Y, Gonda K, Sugihara K, Fukamizu A, Asano M, Takuwa Y. S1p(2), the G protein-coupled receptor for sphingosine-1-phosphate, negatively regulates tumor angiogenesis and tumor growth in vivo in mice. *Cancer Res* 2010;70:772-781.
 11. Takuwa Y, Okamoto Y, Yoshioka K, Takuwa N. Sphingosine-1-phosphate signaling in physiology and diseases. *Biofactors* 2012;38(5):329-37
 12. Dejana E, Tournier-Lasserre E, Weinstein BM. The control of vascular integrity by endothelial cell junctions: Molecular basis and pathological implications. *Dev Cell* 2009;16:209-221.
 13. Finkelman FD. Anaphylaxis: Lessons from mouse models. *J Allergy Clin Immunol* 2007;120:506-515.
 14. Garcia JG, Liu F, Verin AD, Birukova A, Dechert MA, Gerthoffer WT, Bamberg JR, English D. Sphingosine 1-phosphate promotes endothelial cell barrier integrity by edg-dependent cytoskeletal rearrangement. *J Clin Invest* 2001;108:689-701.
 15. Sanna MG, Wang SK, Gonzalez-Cabrera PJ, Don A, Marsolais D, Matheu MP, Wei SH, Parker I, Jo E, Cheng WC, Cahalan MD, Wong CH, Rosen H. Enhancement of capillary leakage and restoration of lymphocyte egress by a chiral s1p1 antagonist in vivo. *Nat Chem Biol* 2006;2:434-441.
 16. Camerer E, Regard JB, Cornelissen I, Srinivasan Y, Duong DN, Palmer D, Pham TH, Wong JS, Pappu R, Coughlin SR. Sphingosine-1-phosphate in the plasma compartment regulates basal and inflammation-induced vascular leak

- in mice. *J Clin Invest* 2009;119:1871-1879.
17. Marsolais D, Rosen H. Chemical modulators of sphingosine-1-phosphate receptors as barrier-oriented therapeutic molecules. *Nat Rev Drug Discov* 2009;8:297-307.
 18. Lee JF, Gordon S, Estrada R, Wang L, Siow DL, Wattenberg BW, Lominadze D, Lee MJ. Balance of S1P1 and S1P2 signaling regulates peripheral microvascular permeability in rat cremaster muscle vasculature. *Am J Physiol Heart Circ Physiol* 2009;296:H33-42.
 19. Oskeritzian CA, Price MM, Hait NC, Kapitonov D, Falanga YT, Morales JK, Ryan JJ, Milstien S, Spiegel S. Essential roles of sphingosine-1-phosphate receptor 2 in human mast cell activation, anaphylaxis, and pulmonary edema. *J Exp Med* 2010;207:465-474.
 20. Olivera A, Eisner C, Kitamura Y, Dillahunt S, Allende L, Tuymetova G, Watford W, Meylan F, Diesner SC, Li L, Schnermann J, Proia RL, Rivera J. Sphingosine kinase 1 and sphingosine-1-phosphate receptor 2 are vital to recovery from anaphylactic shock in mice. *J Clin Invest* 2010;120:1429-1440.
 21. Olivera A, Dillahunt SE, Rivera J. Interrogation of sphingosine-1-phosphate receptor 2 function in vivo reveals a prominent role in the recovery from IgE and IgG-mediated anaphylaxis with minimal effect on its onset. *Immunol Lett* 2013;150:89-96.
 22. Fukumura D, Gohongi T, Kadambi A, Izumi Y, Ang J, Yun CO, Buerk DG, Huang PL, Jain RK. Predominant role of endothelial nitric oxide synthase in vascular endothelial growth factor-induced angiogenesis and vascular permeability. *Proc Natl Acad Sci U S A* 2001;98:2604-2609.
 23. Hatakeyama T, Pappas PJ, Hobson RW, 2nd, Boric MP, Sessa WC, Duran WN.

- Endothelial nitric oxide synthase regulates microvascular hyperpermeability in vivo. *J Physiol* 2006;574:275-281.
24. Takano H, Liu W, Zhao Z, Cui S, Zhang W, Shibamoto T. N(G)-nitro-L-arginine methyl ester, but not methylene blue, attenuates anaphylactic hypotension in anesthetized mice. *J Pharmacol Sci* 2007;104:212-217.
 25. Duran WN, Breslin JW, Sanchez FA. The NO cascade, eNOS location, and microvascular permeability. *Cardiovasc Res* 2010;87:254-261.
 26. Cauwels A, Janssen B, Buys E, Sips P, Brouckaert P. Anaphylactic shock depends on PI3K and eNOS-derived NO. *J Clin Invest* 2006;116:2244-2251.
 27. Korhonen H, Fisslthaler B, Moers A, Wirth A, Habermehl D, Wieland T, Schutz G, Wettschureck N, Fleming I, Offermanns S. Anaphylactic shock depends on endothelial G_q/G₁₁. *J Exp Med* 2009;206:411-420.
 28. Igarashi J, Bernier SG, Michel T. Sphingosine 1-phosphate and activation of endothelial nitric-oxide synthase. Differential regulation of Akt and MAP kinase pathways by EDG and bradykinin receptors in vascular endothelial cells. *J Biol Chem* 2001;276:12420-12426.
 29. Ishii S, Kuwaki T, Nagase T, Maki K, Tashiro F, Sunaga S, Cao WH, Kume K, Fukuchi Y, Ikuta K, Miyazaki J, Kumada M, Shimizu T. Impaired anaphylactic responses with intact sensitivity to endotoxin in mice lacking a platelet-activating factor receptor. *J Exp Med* 1998;187:1779-1788.
 30. Lorenz JN, Arend LJ, Robitz R, Paul RJ, MacLennan AJ. Vascular dysfunction in S1P2 sphingosine 1-phosphate receptor knockout mice. *Am J Physiol Regul Integr Comp Physiol* 2007;292:R440-446.
 31. Thibeault S, Rautureau Y, Oubaha M, Faubert D, Wilkes BC, Delisle C,

- Gratton JP. S-nitrosylation of beta-catenin by eNOS-derived NO promotes VEGF-induced endothelial cell permeability. *Mol Cell* 2010;13;39(3):468-76.
32. Lucas KA, Pitari GM, Kazerounian S, Ruiz-Stewart I, Park J, Schulz S, Chepenik KP, Waldman SA. Guanylyl cyclases and signaling by cyclic GMP. *Pharmacol Rev* 2000;52:375-414.
33. Hess DT, Stamler JS. Regulation by s-nitrosylation of protein post-translational modification. *J Biol Chem* 2012;287:4411-4418.
34. Marin N, Zamorano P, Carrasco R, Mujica P, Gonzalez FG, Quezada C, Meininger CJ, Boric MP, Duran WN, Sanchez FA. S-nitrosation of beta-catenin and p120 catenin: A novel regulatory mechanism in endothelial hyperpermeability. *Circ Res* 2012;111:553-563.
35. Fleming I. Molecular mechanisms underlying the activation of eNOS. *Pflugers Arch* 2010;459:793-806.
36. Sugimoto N, Takuwa N, Okamoto H, Sakurada S, Takuwa Y. Inhibitory and stimulatory regulation of rac and cell motility by the G12/13-Rho and Gi pathways integrated downstream of a single G protein-coupled sphingosine-1-phosphate receptor isoform. *Mol Cell Biol* 2003;23:1534-1545.
37. Sanchez T, Thangada S, Wu MT, Kontos CD, Wu D, Wu H, Hla T. PTEN as an effector in the signaling of antimigratory G protein-coupled receptor. *Proc Natl Acad Sci U S A* 2005;102:4312-4317.
38. Sanchez T, Estrada-Hernandez T, Paik JH, Wu MT, Venkataraman K, Brinkmann V, Claffey K, Hla T. Phosphorylation and action of the immunomodulator FTY720 inhibits vascular endothelial cell growth factor-induced vascular permeability. *J Biol Chem* 2003;278:47281-47290.
39. Oo ML, Chang SH, Thangada S, Wu MT, Rezaul K, Blaho V, Hwang SI, Han

- DK, Hla T. Engagement of S1P1-degradative mechanisms leads to vascular leak in mice. *J Clin Invest* 2011;121:2290-2300.
40. Igarashi J, Michel T. The enigma of sphingosine 1-phosphate synthesis: A novel role for endothelial sphingosine kinases. *Circ Res* 2008;102:630-632.

Footnote

¹Disclosure of potential conflict of interest: None

²On leave from Center of Medical Functional Experiment, Yanbian University
College of Medicine, China

Figure Legends

Figure 1. *Slpr2* deficiency exacerbates vascular leakage in active anaphylaxis. Mice that had been sensitized with BSA were challenged with i.v. injection of BSA. (A and B) Evans blue leakage in lungs of mice 10 min after BSA challenge or saline infusion. (A) Representative lung photographs. (B) Quantification of Evans blue leakage in lung. (C) HCT 10 min after BSA challenge. (D) Survival of BSA-challenged WT and *Slpr2*^{-/-} mice. Numbers in parentheses denote numbers of mice or samples analyzed throughout the Figures. The following symbols are used throughout the Figures. * p < 0.05; ** p < 0.01; *** p < 0.001. If not marked, differences were not statistically significant.

Figure 2. *Slpr2* deficiency augments PAF injection-induced phosphorylation of Akt and eNOS in aorta and lung and production of cGMP in aorta. (A-D) PAF-stimulated phosphorylation of Akt at Ser⁴⁷³ and eNOS at Ser¹¹⁷⁷ in aorta (A and B) and lung (C and D) of WT and *Slpr2*^{-/-} mice that were i.v. injected with saline or PAF. (E) PAF injection-stimulated increase in cGMP level in aorta. Cyclic GMP was extracted and determined using an ELISA kit.

Figure 3. Genetic eNOS deficiency restores aggravation of vascular leakage after PAF injection and antigen challenge in *Slpr2* deficient mice. (A) HCT 10 min after PAF injection in WT and *Slpr2*^{-/-} mice with or without *Nos3* deficiency. (B) Survival after PAF injection of WT and *Slpr2*^{-/-} mice with or without *Nos3* deficiency. (C) HCT 10 min after BSA challenge in *Slpr2*^{+/+} and *Slpr2*^{-/-} mice with or without *Nos3* deficiency. (D) Changes in mean arterial blood pressure after BSA challenge in *Slpr2*^{+/+} and *Slpr2*^{-/-} mice with or without *Nos3* deficiency. (E) Survival after BSA

challenge of *Slpr2*^{+/+} and *Slpr2*^{-/-} mice with or without *Nos3* deficiency.

Figure 4. PAF- and S1P-stimulated phosphorylation of Akt and eNOS and production of NO are exaggerated in *Slpr2* deficient EC. (A and B) PAF (100 nM)- and S1P (100 nM)-stimulated phosphorylation of Akt at Ser⁴⁷³ (A) and eNOS at Ser¹¹⁷⁷ (B) in lung endothelial cells (MLEC) isolated from WT and *Slpr2*^{-/-} mice. Serum-starved MLEC were stimulated as indicated for 20 min. Top: representative blots. Bottom: quantified data. (C) PAF (50 nM)-, VEGF-A₁₆₅ (10 ng/ml)- and S1P (100 nM)-stimulated increase in NO production in MLEC. NO released into media for 3 h was determined as NO₂⁻ and NO₃⁻.

Figure 5. PAF-induced disruption of AJ is aggravated by *Slpr2* deficiency and rescued by a NOS inhibitor in EC. (A) Immunofluorescent staining using anti-VE-cadherin (green) or anti- β -catenin antibody (red) in MLEC. MLEC were stimulated or not with PAF (100 nM) for 30 min with or without L-NAME (1 mM) in the presence of 1% fetal bovine serum. Bars, 20 μ m. (B) PAF- and S1P-induced S-nitrosylation of β -catenin in MLEC. Serum-starved MLEC were pretreated or not with S1P (30 nM) plus L-NAME (1 mM) or S1P alone for 10 min and then stimulated with PAF (100 nM) for 30 min, followed by analysis using the biotin-switch method and western blotting with anti- β -catenin antibody. Left: representative blots. Right: quantified data of S-nitroso- β -catenin.




Figure 1




Figure 2




Figure 3




Figure 4





Figure 5




Figure E1







Figure E2

A**B****C****Figure E3**









Figure E4

A**B****Figure E5**

A**B****Figure E6**