

Syntheses of cis-Zeatin and Its
9-(2-Deoxy- β -D-ribofuranosyl) Derivative : A
Novel Synthetic Route to the Side Chain at C(6),
and Cytokinin Activity

メタデータ	言語: eng 出版者: 公開日: 2017-10-03 キーワード (Ja): キーワード (En): 作成者: メールアドレス: 所属:
URL	http://hdl.handle.net/2297/7665

Syntheses of *cis*-Zeatin and Its 9-(2-Deoxy- β -D-ribofuranosyl) Derivative: A Novel Synthetic Route to the Side Chain at C(6), and Cytokinin Activity

Antonio EVIDENTE,^{*a} Gennaro PICCIALI,^b Angelo SISTO,^c Masashi OHBA,^d Kei HONDA,^d and Tozo FUJII^{*d}

Dipartimento di Scienze Chimico-Agrarie, Università di Napoli "Federico II",^a 80055 Portici, Italy, Dipartimento di Chimica Organica e Biologica, Università di Napoli "Federico II",^b 80134 Napoli, Italy, Istituto Tossine e Micotossine da Parassiti Vegetali del CNR,^c 70126 Bari, Italy, and Faculty of Pharmaceutical Sciences, Kanazawa University,^d Takara-machi, Kanazawa 920, Japan. Received January 10, 1992

cis-Zeatin (3a) and its 9-(2-deoxy- β -D-ribofuranosyl) derivative (3c) have been synthesized from *N*-[(1,1-dimethylethoxy)carbonyl]glycine methyl ester (5) in 5 steps by adopting the " α -amino aldehyde/olefination" technology. The new *cis*-zeatin derivative (3c), its *trans* isomer (1c), and known *trans*-zeatin 9- β -D-ribose (1b) were tested for cytokinin activity in the stimulation of chlorophyll biosynthesis in etiolated cucumber cotyledons. The riboside 1b turned out to be the most active cytokinin among them, while the deoxyribosides 1c and 3c showed a marked decrease and a total loss, respectively, of cytokinin activity.

Keywords *cis*-zeatin synthesis; *cis*-zeatin 9-(2-deoxy- β -D-ribofuranosyl); *trans*-zeatin 9- β -D-ribofuranosyl; *trans*-zeatin 9-(2-deoxy- β -D-ribofuranosyl); glycine; α -amino aldehyde olefination; *Z* stereoselectivity; diisobutylaluminum hydride reduction; cytokinin activity; chlorophyll biosynthesis stimulation

In 1989, Evidente *et al.*¹⁾ reported the isolation of a new cytokinin from the culture filtrate of *Pseudomonas amygdali*, and its structure was established as 9-(2-deoxy- β -D-ribofuranosyl)-*trans*-zeatin (1c) by means of chemical synthesis.¹⁻³⁾ The natural occurrence of both *cis*-zeatin (3a) and *trans*-zeatin (1a) at the aglycone and the 9- β -D-ribofu-

ranoside (3b and 1b) levels⁴⁾ suggests that the *cis* isomer (3c) of the new cytokinin 1c may also occur in nature, and the availability of a synthetic reference sample would greatly facilitate the search for this *cis* isomer as a natural product. We therefore investigated the synthesis of 3c in the present work by adopting the " α -amino aldehyde/olefination" technology, which had been shown by us to be very effective for the construction of the side chains at C(6) of both enantiomers of *trans*-1'-methylzeatin (2a) and *cis*-1'-methylzeatin (4a), and of their 9- β -D-ribofuranosides (2b and 4b)^{5,6)} and 1c.²⁾

The synthetic route to 3c started with reduction of the *N*-protected glycine methyl ester 5 with diisobutylaluminum hydride (DIBAH) in CH₂Cl₂-hexane at -78 °C for 1 h, giving the aldehyde 6 in 67% yield. Olefination of 6 by the Still-Gennari modification⁷⁾ [(CF₃CH₂O)₂P(O)CH(Me)-CO₂Me, KN(SiMe₃)₂, 18-crown-6/MeCN, tetrahydrofuran (THF), -78 °C, 30 min] of the Horner-Wadsworth-Emmons reaction⁸⁾ produced the desired (*Z*)-ester 7 and the undesired (*E*)-ester 9 in 58% and 0.5% yields, respectively. On the other hand, Wittig reaction⁸⁾ of 6 with methyl 2-




Chart 1

TABLE I. Cytokinin Activity of 9-(2-Deoxy- β -D-ribofuranosyl)-*cis*-zeatin (3c), Its *trans* Isomer (1c), and *trans*-Zeatin 9- β -D-Riboside (1b) Tested by Cucumber Cotyledon Bioassay

Compound ^{a)}	Averaged chlorophyll yield ^{b)} (% of control)
Control	100.00 a
3c	108.87 aB
1c	164.91 C
1b	217.53 D

a) Solutions of cytokinins were 4.5 μ M and contained 0.1% methanol. b) Values scored by the same roman letters are not statistically significantly different at the 5% (lower case letters) or at the 1% (capital letters) level in Duncan's multiple range test.

(triphenylphosphoranylidene)propionate according to the literature procedure²⁾ but with some modification (CH₂Cl₂, room temp., 1 h) gave **7** and **9** in 6% and 74% yields, respectively. On reduction with DIBAH in CH₂Cl₂-hexane at -78 °C for 1 h, the (*Z*)-ester **7** furnished the allyl alcohol **8** in 85% yield. The carbamate **8** was then hydrolyzed with 10% aqueous HCl at room temperature for 1 h, and the resulting amino alcohol was isolated in the form of the oxalate **10** in 85% yield (from **8**).

Finally, coupling of **10** with 6-chloro-9-(2-deoxy- β -D-ribofuranosyl)purine in boiling 1-butanol containing Et₃N for 2.5 h afforded the desired deoxyriboside **3c** in 89% yield as a crystalline solid. In order to confirm the *Z* configuration of **10** and hence that of **3c**, **10** was purinylated with 6-chloropurine under similar conditions to give *cis*-zeatin (**3a**) in 72% yield. The new synthesis of **3a** from **5** through **6**, **7**, **8**, and **10** by taking advantage of the " α -amino aldehyde/olefination" technology represents a convenient alternative to previous synthetic methods.⁹⁻¹¹⁾

Thus, the total syntheses of natural 9-(2-deoxy- β -D-ribofuranosyl)-*trans*-zeatin (**1c**) and its *cis* isomer (**3c**) have made it possible to obtain sufficient amounts of **1c** and **3c** to test their cytokinin activity, compared to that of *trans*-zeatin 9- β -D-ribose (**1b**), in the stimulation of chlorophyll biosynthesis in etiolated cucumber cotyledons. The activity of the compounds, assayed at a concentration of 4.5 μ M, was then evaluated by using Duncan's statistic multiple range test. The results listed in Table I showed that **1c**, compared to **1b** which was the most active cytokinin among the compounds assayed, exhibited a marked decrease in the biosynthesis of chlorophyll, while **3c** was not statistically significantly different from the control. These findings indicate that, in general agreement with previously reported data,^{6b,12)} both the *trans* stereochemistry of the double bond of the side chain at C(6) and an unaltered ribosyl residue, when it is present, should be two structural features important to stimulating the biosynthesis of chlorophyll in cucumber cotyledons.

Experimental

General Notes All melting points were taken on a Yamato MP-1 capillary melting point apparatus and are corrected. Thin-layer chromatography (TLC) was conducted on Merck silica gel 60 F₂₅₄ plates (0.25-mm thickness), and spots were detected by means of ultraviolet (UV) absorbance measurement (at 254 nm) and/or by spraying with the standard KMnO₄ or I₂-KI reagent. Flash chromatography¹³⁾ was carried out by using Merck silica gel 60 (No. 9385). UV spectra were recorded on a Hitachi 320 UV spectrophotometer [on solutions in 95% (v/v) aqueous EtOH, 0.1 N aqueous HCl (pH 1), 0.005 M phosphate buffer (pH 7), and

0.1 N aqueous NaOH (pH 13)]. Infrared (IR) spectra were measured with a JASCO A-202 IR spectrophotometer. Mass spectra (MS) were obtained on a Hitachi M-80 mass spectrometer. ¹H- and ¹³C-nuclear magnetic resonance (NMR) spectra were recorded on either a JEOL JNM-FX-100 (¹H 100 MHz), a JEOL JNM-EX-270 (¹H 270 MHz, ¹³C 67.8 MHz), or a JEOL JNM-GSX-500 (¹H 500 MHz, ¹³C 125.65 MHz) instrument. Chemical shifts are reported in δ values relative to Me₄Si. Optical rotations were measured with a JASCO DIP-181 polarimeter using a 1-dm sample tube. Elemental analyses were performed by Mr. Y. Itatani and his associates at Kanazawa University. The following abbreviations are used: br = broad, d = doublet, dd = doublet-of-doublets, ddd = doublet-of-doublets-of-doublets, s = singlet, sh = shoulder, t = triplet, tq = triplet-of-quartets.

2-Oxoethylcarbamic Acid *tert*-Butyl Ester (6) A stirred solution of **5** (6.87 g, 36 mmol) in dry CH₂Cl₂ (120 ml) was cooled to -78 °C in an atmosphere of N₂, and a 0.95 M solution (76.5 ml, 73 mmol) of DIBAH in hexane was added dropwise over 10 min. After the mixture had been stirred at -78 °C for 1 h, the reaction was quenched by adding 2 N aqueous HCl (36 ml). The resulting mixture was neutralized with saturated aqueous NaHCO₃ and then stirred at room temperature for 1.5 h. The aqueous layer was separated from the organic layer and extracted with CH₂Cl₂. The CH₂Cl₂ extracts and the above organic layer were combined, washed with saturated aqueous NaCl, dried over anhydrous Na₂SO₄, and concentrated *in vacuo* to leave a colorless oil. The oil was purified by means of flash chromatography¹³⁾ [hexane-AcOEt (1:1, v/v)] to give **6**²⁾ (3.89 g, 67%) as a colorless oil, IR $\nu_{\text{max}}^{\text{CHCl}_3}$ cm⁻¹: 3450 (NH), 1709 (br, CO₂), ¹H-NMR (100 MHz, CDCl₃) δ : 1.46 (9H, s, CMe₃), 4.06 (2H, d, *J* = 5 Hz, CH₂), 5.16 (1H, br, NH), 9.65 (1H, s, CHO).

(Z)-4-[[[1,1-Dimethylethoxy]carbonyl]amino]-2-methyl-2-butenic Acid Methyl Ester (7) and (E)-4-[[[1,1-Dimethylethoxy]carbonyl]amino]-2-methyl-2-butenic Acid Methyl Ester (9) i) *Via* the Still-Gennari Modification of the Horner-Wadsworth-Emmons Reaction: A solution of 18-crown-6-MeCN complex¹⁵⁾ (8.66 g, 25 mmol) and methyl 2-[bis-(2,2,2-trifluoroethoxy)phosphinyl]propionate [(CF₃CH₂O)₂P(O)CH(Me)CO₂Me]¹⁷⁾ (1.83 g, 5.5 mmol) in dry THF (75 ml) was stirred at -78 °C in an atmosphere of N₂, and a 0.5 M solution (12 ml, 6.0 mmol) of potassium bis(trimethylsilyl)amide [KN(SiMe₃)₂] in toluene was added dropwise over 10 min. The mixture was stirred at -78 °C for 25 min, and then a solution of **6** (796 mg, 5.0 mmol) in dry THF (15 ml) was added dropwise over 5 min. After the resulting mixture had been stirred at the same temperature for a further 30 min, the reaction was quenched by adding saturated aqueous NH₄Cl (40 ml). The aqueous layer was separated from the organic layer and extracted with ether (3 \times 20 ml). The ethereal extracts and the above organic layer were combined, dried over anhydrous Na₂SO₄, and concentrated *in vacuo* to leave a pale yellow oil. The oil was purified by means of flash chromatography¹³⁾ [hexane-AcOEt (4:1, v/v)] afforded **7** (661 mg, 58%) as a colorless solid. Recrystallization of the solid from hexane yielded an analytical sample as colorless needles, mp 67-67.5 °C; IR $\nu_{\text{max}}^{\text{Nujol}}$ cm⁻¹: 3330 (NH), 1715 (α,β -unsaturated ester CO), 1685 (carbamate CO), 1645 (C=C); ¹H-NMR (500 MHz, CDCl₃) δ : 1.45 (9H, s, CMe₃), 1.91 (3H, d, *J* = 1.5 Hz, CH = CMe), 3.75 (3H, s, CO₂Me), 4.07 (2H, dd, *J* = 6 Hz each, NCH₂), 4.89 (1H, br, NH), 6.02 (1H, br, CH = CMe); ¹³C-NMR (67.8 MHz, CDCl₃) δ : 20.1 (CH = CMe), 28.4 (CMe₃), 39.6 (NCH₂), 51.6 (CO₂Me), 79.4 (CMe₃), 128.8 (CH = CMe), 140.6 (CH = CMe), 156.0 (CO₂CMe₃), 167.8 (CO₂Me). *Anal.* Calcd for C₁₁H₁₉NO₄: C, 57.63; H, 8.35; N, 6.11. Found: C, 57.87; H, 8.50; N, 6.16.

Later fractions in the above chromatography furnished **9**²⁾ (6 mg, 0.5%) as a colorless solid, mp 54-56 °C. This sample was identical (by comparison of the IR spectrum and TLC mobility) with the one obtained by method (ii) described below.

ii) *Via* the Wittig Reaction: A solution of methyl 2-(triphenylphosphoranylidene)propionate¹⁶⁾ (1.15 g, 3.3 mmol) in CH₂Cl₂ (3 ml) was added to a stirred solution of **6** (478 mg, 3.0 mmol) in CH₂Cl₂ (3 ml) over 10 min at 16-18 °C with occasional ice-cooling. The resulting mixture was stirred at room temperature for 1 h and then concentrated *in vacuo*. The yellow oily residue was extracted with hot hexane (5 \times 8 ml). The hexane extracts were combined and concentrated *in vacuo* to leave a slightly yellow solid. The solid was purified by means of flash chromatography¹³⁾ using hexane-AcOEt (4:1, v/v) as the eluent. Earlier fractions yielded the (*Z*)-isomer (**7**) (44 mg, 6%) as a colorless solid, mp 66-67 °C. This sample was identical (by comparison of the IR spectrum and TLC mobility) with the one prepared by method (i).

Later fractions obtained from the above chromatography gave the (*E*)-isomer (**9**)²⁾ (508 mg, 74%) as a colorless solid. Recrystallization of the solid from hexane afforded an analytical sample as colorless plates,

mp 57–58°C; IR $\nu_{\text{max}}^{\text{Nujol}}$ cm^{-1} : 3340 (NH), 1713 (α,β -unsaturated ester CO), 1685 (carbamate CO), 1655 (sh) (C=C); $^1\text{H-NMR}$ (500 MHz, CDCl_3) δ : 1.45 (9H, s, CMe_3), 1.87 (3H, d, $J=1$ Hz, $\text{CH}=\text{CMe}$), 3.74 (3H, s, CO_2Me), 3.90 (2H, br, NCH_2), 4.73 (1H, br, NH), 6.66 (1H, tq, $J=6.5$, 1 Hz, $\text{CH}=\text{CMe}$); $^{13}\text{C-NMR}$ (67.8 MHz, CDCl_3) δ : 12.6 ($\text{CH}=\text{CMe}$), 28.4 (CMe_3), 38.9 (NCH_2), 51.9 (CO_2Me), 79.7 (CMe_3), 129.2 ($\text{CH}=\text{CMe}$), 138.3 ($\text{CH}=\text{CMe}$), 155.8 (CO_2CMe_3), 168.0 (CO_2Me). *Anal.* Calcd for $\text{C}_{11}\text{H}_{19}\text{NO}_4$: C, 57.63; H, 8.35; N, 6.11. Found: C, 57.47; H, 8.58; N, 6.08.

(Z)-4-Hydroxy-3-methyl-2-butenyl)carbamic Acid tert-Butyl Ester (8) A stirred solution of **7** (459 mg, 2.0 mmol) in dry CH_2Cl_2 (5 ml) was cooled to -78°C in an atmosphere of N_2 , and a 0.94 M solution (6.4 ml, 6.0 mmol) of DIBAH in hexane was added dropwise over 10 min. After the mixture had been stirred at -78°C for 1 h, the reaction was quenched by adding a 5 M solution (4 ml) of AcOH in CH_2Cl_2 . The resulting mixture was allowed to warm to room temperature, and 10% aqueous tartaric acid (10 ml) and H_2O (10 ml) were added in that order. The aqueous layer was separated from the organic layer and extracted with CH_2Cl_2 (3×15 ml). The CH_2Cl_2 extracts and the above organic layer were combined, washed successively with saturated aqueous NaHCO_3 and saturated aqueous NaCl, dried over anhydrous Na_2SO_4 , and concentrated *in vacuo* to leave a colorless oil. Purification of the oil by means of flash chromatography¹³ [hexane–AcOEt (3:2, v/v)] gave **8** (342 mg, 85%) as a colorless oil, IR $\nu_{\text{max}}^{\text{film}}$ cm^{-1} : 3350 (NH and OH), 1690 (carbamate CO); $^1\text{H-NMR}$ (500 MHz, CDCl_3) δ : 1.43 (9H, s, CMe_3), 1.82 (3H, s, $\text{CH}=\text{CMe}$), 1.9 (1H, br, OH), 3.72 (2H, d, $J=7$ Hz, NCH_2), 4.12 (2H, s, CH_2OH), 4.74 (1H, br, NH), 5.27 (1H, t, $J=7$ Hz, $\text{CH}=\text{CMe}$).

(Z)-4-Amino-2-methyl-2-buten-1-ol Ethanediolate (2:1) (Salt) (10) A mixture of **8** (322 mg, 1.6 mmol) and 10% aqueous HCl (3.2 ml) was stirred at room temperature for 1 h, giving a slightly pinkish solution. The solution was passed through a column of Amberlite IRA-402 (HCO_3^-) (16 ml), and the column was eluted with H_2O . The eluate (75 ml) was concentrated to dryness *in vacuo* to leave a pale yellow oil (158 mg), which was dissolved in EtOH (1.5 ml). The resulting ethanolic solution was exactly neutralized by addition of a solution of oxalic acid (72 mg, 0.8 mmol) in EtOH (0.5 ml) and, if necessary, with Et_3N . The precipitate that resulted was filtered off, washed with EtOH (0.3 ml), and dried to give **10** (199 mg, 85%), mp 180–182°C. Recrystallization of crude **10** from EtOH– H_2O (10:1, v/v) afforded an analytical sample as colorless scales, mp 183.5–185°C; IR $\nu_{\text{max}}^{\text{max}}$ cm^{-1} : 3050 (sh) (OH), 1655 (C=C), 1580 (CO_2^-); $^1\text{H-NMR}$ (270 MHz, D_2O) δ : 1.73 (3H, s, Me), 3.67 (2H, d, $J=7.5$ Hz, NCH_2), 4.15 (2H, s, CH_2OH), 5.45 (1H, t, $J=7.5$ Hz, $\text{CH}=\text{CMe}$). *Anal.* Calcd for $\text{C}_{12}\text{H}_{24}\text{N}_2\text{O}_6$: C, 49.30; H, 8.27; N, 9.58. Found: C, 49.54; H, 8.46; N, 9.66.

(Z)-2-Methyl-4-(9H-purin-6-ylamino)-2-buten-1-ol (cis-Zeatin) (3a) A mixture of **10** (64 mg, 0.22 mmol), 6-chloropurine (62 mg, 0.4 mmol), Et_3N (0.5 ml), and 1-butanol (5 ml) was heated under reflux for 3 h. The reaction mixture was concentrated *in vacuo* to leave a pale yellow oil, which was dissolved in H_2O (1 ml). The resulting aqueous solution was passed through a column of Amberlite IRA-402 (HCO_3^-) (4 ml), and the column was eluted with H_2O (50 ml). The eluate was concentrated *in vacuo*, and the residual solid (82 mg) was recrystallized from EtOH (1 ml), giving a first crop (46 mg) of **3a**. A second crop (17 mg) was obtained by concentration of the mother liquor from the above recrystallization under reduced pressure and subsequent purification of the residue by flash chromatography¹³ [AcOEt–EtOH (4:1, v/v)]. The total yield of **3a** was 63 mg (72%). Further recrystallization of crude **3a** from EtOH yielded an analytical sample as colorless prisms, mp 212–214°C (lit. mp 206–208°C^{9a}); mp 209°C^{9b}); MS m/z : 219 (M^+); UV $\lambda_{\text{max}}^{95\% \text{ EtOH}}$ 269 nm (ϵ 19100); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 1) 274 (18000); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 7) 269 (19300); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 13) 275 (18600), 283 (sh) (14300); $^1\text{H-NMR}$ [500 MHz, (CD_3) $_2\text{SO}$] δ : 1.71 (3H, s, Me), 4.04 (2H, s, CH_2OH), 4.12 (2H, br, NCH_2), 4.73 (1H, br, OH), 5.35 (1H, t, $J=6.5$ Hz, $\text{CH}=\text{CMe}$), 7.60 (1H, br, NHCH_2), 8.08 and 8.17 (1H each, s, purine protons), 12.88 [1H, br, N(9)H]. *Anal.* Calcd for $\text{C}_{10}\text{H}_{13}\text{N}_5\text{O}$: C, 54.78; H, 5.98; N, 31.94. Found: C, 54.59; H, 6.04; N, 31.91. This sample was identical [by mixture melting point test and comparison of the IR (Nujol) and $^1\text{H-NMR}$ spectra and TLC mobility (in four solvent systems)] with authentic *cis*-zeatin (Sigma Chemical Co.).

(Z)-N-(4-Hydroxy-3-methyl-2-butenyl)-2'-deoxyadenosine [9-(2-Deoxy- β -D-ribofuranosyl)-*cis*-zeatin] (3c) A stirred solution of **10** (64 mg, 0.22 mmol) and 6-chloro-9-(2-deoxy- β -D-ribofuranosyl)purine¹⁸ (108 mg, 0.4 mmol) in 1-butanol (5 ml) containing Et_3N (0.5 ml) was heated under reflux for 2.5 h. The reaction mixture was concentrated *in vacuo* to leave a slightly yellow oil, which was dissolved in H_2O (1 ml). The aqueous solution was passed through a column of Amberlite IRA-402 (HCO_3^-)

(4 ml), and the column was eluted with H_2O (50 ml). The eluate was concentrated *in vacuo*, and the residual oil was purified by flash chromatography¹³ [AcOEt–EtOH (4:1, v/v)] to give **3c** (119 mg, 89%) as a colorless solid, mp 139–141°C. Recrystallization of the solid from MeCN–EtOH (10:1, v/v) yielded an analytical sample as colorless needles, mp 142.5–143.5°C; $[\alpha]_{\text{D}}^{22} -20.4^\circ$ ($c=0.322$, EtOH); MS m/z : 335 (M^+); UV $\lambda_{\text{max}}^{95\% \text{ EtOH}}$ 269 nm (ϵ 19700); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 1) unstable; $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 7) 269 (20500); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 13) 269 (20400); $^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 1.82 [3H, d, $J=1$ Hz, $\text{C}(5'')\text{-H's}$], 2.39 (1H, ddd, $J=13.5$, 6, 2.5 Hz) and 2.80 (1H, ddd, $J=13.5$, 8, 6 Hz) [$\text{C}(2')\text{-H's}$], 3.74 and 3.84 [1H each, dd, $J=12.5$, 3 Hz, $\text{C}(5')\text{-H's}$], 4.07 [1H, ddd, $J=3$ Hz each, $\text{C}(4')\text{-H}$], 4.20 [2H, s, $\text{C}(4'')\text{-H's}$], 4.25 [2H, br, $\text{C}(1'')\text{-H's}$], 4.57 [1H, ddd, $J=6$, 3, 2.5 Hz, $\text{C}(3')\text{-H}$], 5.47 [1H, tq, $J=7$, 1 Hz, $\text{C}(2'')\text{-H}$], 6.41 [1H, dd, $J=8$, 6 Hz, $\text{C}(1')\text{-H}$], 8.20 and 8.25 (1H each, s, purine protons)¹⁹; $^{13}\text{C-NMR}$ (125.65 MHz, CD_3OD) δ : 21.7 [$\text{C}(5'')$], 38.9 [$\text{C}(1'')$], 41.6 [$\text{C}(2'')$], 61.7 [$\text{C}(4'')$], 63.6 [$\text{C}(5')$], 73.0 [$\text{C}(3')$], 87.2 [$\text{C}(1')$], 89.9 [$\text{C}(4')$], 121.3 [$\text{C}(5)$], 124.3 [$\text{C}(2'')$], 140.0 [$\text{C}(3'')$], 140.9 [$\text{C}(8)$], 149.1 [$\text{C}(4)$], 153.3 [$\text{C}(2)$], 155.9 [$\text{C}(6)$].¹⁹ *Anal.* Calcd for $\text{C}_{15}\text{H}_{21}\text{N}_5\text{O}_4$: C, 53.72; H, 6.31; N, 20.88. Found: C, 53.48; H, 6.46; N, 20.85.

Bioassay Procedure The cytokinin activities (*i.e.*, stimulation of chlorophyll biosynthesis) of compounds **3c**, **1c**, and **1b** were tested in the cucumber cotyledon bioassay as described by Fletcher *et al.*,²⁰ with some minor modifications. The results are given in Table I.

Acknowledgment We thank the Italian National Research Council (CNR) for its financial support under the special *ad hoc* program 'Chimica Fine II', subproject 3.

References and Notes

- 1) A. Evidente, N. S. Iacobellis, R. Vellone, A. Sisto, and G. Surico, *Phytochemistry*, **28**, 2603 (1989).
- 2) L. De Napoli, A. Evidente, G. Piccialli, C. Santacroce, and R. Vellone, *Phytochemistry*, **29**, 701 (1990).
- 3) An enzymatic synthesis of this deoxyriboside from *trans*-zeatin and 2-deoxyribose 1-phosphate has recently been reported: Y. Inoue, F. Ling, and A. Kimura, *Agric. Biol. Chem.*, **55**, 629 (1991).
- 4) For a review, see D. S. Letham and L. M. S. Palni, *Annu. Rev. Plant Physiol.*, **34**, 163 (1983).
- 5) a) T. Itaya, T. Fujii, A. Evidente, G. Randazzo, G. Surico, and N. S. Iacobellis, *Tetrahedron Lett.*, **27**, 6349 (1986); b) T. Fujii, T. Itaya, and S. Matsubara, *Chem. Pharm. Bull.*, **37**, 1758 (1989); c) T. Fujii, T. Itaya, S. Yoshida, and S. Matsubara, *ibid.*, **37**, 3119 (1989).
- 6) a) T. Fujii, M. Ohba, and M. Sakari, *Heterocycles*, **27**, 2077 (1988); b) T. Fujii, M. Ohba, M. Sakari, and S. Matsubara, *Chem. Pharm. Bull.*, **38**, 2702 (1990), and references cited therein.
- 7) W. C. Still and C. Gennari, *Tetrahedron Lett.*, **24**, 4405 (1983).
- 8) For a review, see B. E. Maryanoff and A. B. Reitz, *Chem. Rev.*, **89**, 863 (1989).
- 9) a) N. J. Leonard, A. J. Playtis, F. Skoog, and R. Y. Schmitz, *J. Am. Chem. Soc.*, **93**, 3056 (1971); b) G. Mackenzie, P. W. Rugg, and G. Shaw, *J. Chem. Soc., Perkin Trans. 1*, **1976**, 1446.
- 10) a) G. Desvages and M. Olomucki, *Bull. Soc. Chim. Fr.*, **1969**, 3229; b) H. J. Vreman, R. Y. Schmitz, F. Skoog, A. J. Playtis, C. R. Frihart, and N. J. Leonard, *Phytochemistry*, **13**, 31 (1974).
- 11) For the synthesis at the 9- β -D-ribofuranoside level, see a) A. J. Playtis and N. J. Leonard, *Biochem. Biophys. Res. Commun.*, **45**, 1 (1971); b) S. David, G. de Sennyey, and B. Sotta, *Tetrahedron Lett.*, **23**, 1817 (1982).
- 12) A. Evidente, T. Fujii, N. S. Iacobellis, S. Riva, A. Sisto, and G. Surico, *Phytochemistry*, **30**, 3505 (1991), and references cited therein.
- 13) W. C. Still, M. Kahn, and A. Mitra, *J. Org. Chem.*, **43**, 2923 (1978).
- 14) R. Schwyzler, P. Sieber, and H. Kappeler, *Helv. Chim. Acta*, **42**, 2622 (1959).
- 15) a) G. W. Gokel, D. J. Cram, C. L. Liotta, H. P. Harris, and F. L. Cook, *J. Org. Chem.*, **39**, 2445 (1974); b) C. L. Liotta, U. S. Patent 3997562 (1976) [*Chem. Abstr.*, **86**, P 121387r (1977)].
- 16) O. Isler, H. Gutmann, M. Montavon, R. Rugg, G. Ryser, and P. Zeller, *Helv. Chim. Acta*, **40**, 1242 (1957).
- 17) Sodium 2,2-dimethyl-2-silapentane-5-sulfonate (DSS) was used as an internal standard.
- 18) M. J. Robins and G. L. Basom, *Can. J. Chem.*, **51**, 3161 (1973).
- 19) See formula **3c** in Chart I for the numbering system.
- 20) R. A. Fletcher, V. Kallidumbil, and P. Steele, *Plant Physiol.*, **69**, 675 (1982).