

# Usnic Acid. XVI. Alkaline Degradation of Dihydrous Usnic Acid

メタデータ	言語: eng 出版者: 公開日: 2017-10-03 キーワード (Ja): キーワード (En): 作成者: メールアドレス: 所属:
URL	<a href="http://hdl.handle.net/2297/7064">http://hdl.handle.net/2297/7064</a>

[Chem. Pharm. Bull.]  
28(5)1590—1596(1980)

## Usnic Acid. XVI.<sup>1)</sup> Alkaline Degradation of Dihydrousnic Acid

KŌTARO TAKAHASHI,<sup>2a)</sup> MASAKO TAKANI,<sup>2a)</sup> and YOSHIKAZU WADA<sup>2b)</sup>

Faculty of Pharmaceutical Sciences, University of Kanazawa<sup>2a)</sup> and Chemical Research Laboratories, Central Research Division, Takeda Chemical Industries Ltd<sup>2b)</sup>

(Received December 6, 1979)

The structure of a new alkaline degradation product of 1-dihydrousnic acid was elucidated as 5-acetyl-1,2,3,3a,7,7a-hexahydro-2,6-dihydroxy-1,2,3a-trimethyl-4-oxo-4*H*-inden-3,3-dicarboxylic acid (I) by chemical and spectral studies, together with X-ray analysis. The mechanism of formation of I from 1-dihydrousnic acid is discussed.

**Keywords**—dihydrousnic acid; alkaline degradation; 5-acetyl-1,2,3,3a,7,7a-hexahydro-2,6-dihydroxy-1,2,3a-trimethyl-4-oxo-4*H*-inden-3,3-dicarboxylic acid; PMR; CMR; MS; X-ray analysis

In a previous paper of this series,<sup>1)</sup> it was reported that usnic acid afforded, on alkaline degradation, 4,6-dihydroxy-3,5-dimethylcoumaran-2-one, in addition to usnetic and pyrousnic acids. This finding prompted us to investigate the alkaline degradation of dihydrousnic acid, since its degradation reactions are remarkably different from those of usnic acid.<sup>3)</sup> This paper deals with the structure elucidation of a new alkaline degradation product of dihydrousnic acid, and with the mechanism of its formation. 1-Dihydrousnic acid,  $[\alpha]_D^{20} -97^\circ$  ( $c=1.01$ , dioxane), was hydrolyzed with 50% (w/v) sodium hydroxide solution in an oxygen or a nitrogen stream to afford  $C_{16}H_{20}O_8$  (I), colorless plates of mp  $145^\circ$  (dec.),  $[\alpha]_D^{20} +75.7^\circ$  ( $c=1.03$ , dioxane). Compound I shows infrared (IR) absorption bands ( $cm^{-1}$ ) at 3520 (OH), 1720 (COOH), 1680 ( $\alpha,\beta$ -unsaturated C=O), 1600 (C=O chelated with enol OH) and 1550 (broad, characteristic of a triketone<sup>1)</sup>) and mass (MS) fragment ions at  $m/e$  296 ( $M^+-CO_2$ ) and  $m/e$  252 ( $M^+-2 \times CO_2$ ). Compound I was methylated with diazomethane to afford a dimethylate,  $C_{18}H_{24}O_8$  (II), colorless triangular crystals of mp  $76-78^\circ$ , and a monomethylate,  $C_{17}H_{22}O_8$  (III), colorless needles of mp  $154^\circ$ . Compound III showed an MS fragment ion at  $m/e$  310 ( $M^+-CO_2$ ) and  $^1H$  nuclear magnetic resonance (PMR) signals ( $\delta$ -value, ppm) at 2.51 (s, 3H), 18.14 (s, 1H), 1.62 (s, 3H), 2.89 (d, 2H), and 2.12 (d, t, 1H), which could be assigned to a chelated  $COCH_3$ , a chelated enol OH, an angular  $CH_3$ , and methylene and methine protons in a  $-CH_2-\dot{C}H-$  group, respectively, as shown in Table I.

Compound I was pyrolyzed at  $130-135^\circ$  *in vacuo* (3 mmHg) to afford  $C_{15}H_{18}O_6$  (IV), colorless needles of mp  $128-129^\circ$ , by decarboxylation and dehydration, and on oximation with hydroxylamine hydrochloride and pyridine, I afforded  $C_{15}H_{17}NO_4$  (V), colorless plates of mp  $187^\circ$  and  $C_{15}H_{18}N_2O_4$  (VI), colorless plates of mp  $228-230^\circ$ ; the former was proved to be identical with a mono-oxime monoanhydride of IV and the latter with a bisoxime monoanhydride of IV. Compound V afforded, on methylation with diazomethane, a monomethylate  $C_{16}H_{19}NO_4$  (VII), colorless plates of mp  $148-149^\circ$ , which showed IR bands at 1710 ( $COOCH_3$ ) and  $1675$  ( $\alpha,\beta$ -unsaturated C=O). These chemical and spectral findings suggest that ring B of dihydrousnic acid (Chart 2) remains unchanged in the molecule of compound I. Compound III showed PMR signals at 1.09 (d,  $J=6$  Hz, 3H), 1.87 (d, q,  $J=9$  and 6 Hz, 1H), 2.12 (d, t,

1) Part XV: K. Takahashi and M. Takani, *Chem. Pharm. Bull.*, 28, 177 (1980).

2) Location: a) 13-1 Takaramachi, Kanazawa, 920, Japan; b) 17-85 Jusohonmachi, Yodogawa-ku, Osaka, 532, Japan.

3) Y. Asahina and S. Shibata, "The Chemistry of Lichen Substances," Japan Society for the Promotion of Science, Tokyo, Japan, 1954, p. 190.

TABLE I. PMR Data ( $\delta$ -Value in  $\text{CDCl}_3$ , 100 MHz,  $J$  in Hz)


	$\text{C}_1\text{-CH}_3$	$\text{C}_{3a}\text{-CH}_3$	$\text{C}_2\text{-CH}_3$	$\text{C}_6\text{-}\overset{\text{X}^b}{\underset{\parallel}{\text{C}}}\text{-CH}_3$	$\text{C}_1\text{-H}$	$\text{C}_{7a}\text{-H}$	$\text{C}_7\text{-H}_2$	OH	OMe
I <sup>a)</sup>	1.09 d, 3H $J=6$	1.61 s, 3H	1.42 s, 3H	2.51 s, 3H	1.98 d,q, 1H $J=8.6$	2.10 d,t, 1H $J=8.8$	2.88 d, 2H $J=8$		
III	1.09 d, 3H $J=6$	1.62 s, 3H	1.32 s, 3H	2.51 s, 3H	1.87 d,q, 1H $J=9, 6$	2.12 d,t, 1H $J=9, 9$	2.89 d, 2H $J=9$	4.36 br, 1H 13.38 br, 1H 18.14 s, 1H	3.75 s, 1H
IV	1.13 d, 3H $J=7$	1.51 s, 3H	1.98 d, 3H $J=1.5$	2.62 s, 3H	2.46 d,q, 1H $J=11, 7$	1.91 d,d,d, 1H $J=11, 6, 2$	3.01 d,d, 1H $J=19, 6$ 2.76 d,d, 1H $J=19, 2$	12.96 br, 1H 18.63 s, 1H	
V	1.16 d, 3H $J=6$	1.53 s, 3H	1.95 d, 3H $J=1$	2.49 s, 3H	2.42 d,q, 1H $J=11, 6$	2.24 d,t, 1H $J=11, 3$	3.20 d, 2H $J=3$	12.60 br, 1H	
VI	1.11 d, 3H $J=6$	1.52 s, 3H	1.94 d, 3H $J=1$	2.57 s, 3H	2.31 d,q, 1H $J=11, 6$	2.09 d,t, 1H $J=11, 4$	3.04 d, 2H $J=4$		
VII	1.12 d, 3H $J=6$	1.58 s, 3H	1.96 d, 3H $J=1$	2.44 s, 3H	2.0-2.4 m, 2H		3.09 s-like 1H 3.13 d, 1H $J=3$		3.84 s, 3H
Dihydro- usinc acid		1.68 s, 3H $\text{C}_{9b}\text{-CH}_3$	1.98 s, 3H $\text{C}_8\text{-CH}_3$	2.56 s, 3H 2.63 s, 3H $2 \times \text{COCH}_3$		4.89 t, 1H $J=5$ $\text{C}_{4a}\text{-H}$	3.13 d, 2H $J=5$ $\text{C}_4\text{-H}_2$	9.53 s, 1H 13.37 s, 1H 18.28 s, 1H $3 \times \text{OH}$	

a) Some  $\text{CD}_3\text{OD}$  was added to dissolve the compound.

b)  $\text{X}=\text{O}$  for I, III, IV and  $\text{X}=\text{N}-\text{O}-$  for V, VI, VII.

Abbreviations: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; br, broad.

$J=9$  and 9 Hz, 1H) and 2.89 (d,  $J=9$  Hz, 2H), suggesting the presence of a  $\text{CH}_3\text{-}\dot{\text{C}}\text{H-}\dot{\text{C}}\text{H-CH}_2\text{-}$  system in the molecule, and also gave PMR signals at 1.32 (s, 3H), and 4.36 (broad, s, 1H), which suggest the presence of a  $-\dot{\text{C}}(\text{OH})\text{-CH}_3$  system, taking into consideration the observation that compound IV (decarboxy-dehydro derivative of I) shows a PMR signal at 1.98 (d,  $J=1.5$  Hz, 3H), which is assignable to methyl protons of a  $=\dot{\text{C}}\text{-CH}_3$  system, as shown in Table I. Compound III showed  $^{13}\text{C}$  NMR (CMR) signals ( $\delta$ -value, ppm) assignable to carbons of



a system, three methyl, one methylene and two methine groups, a  $\text{COOH}$ , a  $\text{COOCH}_3$  and three quaternary carbons, but III showed no signals assignable to aromatic carbons, as shown in Table II.

Based on the evidence described above, some structural features of the degradation product could be assumed, and X-ray analysis (Table III, Fig. 1 and Fig. 2) finally elucidated the structure of the degradation product as 5-acetyl-1,2,3,3a,7,7a-hexahydro-2,6-dihydroxy-1,2,3a-trimethyl-4-oxo-4*H*-inden-3,3-dicarboxylic acid (I) (final R-factor=4.6%) as shown in Chart

1. However, the absolute configuration still remains to be determined. Thus, compounds III, IV, V, VI, and VII could be formulated as shown in Chart 1, and their PMR and CMR spectra could be interpreted as shown in Table I and Table II, respectively.

The formation of I from dihydrousnic acid can be explained by the reaction mechanism postulated in Chart 2. In this mechanism, intermediate 1 derived from *Z*-dihydrousnic acid

TABLE II. CMR Data ( $\delta$  Value in  $\text{CDCl}_3$ , 25.15 MHz,  $J$  in Hz)

Carbon	Shielding <sup>a)</sup>				
	I (X=O) <sup>b)</sup>	III (X=O)	IV (X=O)	V (X=N-O-)	Dihydrousnic acid
C <sub>1</sub> -CH <sub>3</sub>	11.4 (q), [s] <sup>b)</sup>	11.4 (q, d) $J=127, 3$	15.6 (q, t), [s] <sup>e)</sup> $J=126, 4$	15.6 (q, t) $J=127, 4$	7.2 (q), [s] <sup>f)</sup> C <sub>8</sub> -CH <sub>3</sub>
C <sub>3a</sub> -CH <sub>3</sub>	30.5 (q), [s] <sup>e)</sup>	30.5 (q, d) $J=131, 8$	22.1 (q, d), [s] <sup>f)</sup> $J=129, 4$	21.8 (q, q) $J=129, 4$	23.3 (q), [s] <sup>f)</sup> C <sub>9b</sub> -CH <sub>3</sub>
X C <sub>5</sub> -C <sup>  </sup> -CH <sub>3</sub>	27.4 (q)	27.4 (q, d) $J=130, 3$	29.7 (q), [s] <sup>g)</sup> $J=130$	10.9 (q) $J=131$	31.2 (q) } 28.0 (q) } $2 \times \text{COCH}_3$
C <sub>2</sub> -CH <sub>3</sub>	23.5 (q), [s] <sup>d)</sup>	23.5 (q) $J=127$	14.1 (q), [s] <sup>h)</sup> $J=128$	14.4 (q) $J=129$	35.5 (t) C <sub>4</sub>
C <sub>3a</sub>	57.0 (s)	57.1 (q) $J=4$	56.0 (s)	57.2 (q) $J=4$	54.4 (s) C <sub>9b</sub>
C <sub>7</sub>	37.2 (t)	37.2 (t, t) $J=123, 3$	30.9 (t, t) $J=129, 3$	21.3 (t) $J=132$	85.8 (d) C <sub>4a</sub>
C <sub>1</sub>	46.4 (d)	46.1 (d, q) $J=132, 5$	45.9 (d, m) $J=128, 3$	45.5 (d, q) $J=132, 4$	101.8 (s) C <sub>6</sub>
C <sub>7a</sub>	49.8 (d)	49.7 (d) $J=131$	48.9 (d, q) $J=134, 4$	53.3 (d, m) $J=133, 4$	105.3 (s) C <sub>9a</sub>
C <sub>3</sub>	75.4 (s)	75.1 (s)	132.9 (q) $J=5$	131.9 (d) $J=4$	106.6 (s) C <sub>3</sub>
C <sub>2</sub>	83.1 (s)	83.2 (q) $J=3$	157.7 (m) $J=4$	158.1 (s) $J=4$	111.6 (s) C <sub>2</sub>
C <sub>5</sub>	111.3 (s)	111.0 (d) $J=6$	112.3 (s)	114.1 (q) $J=3$	158.6 (s) C <sub>5a</sub>
C <sub>4</sub>	198.6 (s)	198.6 (m) $J=4$	198.5 (s)	180.8 (t) $J=9$	159.3 (s) C <sub>9</sub>
C <sub>6</sub>	201.3 (s)	201.4 (m) $J=3$	201.6 (q) $J=4$	200.0 (q) $J=4$	163.5 (s) C <sub>7</sub>
X C <sub>5</sub> -C <sup>  </sup> -CH <sub>3</sub>	205.9 (s)	205.7 (q) $J=3$	204.5 (q) $J=6$	156.8 (q) $J=4$	195.5 (s) } 197.4 (s) } C <sub>1</sub> and C <sub>3</sub>
COOH and COOCH <sub>3</sub>	171.5 (s)	170.3 (s)	166.4 (s)	166.2 (s)	201.4 (s) } 202.8 (s) } $2 \times \text{COCH}_3$
COOCH <sub>3</sub>	172.4(s)	171.7(s) $J=4$ 53.2 (q) $J=149$			

a) ppm relative to TMS measured with complete decoupling. The letters in parentheses, in the cases of I and dihydrousnic acid, designate the multiplicity of the carbon signal with off-resonance decoupling and in the cases of III, IV, and V, the multiplicity without decoupling, unless otherwise indicated. Multiplicity with off-resonance decoupling was determined on irradiation at b) 1.09, c) 1.61, d) 1.42, e) 1.13, f) 1.51, g) 2.62, h) 1.98, i) 1.98, j) 1.68 by means of the proton selective decoupling technique (see Table I).

b) Some  $\text{CD}_3\text{OD}$  was added to dissolve the compound.

Abbreviations: s, s-like; d, d-like; t, t-like; q, q-like (others, see Table I).







Chart 2


## Experimental


The following instruments were used to obtain physical data. Melting point: Yanagimoto micro-melting apparatus (a hot plate type); UV (in EtOH): Hitachi 323 recording spectrometer; IR spectra (in KBr pellet): Nippon Bunko IR-G spectrometer; PMR: JNM-FX-100S instrument at 100 MHz (CMR at 25.15 MHz), with  $(\text{CH}_3)_4\text{Si}$  as an internal reference; MS spectra: JMS-OISG spectrometer [direct inlet: ionizing current, 150  $\mu\text{A}$ ; ionizing energy, 30 eV; source temperature, 140° (I), 115° (III), 95° (IV), 110° (V), 130° (VI) or 110° (VII)]. Thin-layer chromatography (TLC) was carried out on glass plates coated with silica gel G (Merck) and column chromatography with silica gel (Merck) and silicic acid (Mallinckrodt), unless otherwise stated.

**Hydrolysis of 1-Dihydrousnic Acid**—1-Dihydrousnic acid (8 g) in 50% (w/v) NaOH solution (48 ml) was hydrolyzed in an oxygen stream on a boiling water bath for 1 hr, then the hydrolysate was poured into ice-water, acidified with HCl and extracted with ethyl acetate. The ethyl acetate layer was evaporated down *in vacuo* to afford a resinous substance (7 g), which was column-chromatographed on silicic acid (350 g) with *n*-hexane-acetone (5:1). The fraction of  $R_f=0.14$  (TLC, silica gel impregnated with 0.1 N  $(\text{COOH})_2$ , *n*-hexane-acetone=5:1) was crystallized from petr-ether-ether to afford colorless plates (I) of mp 145° (dec.). Yield: 1.8 g. *Anal.* Calcd for  $\text{C}_{16}\text{H}_{20}\text{O}_8$ : C, 56.46; H, 5.92. Found: C, 56.46; H, 6.00.  $\text{FeCl}_3$  reaction: red-brown. IR ( $\text{cm}^{-1}$ ): 3500 (OH), 1720 (COOH), 1680 ( $\alpha,\beta$ -unsaturated C=O), 1800, 1550 (broad, triketone), 1460, 1220, 1180, 980, 925, 915, 780, 710. UV  $\lambda_{\text{max}}$  (nm, log  $\epsilon$ ): 242 (3.86), 274 (3.95). MS  $m/e$  (relative intensity %): 340 ( $\text{M}^+$ , 8), 322 ( $\text{M}^+ - \text{H}_2\text{O}$ , 1), 296 ( $\text{M}^+ - \text{CO}_2$ , 35), 278 (296  $-\text{H}_2\text{O}$ , 58), 260 (278  $-\text{H}_2\text{O}$ , 27), 252 (296  $-\text{CO}_2$ , 9), 245 (260  $-\text{CH}_3$ , 8), 237 (252  $-\text{CH}_3$ , 5), 234 (278  $-\text{CO}_2$ , 11), 219 (234  $-\text{CH}_3$ , 6), 216




The fraction of *Rf* 0.04 (TLC, silica gel impregnated with 0.1 N (COOH)<sub>2</sub>, *n*-hexane-acetone=5:1) was chromatographed on silica gel with CHCl<sub>3</sub>-acetone (5:1) and the resulting fraction of *Rf* 0.29 (TLC, CHCl<sub>3</sub>-acetone=5:1) (20 mg) was sublimed *in vacuo* at 70° to afford colorless needles (15 mg) of mp 106°; this material was identical with 4-methylresorcinol, obtained from 2,6-dihydroxy-3-methyl-acetophenone by hydrolysis with 50% NaOH (IR and mixed fusion). *Anal.* Calcd for C<sub>7</sub>H<sub>8</sub>O<sub>2</sub>: C, 67.73; H, 6.50. Found: C, 67.27; H, 6.40. Compound I was also obtained from 1-dihydrousnic acid by hydrolysis in a nitrogen stream.

**Methylation of I**—Compound I (0.5 g) in MeOH (30 ml) was methylated with diazomethane and the product was column-chromatographed on silicic acid (100 g) with benzene-acetone (30:1) to afford fractions of *Rf* 0.18 and 0.33 (TLC, silica gel impregnated with 0.1 N (COOH)<sub>2</sub>, benzene-acetone=50:1). The fraction of *Rf* 0.18 was chromatographed on silica gel (80 g) with *n*-hexane-acetone (5:1) and the resulting fraction of *Rf* 0.84 (TLC, *n*-hexane-acetone=5:1) was crystallized from petr-ether-ether to afford the dimethylate (II), colorless triangular crystals of mp 76–78°. *Anal.* Calcd for C<sub>18</sub>H<sub>24</sub>O<sub>8</sub>: C, 58.69; H, 6.57. Found: C, 58.37; H, 6.60. The fraction of *Rf* 0.33, one of the two fractions mentioned above, was further chromatographed on silicic acid (50 g) with benzene-AcOEt (50:1) and the resulting fraction of *Rf* 0.41 (TLC, silica gel impregnated with 0.1 N (COOH)<sub>2</sub>, benzene-AcOEt=50:1) was crystallized from petr-ether-ether to afford the monomethylate (III), colorless needles of mp 154°. Yield: 130 mg. *Anal.* Calcd for C<sub>17</sub>H<sub>22</sub>O<sub>8</sub>: C, 57.62; H, 6.55. Found: C, 57.73; H, 6.46. IR (cm<sup>-1</sup>): 3500 (OH), 1750 (COOCH<sub>3</sub>), 1715 (C=O), 1610 (C=O chelated with enol OH), 1540 (broad, triketone). UV λ<sub>max</sub> (nm, log ε): 274 (4.25). MS *m/e* (rel. intensity %): 354 (M<sup>+</sup>, 47), 336 (M<sup>+</sup> -H<sub>2</sub>O, 5), 310 (M<sup>+</sup> -CO<sub>2</sub>, 13), 292 (310 -H<sub>2</sub>O, 53), 260 (292 -CH<sub>3</sub>OH, 27), 232 (260 -CO, 12), 218 (260 -C<sub>2</sub>H<sub>2</sub>O, 16), 217 (260 -CH<sub>3</sub>CO, 13), 180 (CH<sub>3</sub>-C<sup>+</sup>-CH<sub>3</sub>, 43), 166 (A and/or ).




**Pyrolysis of I**—Compound I (0.5 g) was heated at 130–135° *in vacuo* (3 mmHg) for 40 min and when bubbling ceased, the resinous product (0.4 g) was chromatographed on silicic acid (200 g) with *n*-hexane-acetone (10:1). The fraction of *Rf* 0.2 (TLC, silica gel impregnated with 0.1 N (COOH)<sub>2</sub>, *n*-hexane-acetone=10:1) was crystallized from petr-ether-ether to afford IV, colorless needles of mp 128–129°. Yield: 110 mg. *Anal.* Calcd for C<sub>15</sub>H<sub>18</sub>O<sub>5</sub>: C, 64.73; H, 6.52. Found: C, 64.50; H, 6.43. IR (cm<sup>-1</sup>): 1700 (COOH), 1665 (α,β-unsaturated C=O), 1590 (C=O chelated with enol OH), 1545 (broad, triketone). UV λ<sub>max</sub> (nm, log ε): 225 (shoulder, 3.99), 272.5 (4.26). MS *m/e* (rel. intensity %): 278 (M<sup>+</sup>, 51), 260 (M<sup>+</sup> -H<sub>2</sub>O, 11), 245 (260 -CH<sub>3</sub>, 3), 234 (M<sup>+</sup> -CO<sub>2</sub>, 3), 218 (260 -C<sub>2</sub>H<sub>2</sub>O, 4), 166 (A, 25), 152 (B, 100), 148 (A -H<sub>2</sub>O, 12), 126 (C, 8), 124 (A -C<sub>2</sub>H<sub>2</sub>O, 9), 108 (B -CO<sub>2</sub>, 22), 107 (B -COOH, 26), 98 (C -CO, 9), 93 (108 -CH<sub>3</sub>, 28), 84 (C -C<sub>2</sub>H<sub>2</sub>O, 4).


**Oximation of I**—A mixture of I (0.5 g), NH<sub>3</sub>OH·HCl (0.5 g) in pyridine (0.5 ml) and EtOH (5 ml) was refluxed on a water bath for 5 hr and the reaction mixture, after removal of the solvent by evaporation, was acidified with dil. HCl and extracted with AcOEt. The ethyl acetate-soluble fraction afforded an oil (430 mg), which was chromatographed on silicic acid (100 g) with CHCl<sub>3</sub>-AcOEt (20:1). The fraction of *Rf* 0.53 (TLC, silica gel impregnated with 0.1 N (COOH)<sub>2</sub>, CHCl<sub>3</sub>-AcOEt=20:1) (70 mg) was crystallized from petr-ether-ether to afford V, colorless plates of mp 187°. This oxime was identical with the mono-oxime monoanhydride of IV (mixed fusion and IR). *Anal.* Calcd for C<sub>18</sub>H<sub>17</sub>NO<sub>4</sub>: C, 65.44; H, 6.22; N, 5.09. Found: C, 65.41; H, 6.05; N, 5.31. MS *m/e* (rel. intensity %): 275 (M<sup>+</sup>, 100), 257 (M<sup>+</sup> -H<sub>2</sub>O, 20), 231 (M<sup>+</sup>



The fraction of *Rf* 0.10 (TLC, silica gel impregnated with 0.1 N (COOH)<sub>2</sub>, CHCl<sub>3</sub>-AcOEt=20:1) (60 mg) was further chromatographed on silica gel with CHCl<sub>3</sub>-EtOH (20:1) and the resulting fraction of *Rf* 0.36 (10 mg) (TLC, CHCl<sub>3</sub>-EtOH=20:1) afforded VI, colorless plates of mp 228–230°. *Anal.* Calcd for C<sub>15</sub>H<sub>16</sub>N<sub>2</sub>O<sub>4</sub>: C, 62.05; H, 6.25; N, 9.65. Found: C, 62.13; H, 6.33; N, 9.59. MS *m/e* (rel. intensity %): 290 (M<sup>+</sup>,

89), 272 ( $M^+ - H_2O$ , 65), 255 (272 - OH, 100), 229 (255 -  $C\equiv N$ , 42), 152 (B, 11), 135 ( , 18).

**Methylation of V**—Compound V (80 mg) in MeOH was methylated with diazomethane and the product was chromatographed on silica gel (80 g) with benzene-acetone (50:1). The fraction of  $R_f$  0.27 (TLC, benzene-acetone=50:1) (50 mg) was crystallized from petr-ether-ether to afford VII, colorless plates of mp 148–149°. *Anal.* Calcd for  $C_{16}H_{19}NO_4$ : C, 66.42; H, 6.62; N, 4.84. Found: C, 66.36; H, 6.56; N, 5.12. IR ( $cm^{-1}$ ): 1710 ( $COOCH_3$ ), 1675 ( $\alpha,\beta$ -unsaturated  $C=O$ ). MS  $m/e$  (rel. intensity %): 289 ( $M^+$ , 100), 258 ( $M^+ - OCH_3$ , 44), 257 ( $M^+ - CH_3OH$ , 57), 230 ( $M^+ - COOCH_3$ , 28), 166 (D, 40), 165 (D - H, 75), 163

( , 4), 134 (D- $CH_3OH$ , 16), 133 (48), 126 (8), 125 (79), 107 (D- $COOCH_3$ , 25).

**Oximation of IV**—Compound IV (80 mg) in EtOH (2 ml) was oximated with  $NH_2OH \cdot HCl$  (50 mg) and pyridine (0.5 ml) and the product was chromatographed on silicic acid (35 g). The resulting fraction of  $R_f$  0.53 (TLC, silica gel impregnated with 0.1 N ( $COOH$ )<sub>2</sub>,  $CHCl_3$ -AcOEt=20:1) (30 mg) was crystallized from petr-ether-ether to afford colorless needles of mp 187°; this material was identical with V (mixed fusion and IR). *Anal.* Calcd for  $C_{15}H_{17}NO_4$ : C, 65.44; H, 6.22; N, 5.09. Found: C, 65.12; H, 6.21; N, 5.27. The fraction of  $R_f$  0.10 (TLC, silica gel impregnated with 0.1 N ( $COOH$ )<sub>2</sub>,  $CHCl_3$ -AcOEt=20:1) was further chromatographed on silica gel with  $CHCl_3$ -EtOH (20:1) and the resulting fraction of  $R_f$  0.36 (TLC,  $CHCl_3$ -EtOH=20:1) afforded VI, mp 228–230° (mixed fusion and TLC).

**Acknowledgement** The authors (K.T and M.T) are grateful to Prof. T. Fujii of this faculty for valuable discussion on the reaction mechanism, and to Drs. H. Morimoto and M. Nishikawa, Central Research Division, Takeda Chemical Industries Co. Ltd. for their kind advice. Thanks are also due to Mr. Y. Itatanl for NMR measurements, to Miss Y. Arano for elemental analyses, to Miss K. Ohata for MS measurements and to Mr. H. Ogata of Wako Junyaku Co. Ltd. for providing usnic acid.