

Auxiliary role for D-alanylated wall teichoic acid in Toll-like receptor 2-mediated survival of *Staphylococcus aureus* in macrophages

メタデータ	言語: eng 出版者: 公開日: 2017-10-04 キーワード (Ja): キーワード (En): 作成者: メールアドレス: 所属:
URL	http://hdl.handle.net/2297/20345

Auxiliary role for D-alanylated wall teichoic acid in Toll-like receptor 2-mediated survival of Staphylococcus aureus in macrophages

Journal:	<i>Immunology</i>
Manuscript ID:	IMM-2009-1362.R1
Manuscript Type:	Original Article
Date Submitted by the Author:	
Complete List of Authors:	Shiratsuchi, Akiko; Kanazawa University, Graduate School of Medical Science; Kanazawa University, Graduate School of Natural Science and Technology Shimizu, Kaori; Kanazawa University, Graduate School of Natural Science and Technology Watanabe, Ikuko; Kanazawa University, Graduate School of Medical Science Hashimoto, Yumi; Kanazawa University, Graduate School of Medical Science Kurokawa, Kenji; The University of Tokyo, Graduate School of Pharmaceutical Sciences; Pusan National University, National Research Laboratory of Defense Proteins, College of Pharmacy Razanajatovo, Iony; The University of Tokyo, Graduate School of Pharmaceutical Sciences Park, Keun Hwa; Pusan National University, National Research Laboratory of Defense Proteins, College of Pharmacy Park, Hae Kyung; Pusan National University, National Research Laboratory of Defense Proteins, College of Pharmacy Lee, Bok Luel; Pusan National University, National Research Laboratory of Defense Proteins, College of Pharmacy Sekimizu, Kazuhisa; The University of Tokyo, Graduate School of Pharmaceutical Sciences Nakanishi, Yoshinobu; Kanazawa University, Graduate School of Medical Science; Kanazawa University, Graduate School of Natural Science and Technology
Key Words:	Phagocytosis, macrophages / Monocytes, BACTERIA/BACTERIAL IMMUNITY

Summary

We previously reported that *Staphylococcus aureus* avoids killing in macrophages by exploiting the action of Toll-like receptor (TLR) 2 that leads to the c-Jun N-terminal kinase (JNK)-mediated inhibition of superoxide production. To search for bacterial components responsible for this event, a series of *S. aureus* mutants, in which the synthesis of cell wall was interrupted, were screened for the level of JNK activation in macrophages. Besides a mutant lacking the putative TLR2 ligand, lipoproteins, two mutant strains were found to activate the phosphorylation of JNK to a lesser extent than the parental strain, and this defect was recovered by acquisition of the corresponding wild-type genes. Macrophages that had phagocytosed the mutant strains produced more superoxide than those engulfing the parental strain, and the mutant bacteria were more efficiently killed in macrophages than the parent. The genes mutated, *dltA* and *tagO*, encoded proteins involved in the synthesis of D-alanylated wall teichoic acid. Unlike a cell wall fraction rich in lipoproteins, D-alanine-bound wall teichoic acid purified from the parent strain by itself did not activate JNK phosphorylation in macrophages. These results suggested that D-alanylated wall teichoic acid of *S. aureus* modulates the cell wall milieu for lipoproteins so that they effectively serve as a ligand for TLR2.

Keywords: bacterial infection; macrophages; phagocytosis; teichoic acids; Toll-like receptor

2

Introduction

Invading microbial pathogens compete with host organisms in the regulation of innate immunity.¹⁻⁵ They try to circumvent host immune responses to achieve an effective infection and prolonged survival through, for example, inhibition of signaling pathways for the activation of nuclear factor (NF)- κ B and mitogen-activated protein kinases, which induce the transcription of genes coding for antimicrobial substances and pro-inflammatory cytokines.¹⁻³ Some bacteria evade phagocytosis by immune cells or do not even concede once phagocytosed: they inhibit phagosomal maturation or escape from phagosomes to avoid digestion by lysosomal enzymes.⁶ To overcome such microbial actions against immune responses, host immune cells adopt an alternative strategy, as exemplified by the induction of autophagy, in which cytoplasmic bacteria are resealed with membranes and subjected to lysis.⁷⁻⁹ It is important to clarify the mechanism underlying the combat between microbial pathogens and host organisms for developing novel and effective medicines against infectious diseases.

We previously reported that *Staphylococcus aureus* inhibits the production of superoxide in macrophages to evade killing after phagocytosis, through Toll-like receptor (TLR) 2-mediated phosphorylation of c-Jun N-terminal kinase (JNK).¹⁰ This is unique in that bacteria exploit the actions of innate immune receptors that recognize microbial pathogens, TLR2 in this case, for their survival in macrophages. Presumably, TLR2 is activated by a component(s) of *S. aureus* residing at the cell wall, [such as lipoproteins and lipopeptides¹¹⁻¹⁷ with some controversies¹⁸](#), to transmit a signal leading to the phosphorylation of JNK and subsequent inhibition of superoxide production in macrophages. In the present study, we took a genetic approach to search for additional bacterial components required for the exploitation of TLR2 by *S. aureus* and obtained evidence that genes responsible for the

1
2
3 synthesis of D-alanylated wall teichoic acid (WTA) play a crucial role.
4
5
6
7

8 **Materials and methods**

9 *Materials*

10
11
12
13
14
15
16 An antibody (#9251) specifically recognizing the phosphorylated form of JNK and another
17
18 (#9252) recognizing both the phosphorylated and unphosphorylated forms were purchased
19
20 from Cell Signaling Technology (Beverly, MA). Using these antibodies, two isoforms of
21
22 JNK with M_r of 46 and 54 kDa, and their phosphorylated forms were detectable.
23
24 pHY300PLK, an *Escherichia coli*-*S. aureus* shuttle vector containing a tetracycline-resistant
25
26 gene, was obtained from Takara-Bio (Ohtsu, Japan). Fluorescein isothiocyanate was
27
28 purchased from Molecular Probes (Eugene, OR); the synthetic lipopeptide
29
30 tripalmitoyl-*S*-glycerylcysteine (Pam3Cys), lipopolysaccharide (LPS) from *Salmonella*
31
32 *enteritidis*, and *N*-acetyl-L-cysteine were from Sigma-Aldrich (St. Louis, MO); mannitol salt
33
34 agar medium was from Nissui (Tokyo, Japan); Diogenes was from National Diagnostics
35
36 (Atlanta, GA); and the Dual Luciferase Assay kit was from Promega Corp. (Madison, WI).
37
38
39
40
41
42
43

44 *Generation of bacterial strains, bacterial culture, and preparation and maintenance of* 45 46 *animal cells*

47
48
49 Cell surface mutants of *S. aureus* are derivatives of the parental wild-type *S. aureus* strain
50
51 RN4220 (a derivative of NCTC8325-4, a restriction and *agr* mutant¹⁹) (Table 1). To
52
53 construct the mutant strains M0614 and M0615, sequences corresponding to portions of the
54
55 *SA0614* and *SA0615* genes (nucleotide positions 50–400 and 32–507, respectively, with the
56
57 first nucleotide of the translation start codon numbered as 1) were amplified by PCR and
58
59 inserted into the *S. aureus* integration vector pSF151.²⁰ RN4220 was then transformed with
60

1
2
3 the resulting plasmids pSFSA0614 and pSFSA0615, and M0614 and M0615 where the
4 cognate genes had been disrupted by homologous recombination were selected. RN4220
5
6 and all the mutant strains were grown with Luria-Bertani medium at 37°C (except for M0702
7
8 that was grown at 30°C) to full growth, washed once with PBS, and used in the subsequent
9
10 experiments. Macrophages from the peritoneal cavity of thioglycollate-injected C57BL/6
11
12 mice were prepared and maintained with RPMI1640 medium supplemented with 10% (v/v)
13
14 heat-inactivated fetal bovine serum at 37°C with 5% (v/v) CO₂ in air.²¹ Mice carrying
15
16 disrupted *tlr2* in a C57BL/6 background²² were provided by Dr. Shizuo Akira of Osaka
17
18 University. HEK293 cells were cultured in DMEM supplemented with 10% heat-inactivated
19
20 fetal bovine serum at 37°C with 5% CO₂ in air. All the experiments involving animals were
21
22 conducted according to protocols that had been approved by the Committee on Animal
23
24 Experimentation of Kanazawa University.
25
26
27
28
29
30
31
32
33

34 *Preparation of WTA and lipoproteins*

35
36 WTA of *S. aureus* that retained D-alanine was prepared as described below. Bacteria were
37
38 disrupted using glass beads and centrifuged at 800 × g for 10 min. The supernatants were
39
40 re-centrifuged at 20,000 × g for 10 min, and the precipitates were suspended with 20 mM
41
42 sodium citrate (pH4.7) containing 0.5% (w/v) SDS, heated at 60°C for 30 min, and centrifuged
43
44 at 20,000 × g for 10 min. The precipitates were suspended with 5% (w/v) trichloroacetic
45
46 acid, kept at room temperature for 18 h, and centrifuged at 20,000 × g for 10 min. The
47
48 supernatants were mixed with acetone, and the resulting precipitates were dissolved with
49
50 water and centrifuged as above. The final supernatants were collected as purified WTA.
51
52 The purity of this WTA preparation was determined based on the amount of phosphorus
53
54 contained in a given dry weight as well as by PAGE followed by staining with silver,
55
56 according to standard procedures.^{23,24} To examine the attachment of D-alanine, the WTA
57
58
59
60

1
2
3 preparation was incubated with 0.1 M NaOH at 37°C for 2 h and separated in thin-layer
4 chromatography on Silica-gel 60 with a solvent of *n*-propanol : pyridine : acetic acid : water =
5 18 : 10 : 5 : 16, and the developed plate was treated with the ninhydrin reagent to visualize
6 amino groups. A fraction rich in lipoproteins was prepared by the Triton X-114
7 phase-partitioning method, as described previously.¹⁴ Briefly, cell lysates were treated with
8 Triton X-114 (2% (v/v)) and centrifuged at 10,000 × *g* for 10 min at 37°C, and materials in
9 the Triton X-114 phase were precipitated with ethanol, dissolved with water, and used as the
10 lipoprotein-rich fraction.
11
12
13
14
15
16
17
18
19
20
21
22
23
24

25 *Assays for JNK phosphorylation, phagocytosis of S. aureus, colony formation of engulfed S.*
26 *aureus, superoxide production, and lysosomal enzyme*

27
28
29 The level of phosphorylated JNK was determined by Western blotting as described.¹⁰ In
30 brief, mouse peritoneal macrophages from either wild-type or *tlr2*-deficient mice were
31 incubated with *S. aureus* (macrophages : bacteria = 1 : 5, except for wild-type macrophages
32 with *tagO* and *lgt* mutants at a ratio of 1 : 10) or cell wall components at 37°C and lysed in a
33 buffer containing SDS and inhibitors of phosphatases and proteases, and the lysates were
34 subjected to SDS-PAGE. The separated proteins were transferred to polyvinylidene
35 difluoride membranes and reacted with antibodies, and specific signals were visualized by a
36 chemiluminescence reaction and processed using Fluor-S MultiImager (Bio-Rad, Hercules,
37 CA). Phagocytosis reactions with peritoneal macrophages and fluorescein
38 isothiocyanate-labeled *S. aureus* as the phagocytes and targets (macrophages : bacteria = 1 :
39 10), respectively, were carried out as described previously.¹⁰ After incubation, the
40 macrophage cultures were washed with PBS to remove unengulfed bacteria, and macrophages
41 attached to culture containers were fixed and treated with trypan blue for quenching
42 fluorescence from bacteria remaining attached to macrophages. The samples were then
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 examined by phase-contrast and fluorescence microscopy for the level of phagocytosis. To
4
5 determine the values of colony-forming units of engulfed *S. aureus*, macrophages incubated
6
7 with bacteria (macrophages : bacteria = 1 : 500) for 30 min were washed to remove
8
9 unengulfed bacteria and further incubated for 30 min. The macrophages 0 and 30 min after
10
11 the wash were lysed with water, and the lysates at serial dilutions were seeded on
12
13 agar-solidified mannitol salt medium or Luria-Bertani medium, the latter of which contained
14
15 tetracycline and was used for bacteria transformed with pHY300PLK-based plasmid. The
16
17 plates were incubated overnight at 37°C, and the number of colonies (only those surrounded
18
19 by yellow rings with mannitol salt medium) was determined and presented relative to that
20
21 obtained at time 0 after the wash. For the examination of superoxide production,
22
23 macrophages maintained on coverslips with serum-free RPMI1640 medium were incubated
24
25 with unlabeled bacteria (macrophages : bacteria = 1 : 1000) at 37°C, and the amount of
26
27 superoxide released into the culture medium was determined by a chemiluminescence
28
29 reaction using Diogenes, as described previously.¹⁰ For determining the activity of
30
31 α -N-acetylglucosaminidase, whole-cell lysates of peritoneal macrophages were incubated
32
33 with a reaction mixture containing 4-methylumbelliferyl N-acetyl- α -D-glucosaminide
34
35 (Sigma-Aldrich), and the level of cleaved substrates was measured with a fluorometer, as
36
37 described previously.²⁵
38
39
40
41
42
43
44
45
46
47

48 *Assay for NF- κ B activation*

49
50
51 HEK293 cells were transfected by the calcium/phosphate method overnight with a mixture of
52
53 plasmid DNA including pELAM²⁶ (a gift from Dr. Douglas Golenbock at University of
54
55 Massachusetts), a reporter gene vector expressing firefly luciferase under the control of a
56
57 promoter activated by NF- κ B; pRL-TK (Promega Corp.), a control reporter constitutively
58
59 expressing Renilla luciferase used for the normalization of transfection efficiency; and mouse
60

1
2
3 TLR2 cDNA in pDisplay (Invitrogen, Carlsbad, CA) (a gift from Dr. Yoshiyuki Adachi at
4
5 Tokyo University of Pharmacy and Life Science).²⁷ The cells were further cultured with
6
7 fresh medium for a day and subsequently incubated with *S. aureus* for 2 h, and the cell lysates
8
9 were examined for the amounts of firefly luciferase and Renilla luciferase using the Dual
10
11 Luciferase Assay kit. The ratio of firefly luciferase to Renilla luciferase was determined and
12
13 considered as the level of NF- κ B activation.
14
15
16
17

18 19 20 *Statistical analysis*

21
22 Data are representative of at least 3 independent experiments ($n = 2-3$ in each experiment)
23
24 that yielded similar results. Data from quantitative analyses are expressed as the mean \pm
25
26 standard deviations of the results from at least 3 independent experiments. Statistical
27
28 analyses were performed using Student's *t*-test, and *P* values of less than 0.05 were
29
30 considered significant. The data significantly different from controls were marked with
31
32 asterisks.
33
34
35
36
37

38 39 **Results**

40 41 42 43 **Inefficient JNK phosphorylation in macrophages after incubation with *S. aureus* lacking** 44 45 **expression of *dltA* or *tagO***

46
47 To search for components of *S. aureus* responsible for the activation of TLR2-mediated
48
49 phosphorylation of JNK in macrophages, we screened a series of *S. aureus* strains with
50
51 mutations that affect the structure of cell wall (Table 1). Peritoneal macrophages from
52
53 thioglycollate-injected mice were incubated with either the parental strain RN4220 or its
54
55 mutant strains, and whole cell lysates were subjected to Western blotting to determine the
56
57 level of the phosphorylated form of JNK. Macrophages showed a rise in the level of
58
59
60

1
2
3 phosphorylated JNK 10 min after the incubation with RN4220, and the increase continued
4
5 during the next 20 min (left panel in Fig. 1a), as we reported previously.¹⁰ Incubation with a
6
7 mutant strain lacking the expression of *dltA* similarly brought about the activation of JNK
8
9 phosphorylation, but the level was much lower than that observed with the parental strain (left
10
11 panel in Fig. 1a). This effect was not due to impaired phagocytosis of the mutant bacteria by
12
13 macrophages because the parental and mutant strains were comparable in their susceptibility
14
15 to phagocytosis (right panels in Fig. 1a). The level of phosphorylated JNK was lower in
16
17 macrophages incubated with the strain T013 (Fig. 1b), in which the *lgt* gene coding for
18
19 lipoprotein diacylglycerol transferase is disrupted.¹⁴ This mutant strain is devoid of lipid
20
21 modification of all lipoproteins at the cell surface, and the result was in line with previous
22
23 reports that lipoproteins serve as a ligand for TLR2. Similar reduction in the level of JNK
24
25 phosphorylation was seen when macrophages were incubated with a *tagO*-deficient strain and,
26
27 though less significantly, mutants for the genes *SA0614* or *SA0615* (Fig. 1b). The other
28
29 mutant strains including one deficient in the *ltaS* gene, which codes for
30
31 polyglycerolphosphate synthase of lipoteichoic acid (LTA), did not differ from the parental
32
33 strain as to the effect on the phosphorylation of JNK in macrophages (Fig. 1b). When
34
35 macrophages were incubated with the *dltA* mutant that had been introduced with a plasmid
36
37 expressing the *dltABCD* operon, the level of phosphorylated JNK became almost equal to that
38
39 in macrophages incubated with the parental strain (left panel in Fig. 1c). Similarly, the
40
41 expression of *tagO* in the *tagO* mutant complemented a defect in the phosphorylation of JNK
42
43 (right panel in Fig. 1c). These results confirmed the importance of *dltA* and *tagO* for *S.*
44
45 *aureus* to induce JNK phosphorylation in macrophages.
46
47
48
49
50
51
52
53
54
55
56
57

58 **Auxiliary action of D-alanylated teichoic acid in the activation of TLR2 by *S. aureus***

59 Unlike TLR4-acting LPS, the parent and mutant strains deficient in *dltA* or *tagO* did not seem
60 to activate macrophages lacking the expression of TLR2 in terms of the induction of JNK

1
2
3 phosphorylation (Fig. 2a). This indicated that the *S. aureus*-activated phosphorylation of
4
5 JNK depends on the action of TLR2. In fact, a fraction rich in lipoproteins, a putative ligand
6
7 for TLR2, obtained from the parent strain, the *dltA* mutant, and the *tagO* mutant, but not from
8
9 a mutant strain lacking the expression of *lgt*, induced the phosphorylation of JNK in wild-type
10
11 macrophages (Fig. 2b).
12
13

14
15 The *dltA* gene codes for one of the proteins responsible for the D-alanylation of
16
17 teichoic acids,²⁸ and *tagO* codes for an enzyme responsible for the transfer of
18
19 *N*-acetylglucosamine phosphate to the lipid carrier,²⁸ an essential step in the synthesis of
20
21 WTA. On the other hand, *SA0614* and *SA0615* code for proteins that compose a
22
23 two-component system of *S. aureus*, which induces the expression of the *dltABCD*
24
25 operon.^{29,30} Considering the results with mutant strains deficient in these genes as well as
26
27 one with the *ltaS* mutant lacking LTA, we anticipated that D-alanylated WTA is required for
28
29 the TLR2-mediated phosphorylation of JNK in macrophages. To more directly determine
30
31 the role of WTA, we isolated WTA from *S. aureus* strains and examined its action. The
32
33 WTA preparation was considered to be nearly pure based on the content of phosphorus and
34
35 the staining pattern in PAGE (left and middle panels in Fig. 2c): note that none was obtained
36
37 from the *tagO* mutant lacking an enzyme essential for WTA synthesis, and that a difference in
38
39 the migration of WTA prepared from the *dltA* mutant was probably due to a lack of D-alanine.
40
41 In fact, WTA of the *dltA* mutant strain seemed to be devoid of D-alanine whereas that of the
42
43 parental and the *lgt* mutant strains retained it (right panel in Fig. 2c). When macrophages
44
45 were incubated with these WTA preparations, the phosphorylation of JNK was not induced
46
47 irrespective of the presence of bound D-alanine in WTA (Fig. 2d), indicating that WTA does
48
49 not serve as a ligand for TLR2. We next tested if WTA influences the action of the TLR2
50
51 ligand. To this end, macrophages were incubated with Pam3Cys, a synthetic TLR2 ligand,
52
53 in the presence and absence of WTA. However, the level of phosphorylated JNK was not
54
55 altered by the addition of WTA (Fig. 2d). These results suggested that D-alanylated WTA
56
57
58
59
60

1
2
3 serves as an auxiliary factor for the TLR2 ligand in the context of bacterial cell wall so that it
4
5 effectively acts on TLR2 to induce the phosphorylation of JNK.
6
7
8
9

10 **Enhanced superoxide production and accelerated killing of engulfed bacteria in** 11 **macrophages incubated with *S. aureus* deficient in *dltA* or *tagO***

12
13 We next determined the level of superoxide production in *S. aureus*-incubated macrophages,
14
15 which we previously showed to be inhibited by phosphorylated and thus activated JNK.¹⁰
16
17

18 The level of superoxide released from macrophages into culture media was significantly
19
20 higher on incubation with a mutant strain lacking the expression of *dltA*, *tagO* or *lgt* than with
21
22 the parental strain (left panel in Fig. 3a). The effect of the mutation in *dltA* was mitigated
23
24 when the *dltABCD* operon was introduced into the mutant strain (right panel in Fig. 3a),
25
26 confirming the requirement of *dltA* for the effective inhibition of superoxide production in
27
28 macrophages by *S. aureus*. The viability of engulfed *S. aureus* was then assessed based on
29
30 their colony-forming ability. The value of colony-forming units obtained with the *dltA*
31
32 mutant was much smaller than that with the parental or the same mutant strain that had
33
34 acquired the corresponding wild-type operon (Fig. 3b), indicating that *S. aureus* lacking the
35
36 expression of *dltA* was more efficiently killed in macrophages. Furthermore, the *dltA* mutant
37
38 survived the killing in macrophages when the cultures were supplemented with
39
40 *N*-acetyl-L-cysteine, a superoxide scavenger (Fig. 3c). We next examined if the recognition
41
42 and engulfment of *S. aureus* alters the activity of macrophages other than superoxide
43
44 production. For this purpose, macrophages were incubated with various *S. aureus* stains,
45
46 and their whole-cell lysates were assayed for α -*N*-acetylglucosaminidase, a major lysosomal
47
48 enzyme. However, its activity did not change after the incubation with any of the bacterial
49
50 strains tested (Fig. 3d), suggesting that the lysosomal activity is not influenced by *S. aureus*.
51
52

53
54
55
56
57
58
59
60 These results indicated that a lack of the expression of *dltA* or *tagO* in *S. aureus* causes
augmented production of superoxide and accelerated killing of engulfed bacteria in

1
2
3 macrophages, and thus suggested a role for the D-alanylation of WTA in the survival of *S.*
4
5
6 *aureus* in macrophages.
7
8
9

10 **Inefficient activation of NF- κ B in TLR2-expressing cells incubated with *S. aureus*** 11 12 **lacking *dltA* or *tagO*** 13 14

15 We next determined the level of NF- κ B-dependent gene expression in TLR2-expressing
16
17 HEK293 cells, examining a role for *dltA* and *tagO* in the activation of TLR2. The
18
19 expression of a NF- κ B-induced gene coding for luciferase depended on the presence of TLR2
20
21 in HEK293 cells as well as the addition of *S. aureus* to them (Fig. 4a), indicating that the level
22
23 of active NF- κ B reflects the activation of TLR2-initiated signaling by bacteria. HEK293
24
25 cells incubated with the *dltA* mutant produced much less luciferase than those treated with the
26
27 parental strain, and this decrease was recovered when the *dltABCD* operon was introduced
28
29 into the mutant (Fig. 4b). A decrease in the level of active NF- κ B was similarly observed
30
31 when the mutants for *tagO*, *SA0614*, and *SA0615*, which all gave a reduced level of
32
33 phosphorylated JNK in macrophages (see Fig. 1b), were tested (Fig. 4c). In contrast, the
34
35 other mutant strains with no effect on the phosphorylation of JNK activated NF- κ B as
36
37 effectively as the parental strain (Fig. 4c). [These results suggested that D-alanylated WTA is](#)
38
39 [required for *S. aureus* to effectively induce the TLR2-mediated activation of NF- \$\kappa\$ B.](#)
40
41
42
43
44
45
46
47
48

49 Taken together, the effects of *dltA* and *tagO* on JNK phosphorylation, superoxide production,
50
51 the survival of engulfed bacteria, and the activation of TLR2-mediated signaling are
52
53 consistent with the concept that a component of *S. aureus*, i.e. D-alanylated WTA, additional
54
55 to an authentic ligand is required for TLR2 to effectively phosphorylate and activate JNK
56
57 leading to the inhibition of superoxide production in *S. aureus*-engulfing macrophages.
58
59
60

Discussion

The study presented here showed that genes responsible for the synthesis and D-alanylation of teichoic acids are required for the TLR2/JNK-dependent survival of *S. aureus* in macrophages. The importance of D-alanylated LTA of *S. aureus* for the production of a pro-inflammatory cytokine by macrophages has been reported.³¹ However, our results clearly indicated that WTA, not LTA, is necessary for the TLR2-mediated phosphorylation of JNK. There are preceding reports showing an *in vivo* role for the D-alanylation of teichoic acids of *S. aureus*³² and *Streptococcus gordonii*³³ in bacterial virulence and TLR2-mediated host defense. Our study provides a reasonable explanation for the observation in these papers that bacteria evoking a higher level of immune response in host organisms are, at the same time, more infectious and virulent. We also showed that the D-alanylation of teichoic acids is necessary for *S. aureus* to effectively activate NF- κ B in TLR2-expressing cells. It can thus be anticipated that D-alanylated WTA plays an important role in the TLR2-initiated signaling pathways in immune cells in favor of both host organisms and invading microbial pathogens. Purified WTA by itself did not induce JNK phosphorylation in macrophages, and exogenously added WTA was not effective on the phosphorylation of JNK induced by a synthetic ligand for TLR2. Therefore, it is probable that D-alanylated WTA does not directly act on TLR2 as a ligand but facilitates the activation of TLR2 by an authentic ligand like lipoproteins or lipopeptides in the context of the bacterial cell wall. There is a report showing that WTA mediates the interaction of *S. aureus* with airway epithelial cells.³⁴ However, this is not the case in our study because the level of the phagocytosis of *S. aureus* by macrophages does not differ between the parental and *dltA* mutant strains. We speculate that WTA modulates the cell wall milieu for lipoproteins so that they effectively serve as a ligand for TLR2.

1
2
3 The stimulation of JNK phosphorylation occurred when TLR2-lacking
4 macrophages were incubated with LPS. This suggests that the JNK-mediated inhibition of
5 killing of engulfed bacteria is not restricted to TLR2-stimulating bacteria, *S. aureus* in our
6 study, but is observed for the bacteria recognized by TLR4. We previously reported that
7 TLR4 delays the fusion between lysosomes and phagosomes that contain engulfed apoptotic
8 cells.²⁵ Other investigators have also reported the involvement of TLR in the regulation of
9 phagosome maturation and thus the fate of engulfed materials including microbial pathogens,
10 microbe-infected cells, and apoptotic cells.³⁵⁻³⁷ They argue that TLR-mediated control of
11 phagosome maturation relates to the regulation of antigen presentation.³⁸⁻⁴¹ It is thus
12 obvious that TLRs, TLR2 and TLR4 in particular, participate in not only humoral but also
13 cellular immune responses. Further investigation will be necessary to obtain a complete
14 picture that illustrates the mechanisms and consequences of TLR-mediated regulation of
15 cellular immunity including phagocytosis.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

34 35 36 **Acknowledgements** 37

38
39
40
41 We thank Douglas Golenbock, Yoshiyuki Adachi, and Shizuo Akira for the materials used in
42 this study. We are also grateful to Masahito Hashimoto for his discussion and suggestions
43 on the analysis of cell wall components. This work was supported by the Grant-in-Aid for
44 Scientific Research from Japan Society for the Promotion of Science (Nos. 16570112,
45 18570123 and 20570127) and from Ministry of Education, Culture, Sports, Science and
46 Technology-Japan (No. 18057009) to A.S., by the Industrial Technology Research Grant
47 Program of the New Energy and Industrial Technology Development Organization of Japan
48 (No. 04A01528) to K.K., and in part by the Bilateral Programme of Joint Research Project
49 from Japan Society for the Promotion of Science to Y.N. and the Joint Research Project under
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

the KOSEF-JSPS Cooperative Programme (F01-2006-000-10016-0) of MOST/KOSEF to
B.L.L.

For Peer Review

References

1. Rosenberver CM, Finlay BB. Phagocyte sabotage: disruption of macrophage signaling by bacterial pathogens. *Nat Rev Mol Cell Biol* 2003; **4**:385–396.
2. Coombes BK, Valdez Y, Finlay BB. Evasive maneuvers by secreted bacterial proteins to avoid innate immune responses. *Curr Biol* 2004; **14**:R856–R867.
3. Bhavsar AP, Guttman JA, Finlay BB. Manipulation of host-cell pathways by bacterial pathogens. *Nature* 2007; **449**:827–834.
4. Sansonetti PJ, Di Santo JP. Debugging how bacteria manipulate the immune response. *Immunity* 2007; **26**:149–161.
5. Nizet V. Understanding how leading bacterial pathogens subvert innate immunity to reveal novel therapeutic targets. *J Allergy Clin Immunol* 2007; **120**:13–22.
6. Ernst JD. Bacterial inhibition of phagocytosis. *Cell Microbiol* 2000; **2**:379–386.
7. Dunn Jr. WA, Dorn BR, Progulsk-Fox A. Trafficking of bacterial pathogens to autophagosomes. In: Klionsky DJ, ed. *Autophagy*. Austin: Landes Biosciences, 2003: chapter 19.
8. Levine B, Deretic V. Unveiling the roles of autophagy in innate and adaptive immunity. *Nat Rev Immunol* 2007; **7**:767–777.
9. Levine B, Kroemer G. Autophagy in the pathogenesis of diseases. *Cell* 2008; **132**:27–42.
10. Watanabe I, Ichiki M, Shiratsuchi A, Nakanishi Y. TLR2-mediated survival of *Staphylococcus aureus* in macrophages: a novel bacterial strategy against host innate immunity. *J Immunol* 2007; **178**:4917–4925.
11. Hashimoto M, Tawaratsumida K, Kariya H, Kiyohara A, Suda Y, Kriksae F, Kirikae T, Götz F. Not lipoteichoic acid but lipoproteins appear to be the dominant immunobiologically active compounds in *Staphylococcus aureus*. *J Immunol* 2006; **177**:3162–3169.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
12. Jin MS, Kim SE, Heo JY, Lee ME, Kim HM, Paik S-G, Lee H, Lee J-O. Crystal structure of the TLR1-TLR2 heterodimer induced by binding of a tri-acylated lipopeptide. *Cell* 2007; **130**:1071–1082.
13. Zähringer U, Lindner B, Inamura S, Heine H, Alexander C. TLR2 — promiscuous or specific? A critical re-evaluation of a receptor expressing apparent broad specificity. *Immunobiology* 2008; **213**:205–204.
14. Kurokawa K, Lee H, Roh K-B, Asanuma M, Kim YS, Nakayama H, Shiratsuchi A, Choi Y, Takeuchi O, Kang HJ, Dohmae N, Nakanishi Y, Akira S, Sekimizu K, Lee BL. The triacylated ATP binding cluster transporter substrate-binding lipoprotein of *Staphylococcus aureus* functions as a native ligand for the Toll-like receptor 2. *J Biol Chem* 2009; **284**:8406–8411.
15. Schmalzer M, Jann NJ, Ferracin F, Landolt LZ, Biswas L, Götz F, Landmann R. Lipoproteins in *Staphylococcus aureus* mediate inflammation by TLR2 and iron-dependent growth in vivo. *J Immunol* 2009; **182**:7110–7118.
16. Tawaratsumida K, Furuyashiki M, Katsumoto M, Fujimoto Y, Fukase K, Suda Y, Hashimoto M. Characterization of N-terminal structure of TLR2-activating lipoprotein in *Staphylococcus aureus*. *J Biol Chem* 2009; **284**:9147–9152.
17. Turner JD, Langley RS, Johnston KL, Gentil K, Ford L, Wu B, Graham M, Sharpley F, Slatko B, Pearlman E, Taylor MJ. Wolbachia lipoprotein stimulates innate and adaptive immunity through toll-like receptors 2 and 6 (TLR2/6) to induce disease manifestations of filariasis. *J Biol Chem*, May 2009; doi:10.1074/jbc.M901528200
18. von Aulock S, Hartung T, Hermann C. Comment on "Not lipoteichoic acid but lipoproteins appear to be the dominant immunobiologically active compounds in *Staphylococcus aureus*". *J Immunol* 2007; **178**:2610.
19. Novick RP, Ross HF, Projan SJ, Kornblum J, Kreiswirth B, Moghazeh S. Synthesis of staphylococcal virulence factors is controlled by a regulatory RNA molecule. *EMBO J* 1993;

1
2
3 12:3967-3975.
4

5
6 20. Tao L, LeBlanc DJ, Ferretti JJ. Novel streptococcal-integration shuttle vectors for gene
7
8 cloning and inactivation. *Gene* 1992; **120**:105–110.
9

10
11 21. Shiratsuchi A, Nakanishi Y. Phosphatidylserine-mediated phagocytosis of anticancer
12
13 drug-treated cells by macrophages. *J Biochem* 1999; **26**:1101–1106.
14

15
16 22. Takeuchi O, Hoshino K, Kawai T, Sanjo H, Takada H, Ogawa T, Takeda K, Akira S.
17
18 Differential roles of TLR2 and TLR4 in recognition of Gram-negative and Gram-positive
19
20 bacteria cell wall components. *Immunity* 1999; **11**:443–451.
21

22
23 23. Chen PS Jr, Toribara TY, Warner H. Microdetermination of phosphorus. *Anal Chem*
24
25 1956; **28**:1756–1758.
26

27
28 24. Wolters PJ, Hildebrandt KM, Dickie JP, Anderson JS. Polymer length of teichuronic acid
29
30 released from cell walls of *Micrococcus luteus*. *J Bacteriol* 1990; **172**:5154–5159.
31

32
33 25. Shiratsuchi A, Watanabe I, Takeuchi O, Akira S, Nakanishi Y. Inhibitory effect of
34
35 Toll-like receptor 4 on fusion between phagosomes and endosomes/lysosomes in
36
37 macrophages. *J Immunol* 2004; **172**:2039–2047.
38

39
40 26. Means TK, Wang S, Lien E, Yoshimura A, Golenbock DT, Fenton MJ. Toll-like receptors
41
42 mediate cellular activation by *Mycobacterium tuberculosis*. *J Immunol* 1999; **163**:3920–3927.
43

44
45 27. Ikeda Y, Adachi Y, Ishibashi K, Miura N, Ohno N. Activation of Toll-like
46
47 receptor-mediated NF- κ B by zymosan-derived water-soluble fraction: possible contribution
48
49 of endotoxin-like substances. *Immunopharmacol Immunotoxicol* 2005; **27**:285–298.
50

51
52 28. Neuhaus FC, Baddiley J. A continuum of anionic charge: structures and functions of
53
54 D-alanyl-teichoic acids in gram-positive bacteria. *Microbiol Mol Biol Rev* 2003; **67**:686-723.
55

56
57 29. Kurokawa K, Kaito C, Sekimizu K. Two-component signaling in the virulence of
58
59 *Staphylococcus aureus*: a silkworm larvae-pathogenic agent infection model of virulence.
60
Methods Enzymol 2007; **422**:233–244.

- 1
2
3 30. Herbert S, Bera A, Nerz C, Kraus D, Peschel A, Goerke C, Meehl M, Cheung A, Götz F.
4
5 Molecular basis of resistance to muramidase and cationic antimicrobial peptide activity of
6
7 lysozyme in staphylococci. *PLoS Pathog* 2007; **3**:e102.
8
9
10 31. Morath S, Stadelmaier A, Geyer A, Schmidt RR, Hartung T. Synthetic lipoteichoic acid
11
12 from *Staphylococcus aureus* is a potent stimulus of cytokine release. *J Exp Med* 2002;
13
14 **195**:1635–1640.
15
16
17 32. Kristian SA, Lauth X, Nizet V, Goetz F, Neumeister B, Peschel A, Landmann R.
18
19 Alanylation of teichoic acids protects *Staphylococcus aureus* against Toll-like receptor
20
21 2-dependent host defense in a mouse tissue cage infection model. *J Infect Dis* 2003;
22
23 **188**:414–423.
24
25
26 33. Chan KG, Mayer M, Davis EM, Halperin SA, Lin T-J, Lee SF. Role of D-alanylation of
27
28 *Streptococcus gordonii* lipoteichoic acid in innate and adaptive immunity. *Infect Immun* 2007;
29
30 **75**:3033–3042.
31
32
33 34. Weidenmaier C, Kokai-Kun JF, Kristian SA, Chanturiya T, Kalbacher H, Gross M,
34
35 Nicholson G, Neumeister B, Mond JJ, Peschel A. Role of teichoic acids in *Staphylococcus*
36
37 *aureus* nasal colonization, a major risk factor in nosocomial infections. *Nat Med* 2004;
38
39 **10**:243–245.
40
41
42 35. Blander JM, Medzhitov R. Regulation of phagosome maturation by signals from Toll-like
43
44 receptors. *Science* 2004; **304**:1014–1018.
45
46
47 36. Blander JM, Medzhitov R. Toll-dependent selection of microbial antigens for presentation
48
49 by dendritic cells. *Nature* 2006; **440**:808–812.
50
51
52 37. Apetoh L, Ghiringhelli F, Tesniere A, Obeid M, Ortiz C, Criollo A, Mignot G, Maiuri MC,
53
54 Ullrich E, Saulnier P, Yang H, Amigorena S, Ryffel B, Barrat FJ, Saftig P, Levi F, Lidereau R,
55
56 Nogues C, Mira J-P, Chompret A, Joulin V, Clavel-Chapelon F, Bourhis J, André F, Delaloge
57
58 S, Tursz T, Kroemer G, Zitvogel L. Toll-like receptor 4-dependent contribution of the
59
60 immune system to anticancer chemotherapy and radiotherapy. *Nat Med* 2007; **13**:1050–1059.

- 1
2
3 38. Blander JM. Coupling Toll-like receptor signaling with phagocytosis: potentiation of
4 antigen presentation. *Trends Immunol* 2007; **28**:19–25.
5
6
7
8 39. Apetoh L, Ghiringhelli F, Tesniere A, Criollo A, Ortiz C, Lidereau R, Mariette C, Chaput
9 N, Mira J-P, Delaloge S, André F, Tursz T, Kroemer G, Zitvogel L. The interaction between
10 HMGB1 and TLR4 dictates the outcome of anticancer chemotherapy and radiotherapy.
11 *Immunol Rev* 2007; **220**:47–59.
12
13
14
15
16
17 40. Blander JM. Signalling and phagocytosis in the orchestration of host defense. *Cell*
18 *Microbiol* 2007; **9**:290–299.
19
20
21
22 41. Apetoh L, Tesniere A, Ghiringhelli F, Kroemer G, Zitvogel L. Molecular interactions
23 between dying tumor cells and the innate immune system determine the efficacy of
24 conventional anticancer therapies. *Cancer Res* 2008; **68**:4026–4030.
25
26
27
28
29 42. Kaito C, Sekimizu K. Colony spreading in *Staphylococcus aureus*. *J Bacteriol* 2007;
30 **189**:2553–2557.
31
32
33
34 43. Oku Y, Kurokawa K, Matsuo M, Yamada S, Lee B-L, Sekimizu, K. Pleiotropic roles of
35 polyglycerolphosphate synthase of lipoteichoic acid in growth of *Staphylococcus aureus* cells.
36 *J Bacteriol* 2009; **191**:141–151.
37
38
39
40
41 44. Takahashi J, Komatsuzawa H, Yamada S, Nishida T, Labischinski H, Fujiwara T, Ohara
42 M, Yamagishi J, Sugai M. Molecular characterization of an *atl* null mutant of *Staphylococcus*
43 *aureus*. *Microbiol Immunol* 2002; **46**:601–612.
44
45
46
47
48 45. Ichihashi N, Kurokawa K, Matsuo M, Kaito C, Sekimizu K. Inhibitory effects of basic or
49 neutral phospholipid on acidic phospholipid-mediated dissociation of adenine nucleotide
50 bound to DnaA protein, the initiator of chromosomal DNA replication. *J Biol Chem* 2003;
51 **278**:28778–28786.
52
53
54
55
56
57
58
59
60

Figure legends

Figure 1. Inefficient phosphorylation of JNK in macrophages incubated with *S. aureus* lacking the expression of D-alanylated WTA. Mouse peritoneal macrophages were incubated with various *S. aureus* strains: the parental strain RN4220 (parent) and mutant strains lacking the expression of the indicated genes with the symbol 'Δ', and responses in macrophages were examined. (a) Macrophages were incubated with the parental strain or the mutant strain deficient in *dltA* for the indicated periods (30 min for phagocytosis), and the levels of phosphorylated JNK and phagocytosis of bacteria were determined. Arrowheads with the terms 'P-JNK' and 'total' denote the positions of the phosphorylated and both the phosphorylated and unphosphorylated forms of JNK, respectively. (b, c) Macrophages were incubated with the indicated *S. aureus* strains for 30 min, and the level of phosphorylated JNK was determined. In (c), the *S. aureus* strains used were the parental strain transformed with pHY300PLK alone (vector), Δ*dltA* transformed with pHY300PLK alone, Δ*dltA* transformed with *dltABCD* operon-expressing pHY300PLK (*dltABCD*), Δ*tagO* transformed with pHY300PLK alone, and Δ*tagO* transformed with *tagO*-expressing pHY300PLK (*tagO*).

Figure 2. Absence of JNK phosphorylation in macrophages treated with D-alanylated WTA. (a) Peritoneal macrophages of *tlr2*-disrupted mice were incubated with the indicated *S. aureus* strains or LPS (0.1 mg/ml) for 30 min, and the level of phosphorylated JNK was determined. (b) Macrophages of wild-type mice were incubated with a fraction rich in lipoproteins (0.1 μg/ml) obtained from the indicated *S. aureus* strains for 30 min, and the level of phosphorylated JNK was determined. (c) WTA prepared from the indicated *S. aureus* strains was characterized: determination of the phosphorus content (left), PAGE of WTA visualized with silver (middle), and thin-layer chromatography of D-alanine alkali liberated

1
2
3 from WTA (visualized by the ninhydrin reaction) (right). The arrowhead with 'D-Ala' in the
4
5 right panel points to the position of D-alanine analyzed as a control. (d) Wild-type
6
7 macrophages were incubated for 15 min with and without Pam3Cys (1 $\mu\text{g}/\text{ml}$), a synthetic
8
9 TLR2 ligand, and WTA (10 $\mu\text{g}/\text{ml}$) purified from the indicated *S. aureus* strains, and the level
10
11 of JNK phosphorylation was determined.
12
13
14
15
16
17

Figure 3. Increased production of superoxide and accelerated killing of engulfed bacteria in
18
19 macrophages incubated with *dltA*-, *tagO*-, or *lgt*-deficient *S. aureus*. Mouse peritoneal
20
21 macrophages were incubated with *S. aureus*, and the levels of superoxide production and
22
23 values of colony-forming units of engulfed bacteria were determined. (a) Macrophages were
24
25 incubated with the indicated *S. aureus* strains for 30 min, and the amount of superoxide in the
26
27 culture medium was determined and presented relative (in percentage terms) to that in
28
29 incubation with the parental strain RN4220 (left panel) or the *dltA* mutant transfected with the
30
31 empty vector (right panel). (b) The value of colony-forming units of the indicated *S. aureus*
32
33 strains that had been engulfed by macrophages was determined 0 and 30 min after elimination
34
35 of unengulfed bacteria and shown relative (in percentage terms) to that obtained at time 0.
36
37 (c) Macrophages that had been pre-treated with *N*-acetyl-L-cysteine (LNAC) (2 mM) or left
38
39 untreated at 37°C for 1 h were incubated with the indicated strains, and the value of
40
41 colony-forming units of engulfed bacteria was determined. (d) Macrophages were incubated
42
43 with the indicated *S. aureus* strains or left untreated (none), and their whole-cell lysates were
44
45 assayed for the activity of α -*N*-acetylglucosaminidase at the indicated time points.
46
47
48
49
50
51
52
53
54
55


Figure 4. Inefficient activation of NF- κ B in TLR2-expressing cells incubated with
56
57 *dltA*-deficient *S. aureus*. (a) (left) HEK293 cells with and without TLR2 expression were
58
59 incubated with RN4220 (HEK293 : bacteria = 1 : 10³) or left untreated, and the level of
60

1
2
3 NF- κ B activation was determined and shown relative to that in cells expressing TLR2 with no
4 added bacteria, taken as 1. (right) HEK293 cells (1×10^5) expressing TLR2 were incubated
5
6 with increasing amounts of RN4220, and the level of NF- κ B activation was determined and
7
8 shown relative to that in the experiment with no added bacteria, taken as 1. (b, c) HEK293
9
10 cells expressing TLR2 were incubated with the indicated *S. aureus* strains (HEK293 : bacteria
11
12 = 1 : 100), and the level of NF- κ B activation was determined and shown relative (in
13
14 percentage terms) to that in the experiment with the parental strain RN4220.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. *S. aureus* strains and plasmids used in this study.


Names	Genotypes and characteristics	References
RN4220	NCTC8325-4, restriction mutant	19
M0793	RN4220 <i>dltA</i> ::pT0793	42
M0702	RN4220 <i>tagO</i> ::pT0702	42
M0614	RN4220 <i>SA0614</i> ::pSFSA0614	this study
M0615	RN4220 <i>SA0615</i> ::pSFSA0615	this study
M0674/pM101	RN4220 <i>ltaS</i> :: <i>phleo</i> /pM101	43
M0875	RN4220 <i>ypfP</i> ::pT0875	42
JT1304	RN4220 <i>atl</i> :: <i>cat</i>	44
CK1001	RN4220 <i>mprF</i> ::pCK20 <i>mprF</i>	45
M0107	RN4220 <i>spa</i> :: <i>phleo</i>	unpublished observation
T013	RN4220 <i>lgt</i> ::pMlgt	14
pHY300PLK	<i>E. coli</i> - <i>S. aureus</i> shuttle vector; <i>Amp^r Tet^r</i>	purchased from Takara-Bio
p0793	pHY300 with intact <i>dltABCD</i> from RN4220	42
p0702	pHY300 with intact <i>tagO</i> from RN4220	42
pSF151	<i>S. aureus</i> integration vector; <i>Km^r</i>	20
pSFSA0614	pSF151 with partial <i>SA0614</i> from RN4220	this study
pSFSA0615	pSF151 with partial <i>SA0615</i> from RN4220	this study

Shiratsuchi et al. Fig. 1



view


Shiratsuchi et al. Fig. 2



new


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Shiratsuchi et al. Fig. 3



Peer Review

Shiratsuchi et al. Fig. 4



Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60