

# Swelling-activated potassium channel in porcine pigmented ciliary epithelial cells.

メタデータ	言語: eng 出版者: 公開日: 2017-10-05 キーワード (Ja): キーワード (En): 作成者: メールアドレス: 所属:
URL	<a href="http://hdl.handle.net/2297/29471">http://hdl.handle.net/2297/29471</a>

# Swelling-Activated Potassium Channel in Porcine Pigmented Ciliary Epithelial Cells

Masayuki Takahira, Mayumi Sakurai, Norimasa Sakurada, and Kazuhisa Sugiyama

Department of Ophthalmology & Visual Science, Kanazawa University, Kanazawa  
920-8640, Japan

Word count: 3123

Supported by Grants-in-Aid 16591742(MT) and 21791674(NS) from Ministry of  
Education, Culture, Sports, Science and Technology, Japan

Corresponding author: Masayuki Takahira

Takara-Machi 13-1, Department of Visual Neuroscience & Ophthalmology, Kanazawa  
University, Kanazawa 920-8640, Japan

Phone: 76-265-2403 Fax: 76-222-9660 Country Code = 81

E-mail: [takahira@kenroku.kanazawa-u.ac.jp](mailto:takahira@kenroku.kanazawa-u.ac.jp)

**ABSTRACT**

**Purpose:** Ion channels in the ciliary epithelium play critical roles in the formation of aqueous humor in the eye. The present study identified a novel, swelling-activated  $K^+$  current in freshly dissociated porcine pigmented ciliary epithelial cells.

**Methods:** Ciliary epithelial cells were freshly dissociated from porcine eyes. Whole-cell currents were recorded using the patch-clamp technique in pigmented and non-pigmented ciliary epithelial cell (PCE-NPCE) pairs or single PCE cells.

**Results:** The zero-current potential was  $-49 \pm 13$  mV in PCE-NPCE cell pairs ( $n = 97$ ), and  $-52 \pm 12$  mV in single PCE cells ( $n = 30$ ). Whole-cell currents in these cells were dominated by an outwardly rectifying  $K^+$  current activated by potentials more positive than  $-90$  mV, which never inactivated during prolonged depolarization. The  $K^+$  current was significantly augmented by hypotonic cell perfusion. External  $Ba^{2+}$  was a blocker of this  $K^+$  conductance ( $IC_{50}$  of  $0.38$  mM), but the conductance was insensitive to external  $TEA^+$ . Linopirdine, a specific inhibitor of KCNQ channels, effectively blocked the  $K^+$  current in these PCE cells. **Conclusions:** Porcine PCE cells express a swelling-activated  $K^+$  channel, which may be a member of the KCNQ/Kv7 channel family. This  $K^+$  channel is active near resting potentials and could contribute to the regulation of cell volume and water transport via the ciliary epithelia.

## INTRODUCTION

The ciliary body epithelium is a transporting epithelium, composed of two epithelial layers, a non-pigmented ciliary epithelial (NPCE) cell layer facing the vitreal space, and a pigmented ciliary epithelial (PCE) cell layer facing the ciliary stroma. By transporting ions and water between the ciliary body and vitreal space, the ciliary epithelium plays critical roles in the formation of aqueous humor in the eye.

Studies on ion transport via the ciliary epithelium have mainly focused on components underlying chloride secretion, which limit aqueous humor production.<sup>1,2</sup> In particular,  $\text{Cl}^-$  channels in NPCE cells mediate the final step of  $\text{Cl}^-$  release and thereby regulate the rate of aqueous humor secretion. The molecular identity of NPCE  $\text{Cl}^-$  channels has not been precisely elucidated, but several lines of evidence indicate that the  $\text{ClC-3}$  member of the  $\text{ClC}$  family of channels and transporters likely plays a major role in  $\text{Cl}^-$  inflow.<sup>3,4</sup> In addition to anion channels, epithelial potassium channels also contribute to critical cellular functions including the secretion of ions and water across the epithelia. The rate of anion secretion depends on the activity of  $\text{K}^+$  channels since cell hyperpolarization caused by the opening of  $\text{K}^+$  channels increases the driving force for anion efflux.<sup>5</sup> Several previous studies have described some  $\text{K}^+$  channels expressed in the ciliary epithelium, including voltage-gated delayed rectifiers,<sup>6-8</sup>  $\text{Ca}^{2+}$ -activated outward  $\text{K}^+$  channels<sup>7,9,10</sup> and an inward rectifier activated by hyperpolarization.<sup>6,11</sup> It is unknown whether these  $\text{K}^+$  currents in the ciliary epithelium are sensitive to cell swelling.

In the retinal pigment epithelium (RPE), another ocular secretory epithelium,  $\text{K}^+$  conductance is mainly composed of two types of  $\text{K}^+$  channels, an inward rectifier  $\text{Kir7.1}$ <sup>12,13</sup> and a M-type  $\text{K}^+$  channel.<sup>14,15</sup> The latter was termed from its resemblance to

the M-current, a  $K^+$  current distributed in neurons and other excitable cells. This M-type  $K^+$  current in RPE is a voltage-gated  $K^+$  channel with an activation threshold near the equilibrium potential for  $K^+$ , and thus, contributes to the resting potential of the cell. One of the characteristic properties of the M-type  $K^+$  current in the RPE is its insensitivity to tetraethylammonium (TEA), a classical  $K^+$  channel blocker.<sup>14,15</sup>

In this study, we identified a swelling-activated, non-inactivating outward  $K^+$  current in freshly dissociated ciliary pigmented epithelial cells. The  $K^+$  current was essentially insensitive to TEA and resembled the M-type  $K^+$  current of RPE in terms of voltage-dependence, kinetics and blocker sensitivities.

## **METHODS**

### *Solutions*

External and internal solutions were buffered with HEPES as in previous patch-clamp studies of corneal epithelium,<sup>16</sup> and retinal pigment epithelium.<sup>15</sup> Briefly, the standard Ringer's solution consisted of (in mM): 135 NaCl, 5.0 KCl, 10 HEPES, 10 glucose, 1.8  $CaCl_2$  and 1.0  $MgCl_2$ , and was titrated to pH 7.4 with NaOH. The osmolarity of standard Ringer's solution was  $302 \pm 6$  mosmol/l. The standard pipette solution consisted of (in mM): 30 KCl, 83 potassium gluconate, 5.0 HEPES, 5.5 EGTA-KOH, 0.5  $CaCl_2$  and 2.0  $MgCl_2$ , and was titrated to pH 7.2 with KOH. The osmolarity of standard pipette solution was  $257 \pm 5$  mosmol/l. An amphotericin B-stock solution (1.2 mg amphotericin B / 50  $\mu$ l DMSO) was added to 2.0 ml of the pipette solution to give a final concentration of 240  $\mu$ g/ml. The cell isolation medium was similar to that used in a previous study on RPE and contained (in mM): 135 N-methyl-D-glucamine (NMDG)-Cl, 5.0 KCl, 10 HEPES, 3.0 EDTA-KOH, 10 glucose, 3.0 cysteine, 1.0

glutathione, 1.0 L(+)-ascorbic acid, 1.0 taurine and 0.2 mg/ml papain (Type III), and was titrated to pH 7.4 with NaOH. In experiments where the concentrations of  $K^+$ , or tetraethylammonium<sup>+</sup> ( $TEA^+$ ) were varied, NaCl was replaced by an equimolar amount of the appropriate  $Cl^-$  salt. A hypotonic solution was made by diluting standard Ringer's solution to 20% and adjusting  $[K^+]$  to 5 mM to give an osmolarity of  $239 \pm 2$  mosmol/l. Linopirdine was dissolved in DMSO and then diluted in Ringer's solution. Throughout the experiments, the final concentration of DMSO in the perfusates was no more than 0.2%, which did not affect whole cell currents in this study.

NaCl, KCl,  $CaCl_2$  and L(+)-ascorbic acid were obtained from Wako Chemical Co. (Osaka, Japan). All the other reagents were obtained from Sigma Chemical Co. (St. Louis, MO, USA).

#### *Cell Isolation*

All experiments were conducted in accordance with the ARVO Statement for the Use of Animals in Ophthalmic and Vision Research. Porcine eyes were enucleated just after death at a local slaughterhouse and transferred to the laboratory within 30 minutes. A piece of ciliary body with epithelium was cut out and incubated in cell isolation medium for 30 to 40 minutes. The tissue was transferred to standard Ringer's solution containing 0.1 % bovine serum albumin for 3 minutes, and then incubated in standard Ringer's solution for 10 minutes followed by gentle vortexing. Isolated cells were observed in a sample of the suspension by phase-contrast microscopy. The cell suspension was stored at 4 °C for up to 24 hours before use.

#### *Cell Perfusion*

Isolated ciliary epithelial cells were transferred to a Lucite perfusion chamber

(RC-5/25, Warner Instrument, Hamden, CT) and settled for 10 to 20 minutes before perfusion. The perfusate flowed into the chamber under gravity at a flow rate of 0.7 ml / min by a perfusion system (BPS-4, ALA Scientific Instruments, Westbury, NY) and was continuously removed by suction. Fluid height was adjusted to give a chamber volume of approximately 0.5 ml and a complete solution exchange within 2 minutes. All experiments were conducted at room temperature (20 to 25 °C).

### *Electrophysiological Methods*

Patch pipettes were pulled from borosilicate glass tubing (BF150-110-10, Sutter Instruments, San Rafael, CA) with a multistage programmable puller (P-97, Sutter Instruments, San Rafael, CA). The pipette input resistance was between 1 and 3 M $\Omega$ . Under phase-contrast microscopy (ECLIPSE TE300, Nikon, Tokyo, Japan), a target cell was selected, and a pipette tip was pressed onto the cell membrane using a micromanipulator (MP-285, Sutter Instruments, San Rafael, CA) to establish a gigaohm seal. Currents under voltage-clamp were recorded by a Heka amplifier system (EPC-8, Heka, Lambrecht/Pfalz, Germany). The built-in low pass filter was set to 3 kHz unless noted otherwise. Recordings were referenced to an Ag-AgCl electrode (EP-2, WPI, Sarasota, FL). The membrane capacitance was compensated by built-in circuits. The apparent membrane potential was corrected for the pipette tip potential (10 mV). Statistical data are presented as mean  $\pm$  S.D. Data were fitted by a non-linear, least-squares fitting using IGOR Pro (Wavemetrics, Lake Oswego, OR).

## **RESULTS**

### *Whole-Cell Currents*

Pairs of pigmented and non-pigmented ciliary epithelial cells (PCE and NPCE cells) with broad contact area (Figure 1) were selected for the recording. The PCE-NPCE cell pairs are typically similar to that observed in a previous report.<sup>17</sup> In the initial and major phase of this study, we found M-type  $K^+$  current<sup>14,15</sup> expressed in the PCE-NPCE cell pairs, as shown in Figure 2. Thereafter, we tried to record from isolated single ciliary epithelial cells, and found the identical M-type  $K^+$  current was observed in single PE cells ( $n = 30$ ) but never in single NPE cells ( $n = 11$ ). Therefore, data obtained from PCE-NPCE cell pairs ( $n = 97$ ) and single PCE cells ( $n = 30$ ) were pooled in this study. The zero-current potential ( $V_0$ ) was measured in the zero-current clamp mode after establishment of a perforated patch configuration: the mean  $V_0$  was  $-49 \pm 13$  mV in PCE-NPCE cell pairs ( $n = 97$ ) and  $-52 \pm 12$  mV in single PCE cells ( $n = 30$ ). The membrane capacitance and input resistance were  $47 \pm 34$  pF and  $489 \pm 324$  Mohm in PCE-NPCE cell pairs ( $n = 93$ ), which were somewhat larger than those values of  $37 \pm 29$  pF and  $389 \pm 277$  Mohm measured in single PCE cells ( $n = 29$ ). Student's *t*-tests showed no significant difference in these values between the two groups.

#### *Effect of Holding Potential.*

Representative families of whole-cell currents in a PCE-NPCE cell pair are depicted in Figure 2. The currents in Figure 2A were generated by a series of voltage pulses from a holding potential of  $-10$  mV. Hyperpolarizing pulses produced time-dependent relaxations in current that reversed around  $-80$  mV. The current relaxations reflect the deactivation of a conductance. The following tail currents elicited by a depolarizing voltage step to the holding potential ( $-10$  mV) from various hyperpolarized prepulses reflect activation of a conductance (see later). Figure 2B shows currents in the same cell evoked by a series of voltage steps to the same test potentials from a holding potential of  $-70$  mV. Voltage pulses to potentials in the range between  $-110$  and  $-70$  mV produced time-independent currents with no rectification. On the other hand,



depolarizing voltage pulses to potentials more positive than -60 mV produced a slowly activating current that did not inactivate during the 2 sec depolarization.

Figure 2C compares the steady-state current-voltage relationships generated from holding potentials of -70 mV (open circle) and -10 mV (closed circle). The two I-V curves were nearly identical, indicating that the conductance underlying this outward current was fully open when the membrane was held at -10 mV and that it did not inactivate during the 10 sec interval between pulses. Electrophysiological properties of this  $K^+$  current in PCE cells were quite similar to those of the M-type  $K^+$  current in RPE<sup>13, 14</sup>.

#### *Selectivity.*

With 5 mM  $K^+$  in the bath, the current relaxations produced by hyperpolarizing pulses reversed between -90 and -80 mV (Figures. 2B and 3A). The reversal potential ( $V_r$ ) was analyzed by interpolation from plots of the deactivation current amplitude versus membrane voltage (not shown). For 4 cells,  $V_r$  averaged  $-84 \pm 2$  mV, which is very close to the theoretical  $E_K$  (-83 mV). To evaluate sensitivities for  $K^+$ ,  $V_r$  was also examined at higher  $[K^+]_o$ . Figure 3B shows an example of the  $K^+$  current activated in 140 mM  $K^+$  Ringer, which reversed between -7 and +3 mV. In 3 cells superfused with 140 mM  $K^+$  Ringer ( $E_K = 0$  mV),  $V_r$  averaged  $-1 \pm 3$  mV. Overall,  $V_r$  was not significantly different from  $E_K$  indicating that the time-dependent outward current is largely carried by  $K^+$ .

#### *Conductance-Voltage Relationship.*

The data in Figure 2B indicated that the M-type  $K^+$  current activated at -60 mV but at more negative potentials, the activation of a conductance could be underestimated because of its small size and slow kinetics. Therefore, we analyzed the tail current elicited by a depolarizing voltage step to a set potential from various hyperpolarized prepulses in order to determine the voltage activation range of  $K^+$  conductance. Figure

4A depicts a family of tail currents produced by voltage steps to -10 mV following 2 sec prepulses to the potentials indicated. When the prepulse potentials were in the range of -110 to -90 mV, the tail current began from the same initial level and then rose exponentially to a new steady-state. In contrast, prepulses to potentials between -80 and -50 mV produced smaller tail currents that began at a higher initial level but rose to the same steady-state. Finally, when the voltage was stepped to -10 mV from -40 and +10 mV, the initial current level was roughly the same as the steady-state current. The results of these experiments indicate that the M-type  $K^+$  current was completely deactivated at -90 mV and more negative potentials, partially activated between -80 and -30 mV, and completely activated at -20 mV and more positive potentials.

The conductance at each prepulse potential,  $G(V)$ , can be expressed as:

$$G(V) = G_{\max} - \Delta G(-10)$$

where  $G_{\max}$  is maximum conductance and  $\Delta G(-10)$  is the change in conductance produced by stepping the voltage to -10 mV, calculated by dividing the tail current amplitude by the driving force on  $K^+$  at -10 mV (= 73 mV). Similar results were obtained in 11 other cells, giving a mean value for  $G_{\max}$  of  $2.3 \pm 0.9$  nS (n=12). Figure 4B summarizes the results of these experiments and plots normalized conductance as a function of prepulse potential. The continuous curve is the least squares fit of the data to the Boltzmann function:

$$\frac{G(V)}{G_{\max}} = \frac{1}{1 + \exp\left[\frac{(V_{1/2} - V_m)}{k_n}\right]}$$

with values for  $V_{1/2}$  and  $k_n$  of -63.3 mV and 10.8 mV, respectively.

#### *Blocker Sensitivity*

To further identify channel type, we examined the sensitivity of the M-type  $K^+$  current in the PCE cells to several  $K^+$  channel blockers. M-type  $K^+$  current in the retinal pigment epithelium<sup>14, 15</sup> is known to be sensitive to  $Ba^{2+}$  and  $Cs^+$  but not to  $TEA^+$ . Figure 5A depicts families of whole-cell currents recorded in the absence (top) and the presence (bottom) of 50 mM external  $TEA^+$ .  $TEA^+$  had little effect on the amplitude or

kinetics of the time-dependent  $K^+$  current. The conductance-voltage relationship obtained from tail currents are depicted in Figure 5B, which shows that even in 50 mM TEA<sup>+</sup> the maximum conductance was 15 % smaller than it was in the absence of TEA<sup>+</sup>. The maximum  $K^+$  conductance changed to  $101 \pm 3$  % for 4 cells in the presence of 20 mM TEA<sup>+</sup>, and  $91 \pm 6$  % for other 4 cells in the presence of 50 mM TEA<sup>+</sup>. Ba<sup>2+</sup>, an effective blocker of the retinal pigment epithelium M-type  $K^+$  current, also inhibited the  $K^+$  current in PCE. Figure 6A shows that 2 mM Ba<sup>2+</sup> considerably inhibited the  $K^+$  current by 82% compared to the control. For 3 cells, the best-fit of the data to a first order equation indicates an apparent IC<sub>50</sub> of 0.38 mM (Figure 6B). Electrophysiological properties and blocker sensitivities indicate that the swelling-activated  $K^+$  current in PCE belongs to a family of KCNQ channels.<sup>18</sup> Thus, we investigated the effect of linopirdine, a specific inhibitor of KCNQ channels. Figure 6C represents families of whole-cell currents recorded in the absence (top) and the presence (bottom) of linopirdine. The  $K^+$  current component (tail currents) was almost completely blocked by 10  $\mu$ M linopirdine. In 4 cells, linopirdine (10 to 25  $\mu$ M) depolarized the resting potentials to  $-41 \pm 8$  mV from the control value of  $-59 \pm 4$  mV. Figure 6D summarizes the results of experiments in which the effects of various concentrations of linopirdine were tested on the  $K^+$  current, which indicated an apparent IC<sub>50</sub> of 0.54  $\mu$ M.

#### *Activation by Hypotonicity*

Modulation of the M-type  $K^+$  current by cell swelling was evaluated. Cell perfusion with a 20% diluted Ringer's solution ( $239 \pm 2$  mosmol/l) induced a significant increase in the  $K^+$  currents in the PCE-NPCE cell pairs as shown in Figure 7. This current augmentation by hypotonicity was reversible when the perfusate returned to the standard osmolarity (Figure 7B). In fourteen PCE-NPCE cell pairs, exposure to 20% hypotonic solution increased the maximum conductance to  $250 \pm 86$ % of the control value. Similar augmentations of *M-type  $K^+$  current* were also observed in single PCE cells ( $n = 3$ , ratio;  $184 \pm 82$ %). The reversal potential of the hypotonicity-induced

current component averaged  $-78 \pm 8$  mV ( $n=7$ ), a value more positive than the theoretical  $E_K$  ( $-83$  mV), which indicates that chloride conductance was also activated somewhat and thereby the reversal potential shifted to the direction of theoretical  $E_{Cl}$  ( $-33$  mV). The current component augmented by exposure to 20 % hypotonic solution as was blocked by linopirdine ( $20 \mu\text{M}$ ,  $n=3$ ), but essentially insensitive to external  $\text{TEA}^+$  ( $40$  mM,  $n=3$ ) as shown in Figure 7C.

## DISCUSSION

In the present study we identified a non-inactivating, outward  $\text{K}^+$  current in porcine pigmented ciliary epithelial cells that is likely different from the previously reported  $\text{K}^+$  channels in the ciliary epithelium. To date, two types of outward  $\text{K}^+$  currents have been identified in the ciliary epithelium. Voltage-activated, delayed rectifier  $\text{K}^+$  currents were reported in studies of cultured rabbit PCE cells,<sup>6</sup> cultured NPCE cells,<sup>8</sup> and fresh bovine PCE cells.<sup>7</sup> All of these currents were inhibited by  $\text{TEA}^+$ . Another type of current is the Ca-activated  $\text{K}^+$  current reported in rabbit PCE cells, which was also sensitive to  $\text{TEA}^+$ .<sup>9</sup> Carbachol-induced transient outward  $\text{K}^+$  currents described in fresh bovine PCE cells was also  $\text{Ca}^{2+}$ -sensitive,<sup>10</sup> but its sensitivities for channel blockers are unknown.

$\text{K}^+$  current in the present study closely resembled the M-type  $\text{K}^+$  current expressed in retinal pigment epithelium<sup>14,15</sup> in terms of activation kinetics, voltage dependence and blocker sensitivities. The common properties of these currents are that they are non-inactivating and activate with a threshold of  $-60$  mV or more negative voltages, and, thus, they contribute to the resting potential. In addition, they are inhibited by  $\text{Ba}^{2+}$  but relatively insensitive to  $\text{TEA}^+$ . The M-type  $\text{K}^+$  current of retinal pigment epithelium was termed from its resemblance to the M-current, a  $\text{K}^+$  current identified in certain neurons and other excitable cells. The underlying channel of the M-

current is encoded by KCNQ genes.<sup>18, 19</sup> The KCNQ/Kv7 channel family is composed of a group of five voltage-gated K<sup>+</sup> channel subunits (Kv7.1-Kv7.5) with six-transmembrane domain-spanning proteins encoded by the KCNQ genes (KCNQ1-5). The channels are formed by homo-tetramers of the same KCNQ  $\alpha$ -subunits or co-assembly of different KCNQ proteins. Linopirdine, a specific blocker of KCNQ channels, inhibited the K<sup>+</sup> channel in this study suggesting that the K<sup>+</sup> current in PCE cells belongs to KCNQ channel members. Sensitivities of KCNQ/KV7 channels to the classical K<sup>+</sup> channel blocker TEA vary by member: KCNQ1 and KCNQ2 are relatively sensitive to TEA, with an IC<sub>50</sub> ranging from 0.3 to 5.0 mM, whereas KCNQ3, KCNQ4 and KCNQ5 are substantially insensitive to TEA, with an IC<sub>50</sub> of 30 mM or more.<sup>18, 20</sup> The K<sup>+</sup> channel in this study was little affected even by treatment with 50 mM TEA; hence, it is likely to belong to the latter members.

One of the physiological roles of this K<sup>+</sup> channel in PCE cells is likely to consist of a major K<sup>+</sup> conductance of the apical and/or basolateral membranes, which is supported by the fact that the channel is open near the resting potential and that linopirdine, a specific blocker of this K<sup>+</sup> channel depolarized the zero-current potential of the PCE cells ( $-59 \pm 4$  to  $-41 \pm 8$  mV). Another role of this K<sup>+</sup> channel in PCE cells may be to mediate cell volume and water transport. As with the K<sup>+</sup> channel in this study, some KCNQ channels have been reported to be activated by cell swelling. For instance, the KCNQ1 channel, which is expressed in secretory epithelia, like the distal colon or airway, is regulated by small changes in cell volume and participates in the secretion of salt and water.<sup>21</sup> In addition, recent evidence suggested that KCNQ3 and KCNQ5 also could mediate basal anion secretion from an airway epithelial cell line<sup>22</sup> and regulate Na<sup>+</sup> transport across lung epithelial cells.<sup>23</sup>

In conclusion, we identified a swelling-activated K<sup>+</sup> channel in freshly dissociated porcine pigmented ciliary epithelial cells. Electrophysiological and

pharmacological properties of this channel indicate that it is a member of the KCNQ/Kv7 channel family. This K<sup>+</sup> channel is active near resting potentials and may contribute to the regulation of cell volume and water transport via the ciliary epithelia.

## References

1. Do CW, Civan MM. Swelling-activated chloride channels in aqueous humor formation: on the one side and the other. *Acta Physiol.* 2006;187:345-52.
2. Do CW, Civan MM. Basis of chloride transport in ciliary epithelium. *J Membr Biol.* 2004 ;200:1-13.
3. Coca-Prados M, Sánchez-Torres J, Peterson-Yantorno K, Civan MM. Association of ClC-3 channel with Cl<sup>-</sup> transport by human nonpigmented ciliary epithelial cells. *J Membr Biol.* 1996 ;150:197-208.
4. Do CW, Civan MM. Species variation in biology and physiology of the ciliary epithelium: similarities and differences. *Exp Eye Res.* 2009; 88:631-640.
5. Moser SL, Harron SA, Crack J, Fawcett JP, Cowley EA. Multiple KCNQ potassium channel subtypes mediate basal anion secretion from the human airway epithelial cell line Calu-3. *J Membr Biol.* 2008;221:153-63.
6. Fain GL, Farahbakhsh NA. Voltage-activated currents recorded from rabbit pigmented ciliary body epithelial cells in culture. *J Physiol.* 1989;418:83-103.
7. Jacob TJ. Two outward K<sup>+</sup> currents in bovine pigmented ciliary epithelial cells: IK(Ca) and IK(V). *Am J Physiol.* 1991;261:C1055-62.
8. Cilluffo MC, Cohen BN, Fain GL. Nonpigmented cells of the rabbit ciliary body epithelium. Tissue culture and voltage-gated currents. *Invest Ophthalmol Vis Sci.* 1991;32:1619-29.
9. Ryan JS, Tao QP, Kelly ME. Adrenergic regulation of calcium-activated potassium current in cultured rabbit pigmented ciliary epithelial cells. *J Physiol.* 1998; 511:145-57.

10. Stelling JW, Jacob TJ. Transient activation of K<sup>+</sup> channels by carbachol in bovine pigmented ciliary body epithelial cells. *Am J Physiol.* 1996;271:C203-9.
11. Stelling JW, Jacob TJ. The inward rectifier K<sup>+</sup> current underlies oscillatory membrane potential behavior in bovine pigmented ciliary epithelial cells. *J Physiol.* 1992;458:439-56.
12. Hughes BA, Takahira M. Inwardly rectifying K<sup>+</sup> currents in isolated human retinal pigment epithelial cells. *Invest Ophthalmol Vis Sci.* 1996;37:1125-39
13. Shimura M, Yuan Y, Chang JT, Zhang S, Campochiaro PA, Zack DJ, Hughes BA. Expression and permeation properties of the K(+) channel Kir7.1 in the retinal pigment epithelium. *J Physiol.* 2001;531:329-46.
14. Hughes BA, Takahira M, Segawa Y. An outwardly rectifying K<sup>+</sup> current active near resting potential in human retinal pigment epithelial cells. *Am J Physiol.* 1995;269:C179-87.
15. Takahira M, Hughes BA. Isolated bovine retinal pigment epithelial cells express delayed rectifier type and M-type K<sup>+</sup> currents. *Am J Physiol.* 1997;273:C790-803.
16. Takahira M, Sakurada N, Segawa Y, Shirao Y. Two types of K<sup>+</sup> currents modulated by arachidonic acid in bovine corneal epithelial cells. *Invest Ophthalmol Vis Sci.* 2001 ;42:1847-54.
17. Walker VE, Stelling JW, Miley HE, Jacob TJ. Effect of coupling on volume-regulatory response of ciliary epithelial cells suggests mechanism for secretion. *Am J Physiol.* 1999 ;276:C1432-8.
18. Robbins J. KCNQ potassium channels: physiology, pathophysiology, and pharmacology. *Pharmacol Ther.* 2001 ;90:1-19.
19. Brown DA. Kv7 (KCNQ) potassium channels that are mutated in human diseases. *J*



Physiol. 2008 ;586:1781-3.

20. Bal M, Zhang J, Zaika O, Hernandez CC, Shapiro MS. Homomeric and heteromeric assembly of KCNQ (Kv7) K<sup>+</sup> channels assayed by total internal reflection fluorescence/fluorescence resonance energy transfer and patch clamp analysis. *J Biol Chem.* 2008;283:30668-76.
21. Grunnet M, Jespersen T, MacAulay N, Jørgensen NK, Schmitt N, Pongs O, Olesen SP, Klaerke DA. KCNQ1 channels sense small changes in cell volume. *J Physiol.* 2003;549:419-27.
22. Moser SL, Harron SA, Crack J, Fawcett JP, Cowley EA. Multiple KCNQ potassium channel subtypes mediate basal anion secretion from the human airway epithelial cell line Calu-3. *J Membr Biol.* 2008;221:153-63.
23. Greenwood IA, Yeung SY, Hettiarachi S, Andersson M, Baines DL. KCNQ-encoded channels regulate Na<sup>+</sup> transport across H441 lung epithelial cells. *Pflugers Arch.* 2009 ;457:785-94.

## Figure legends

### FIGURE 1

Photomicrograph of isolated porcine ciliary epithelial cells.

A pair of pigmented and non-pigmented ciliary epithelial cells (PCE and NPCE cells) freshly isolated from porcine eyes. In this experiment, a PCE cell membrane was clamped by a patch-pipette. Bar, 50  $\mu\text{m}$ .

### FIGURE 2

Whole-cell current in porcine ciliary epithelium and its dependence on holding potential.

(A) Whole-cell currents in a porcine PCE-NPCE pair were elicited by 1s-voltage steps ranging from +10 to -110 mV in -10 mV increments from a holding potential (HP) of -10 mV. (B) In the same cell, currents were elicited by same range voltage steps from a HP of -70 mV. (C) I-V relationships are depicted with a HP of -10 (●) and -70 mV (○). The steady-state current amplitudes were averaged between 900 and 1000 ms in the whole-cell currents in A.

### FIGURE 3

Reversal potential of the time-dependent  $\text{K}^+$  current and its relationship to extracellular  $\text{K}^+$  concentration.

(A) Superfused with standard Ringer solution ( $[\text{K}^+]_o = 5 \text{ mM}$ ), currents elicited by membrane voltage steps ranging from -100 to -70 mV from a holding potential of -10 mV. The polarity of the deactivating currents reversed between -90 and -80 mV. (B)

Superfused with a high  $K^+$  solution ( $[K^+]_o = 140$  mM),  $K^+$  currents elicited by membrane voltage steps ranging from  $-37$  to  $+33$  mV from a holding potential of  $-107$  mV. In  $140$  mM  $[K^+]_o$  solution, the activating currents reversed between  $-7$  and  $+3$  mV.

#### FIGURE 4

Voltage dependence of the  $K^+$  conductance.

(A) Tail current activation at  $-10$  mV following prepulses to the potentials indicated. Tail current amplitude is inversely proportional to the fraction of  $K^+$  channels activated at prepulse potential. (B) Conductance-voltage relationship.  $K^+$  conductance at each voltage is normalized to maximum conductance. Data represents mean  $\pm$  SEM for 12 cells. The continuous curve is the least squares fit of the data to a Boltzmann function:

$$\frac{G(V)}{G_{\max}} = \frac{1}{1 + \exp[(V_{1/2} - V_m)/k_n]}, \quad \text{where } V_{1/2} \text{ and } k_n \text{ are } -63.3 \text{ mV and } 10.8 \text{ mV,}$$

respectively.

#### FIGURE 5

Effect of TEA on the  $K^+$  current. (A) Families of whole-cell currents recorded in the absence (top) and the presence (bottom) of  $50$  mM TEA. Currents were elicited by membrane voltage steps ranging from  $+10$  to  $-130$  mV from a holding potential of  $-10$  mV. (B) Effect of TEA on steady-state conductance-voltage relationship. The  $K^+$  conductance in the absence (closed circle) and the presence (open circle) of  $50$  mM TEA was calculated from tail currents in A.

#### FIGURE 6

Inhibitory effects of  $\text{Ba}^{2+}$  and linopirdine on the  $\text{K}^+$  channels. (A) Whole cell currents elicited by voltage steps ranging from +10 to -130 mV were obtained in the absence and the presence of 2 mM  $\text{Ba}^{2+}$ . (B) Dose response curve. Relative current amplitude at each concentration was measured and summarized in 3 cells. Smooth curves represent fitting the data to the first order equation:  $I_{\text{Inhibitor}} / I_{\text{control}} = 1 - \{[\text{Inhibitor}] / ([\text{Inhibitor}] + \text{IC}_{50})\}$ , with an  $\text{IC}_{50}$  of 0.38 mM. All error bars represent SEM. (C) Whole cell currents elicited by voltage steps ranging from +10 to -110 mV were obtained in the absence and the presence of 10  $\mu\text{M}$  linopirdine. (D) Dose response curve. Each data point depicts measurements in 3 cells. Smooth curves represent fitting the data to the same equation above, where the  $\text{IC}_{50}$  was 0.54  $\mu\text{M}$ . All error bars represent SEM.

## FIGURE 7




Effect of hypotonicity on the  $\text{K}^+$  current.

(A) Families of whole-cell currents recorded when a cell was superfused with standard Ringer's solution ( $302 \pm 6$  mosmol/l, left panel) and 20% diluted hypotonic solution ( $239 \pm 2$  mosmol/l, right panel). Currents were elicited by membrane voltage steps ranging from +10 to -120 mV from a holding potential of -10 mV. (B) Time course of the M-type  $\text{K}^+$  current. Maximum amplitude of the tail currents was monitored in the same cell in A. (C) Effects of high TEA and linopirdine in hypotonic condition. In another cell, families of whole cell current were recorded in standard Ringer's solution, 20% diluted hypotonic solution, hypotonic (20% dilution) 40 mM TEA solution, and 20  $\mu\text{M}$  linopirdine in hypotonic solution. Currents were elicited by membrane voltage steps ranging from +10 to -110 mV from a holding potential of -10 mV.


précis summary

A swelling-activated potassium channel was identified in freshly dissociated porcine pigmented ciliary epithelial cells. This potassium channel may be a member of the KCNQ/Kv7 and could contribute to the regulation of water transport via the ciliary epithelia.





A



B

