

# Pre-Miocene Paleogeographic Linkage Between e Korean peninsula and the Japanese Islands

メタデータ	言語: eng 出版者: 公開日: 2017-10-05 キーワード (Ja): キーワード (En): 作成者: メールアドレス: 所属:
URL	<a href="http://hdl.handle.net/2297/6361">http://hdl.handle.net/2297/6361</a>

# Pre-Miocene Paleogeographic Linkage Between the Korean Peninsula and the Japanese Islands

Yong Il LEE

*School of Earth and Environmental Sciences, Seoul National University NS80, Seoul 151-747, KOREA*

**Abstract** – At present the Japanese Islands are separated from the East Asia continent by the East/Japan Sea. This study presents five pieces of geological evidence indicating that prior to an opening of the East/Japan Sea the Japanese Islands were connected to the East Asian continent through the Korean Peninsula. They are 1) Precambrian rocks in the Oki-Hida belt, Japan, 2) orthogneiss and limestone clasts in Kamiasso Conglomerate in the Mino terrane, Japan, 3) distribution of dinosaur fossils in Korea and Japan, 4) quartzite gravels in Japan and 5) radiolaria-bearing chert pebbles in Early Cretaceous sediments in Korea.

## I. Introduction

The East/Japan Sea is situated between the East Asia continent and Japanese Islands separating Japanese Islands from the continent (Fig. 1). The location of the Japanese Islands near the Asian continent during the Cretaceous with subsequent southward drift in the Tertiary is generally accepted [1, 2, 3]. Southward drift is explained by differential rotation of Southwest and Northeast Japan based on paleomagnetic studies [1, 3] and by strike-slip fault systems initiated in the Eocene [2]. Southwest Japan was rotated clockwise through about 45° with respect to the Korean Peninsula and Northeast Japan was rotated about 40° [4]. This differential rotation occurred concurrently at about 15Ma [4]. According to evidence from the East/Japan Sea including geochronological data [5], marine magnetic anomaly patterns [6, 7, 8] and ODP Leg 127 drilling results [5, 9, 10, 11, 12, 13] the Japan Arc rifted from the Asian continent due to an opening of the East/Japan Sea, and was drifted to the present position later than 19Ma.

The Korean Peninsula and Japanese Islands have quite different geological histories. Nonetheless both regions share several lines of geological evidence indicating that they were once connected to each other prior to an opening of the East/Japan Sea that occurred in the Middle Miocene. This presentation is not concerned with the timing and mode of the opening of the East/Japan Sea, but is concerned with the paleogeography of pre-Miocene time, including how the margins of the rifted arcs are fitted to continental margins before drift. Five pieces of geological evidence will be provided to relate Japanese Islands with East Asia continent through the Korean Peninsula. The areas mentioned in this presentation are shown in Fig. 1.

## II. The Oki-Hida Belt, Japan

### A. The Oki belt

The Oki belt comprises 2.0 Ga to 250 Ma medium-pressure-type gneiss and granite complex [13]. The highest metamorphic grade reaches the upper amphibolite to granulite facies. Protolith is continental sedimentary rocks with minor amount of mafic igneous rocks [14]. The Oki belt is considered to be of continental affinity and is at present isolated from mainland Asia, as it was rifted and detached by the Miocene opening of a back-arc basin, the East/Japan Sea [4, 15]. Judging from the lithologic and chronological similarity, the Oki belt is regarded as an eastern extension of the Precambrian rocks of the Sobaeksan (Yeongnam) massif in South Korea [16, 17].

### B. The Hida belt

The Hida belt in central Japan appears to extend westwards to Oki Island and comprises various types of medium-pressure-type metamorphic and igneous rocks with ages from Precambrian (?) to Mesozoic. Non- to weakly metamorphosed middle to upper Paleozoic shelf sequences occur fragmentally in the periphery of the belt. Many Japanese geologists have long recognized the notion that the Hida gneiss region contains Precambrian basement rocks formed in the southern margin of the Sino-Korean (North China) block (e.g., [18]). Some studies considered that the Hida belt was a microcontinent that collided with the eastern edge of the Sino-Korean block in the late Paleozoic [19, 20, 21]. Structural studies revealed that the Hida belt in the craton margin later thrust southward onto the Permian-Jurassic accretionary complexes, forming a large nappe [16, 22, 23]. The existence of medium-pressure type metamorphism at 250-220 Ma in the Hida belt was interpreted to be a consequence of collision between Sino-Korean and South China blocks [16, 24]. In contrast to common belief, the majority of radiometric age data obtained by various methods indicate Paleozoic to Mesozoic ages, and Precambrian age data are quite limited. Suzuki and Adachi reported CHIME (Chemical Th-U-total Pb isochron) ages ranging from 2000 to 300 Ma for zircon grains in the Hida gneisses and interpreted that the Hida gneiss was not metamorphosed in Precambrian time but in 250 Ma, and that the older isotopic dates are ages of detrital zircons [13]. A recent




Fig. 1. Map showing tectonic divisions of the Korean Peninsula and Japanese Islands described in the text. 1: Oki/Hida belt, 2: Kamiasso Conglomerate, 3: Dinosaur fossils, 4: Quartzites and 5: Radiolaria-bearing chert pebbles.

study by Arakawa, Saito and Amakawa identified three main stages of igneous activities in the Hida belt: mafic meta-volcanic rocks (amphibolites) at late Silurian, mafic dikes at Middle Carboniferous and granite-diorite plutons at Late Triassic to Early Jurassic, all formed at the continental margin or continental arc tectonic environment [25]. They proposed two possible models for the Paleozoic to early Mesozoic formation and evolution of the Hida belt. One is that the Hida belt was formed in an eastern margin of the Sino-Korean block (northern part of the Korean Peninsula) and later developed as the eastern extension of the suture zone between the Sino-Korean and South China blocks. The unique occurrence of middle to late Paleozoic shelf strata with Boreal fauna in the periphery of the Hida belt [26, 27] suggests a strong link between the Hida belt and Sino-Korean block. The other is the model in which the Hida belt was developed as a part of the vast suture zone (East Central Asian Orogenic Belt) to the north, which is situated between the Siberian and Sino-Korean blocks.

### III. Kamiasso Conglomerate, Central Japan

Some intraformational conglomerates are known to occur in the Mino terrane, a Jurassic accretionary complex; one of the best studied examples is the Kamiasso conglomerate exposed along the Hida River gorge in the Kamiasso area, north of Nagoya City. The Kamiasso conglomerate is known as the oldest-rock-bearing conglomerate in Japan. The Kamiasso conglomerate consists of four conglomerate beds within the turbidite [28]. Clasts of the conglomerate are angular to well-rounded pebbles, cobbles and boulders of sedimentary, igneous and metamorphic rocks in coarse-grained graywacke matrix, of which sedimentary clasts of sandstone, shale, chert limestone and marl are most common. The Kamiasso conglomerate is interpreted to be submarine debris-flow deposits [29]. Paleocurrent analyses of the turbidites in the Kamiasso conglomerate indicate that the clastic materials were transported from the north [30, 31]. Among various clasts orthogneiss and limestone clasts are dealt here for consideration.

### A. *Orthogneiss clasts*

Orthogneiss clast comprises about 10% of the total clasts [28]. A Rb-Sr whole-rock isochron age of  $2050 \pm 30$  Ma has been reported by Shibata and Adachi [32]. A well-rounded zircon grain yields an exceptionally old CHIME age of  $3040 \pm 180$  Ma, but most euhedral zircons of magmatic origin are concentrated around 2050 Ma [33]. CHIME ages of many monazite grains are between 1750 and 1500 Ma, but rarely a narrow portion of several monazite cores gives a CHIME age of  $2670 \pm 100$  Ma [33]. The ages of the 2050 Ma isochron age for the orthogneiss clasts closely correspond to the Rb-Sr whole-rock isochron age ( $2060 \pm 50$  Ma) of Imwon granite gneiss, central eastern Korea (northeastern end of the Yeongnam massif) whose age was recalculated by Adachi and Suzuki [33] from the data of Kim, Joo and Jo [34].

### B. *Limestone clasts*

Some limestone clasts of the Kamiaso conglomerate contain Middle Carboniferous fusulinaceans: *Fusulinella laxa*, *Eoschubertella obscura*, *Pseudoendothyra* sp., *Profusulinella* cf. *prisca timanica*, *Profusulinella* cf. *fukujiensis*, *Pseudostaffella kremisi* and *Verella* sp. [33]. The fusulinacean faunas are similar to those of limestone beds of the Pyeongan Supergroup in the Yeongweol and Samcheok coalfields, eastern central Korea, rather than those of the Hida Marginal Belt [33, 35].

## IV. Dinosaur Fossils in Korea and Japan

Dinosaurs are terrestrial animals and thus the occurrence of dinosaur fossils in Japanese Islands indicates that the Japanese Islands were connected to the continent during the time when the dinosaurs roamed on the Earth surface. In this section, the occurrence of dinosaur fossils in Korea and Japan will be reviewed.

### A. *Korea*

Abundant dinosaur fossils including dinosaur footprints, eggs and nests, teeth and bones have been found from the Cretaceous non-marine deposits of Korea. Dinosaur bone fossils were found in fluvial deposits of Early Cretaceous age. Characteristically, preservation of dinosaur bone fossils was aided by calcareous pedogenesis. Dinosaur tracks are the most distinctive, and some tracksites are among the most famous in the world. Until now, 27 dinosaur track localities have been discovered from Upper Cretaceous nonmarine deposits including the Gyeongsang Basin and several small basins, and mainly dinosaur tracks have been found from the coastal areas in southern part of the Korean Peninsula [36]. Dinosaur tracks are of ornithopods, theropods,

pterosaurs and sauropods. Dinosaur tracks usually occur in the lacustrine deposits in which thinly interlaminated fine-grained sandstones and siltstones-mudstones are prevailing. Dinosaur eggshell fragments were reported from Lower Cretaceous deposits, but several complete fossil eggs and clutches were recently found in Upper Cretaceous deposits (e.g., [37]).

### B. *Japan*

Many dinosaur body and footprint fossils have been found at various localities in Japan. Japanese dinosaur fossils, except in a very few cases, are mostly isolated teeth and bones [38]. All dinosaur fossils are found in Cretaceous strata. The dinosaur localities other than the Tetori Group, Kanmon Group, Mifune Group and Goshoura Group are located within the Outer Zone of the Japanese tectonic divisions. The localities in the Outer Zone are of shallow marine sediments, and the dinosaur fossils are allochthonous. On the other hand, the Tetori, Kanmon, Mifune and Goshoura groups are terrestrial sediments, and dinosaur fossils are more or less autochthonous. Also, dinosaur fossils from these groups are all from the Early Cretaceous. Among these groups, the Tetori Group, which ranges from the middle Jurassic to Early Cretaceous of central Japan, has produced most of the dinosaur fossils found in Japan to date [38]. Dinosaur fossils have been found from the upper parts of the Tetori Group. They include theropods, sauropods, and ornithopods, and important members include a dromaeosaurid and an iguanodontid [38]. Co-occurring taxa include crocodiles, turtles, ganoid fishes and birds. The dinosaur fauna of the Tetori Group is Early Cretaceous in age, and it represents a fauna that lived in wet lowland area in a warm, humid climate located on the Far East Asian continental edge [38].

## V. Quartzite Gravels in Japan

There are many strata containing quartzite gravels across the Japanese Islands. The question then arises as to the provenance of the quartzite gravels. In the present Japanese Islands there have been discovered no such matured quartzite rocks [39]. Accordingly, the source of the quartzite gravels in Japanese Islands must be thought in some other areas. The quartzitic rocks from which the quartzitic gravels were derived must have been originally formed in eolian to beach environments in a continental setting. Actually, orthoquartzite is one of the most typical cratonic sedimentary rocks. In addition, quartzose rocks are more common and abundant in the Proterozoic to lower Paleozoic due to the tectonically stable continental blocks superimposed by intense weathering and strong abrading transport agents such as wind, surf, and longshore currents. The lack of land plants cover until the Silurian or Devonian also must have helped to produce these supermature quartz arenites, which virtually disappear




Fig. 2. The simplified map showing the distribution of quartzite types in the southern Korean Peninsula [42]. IB: Imjingang belt, GM: Gyeonggi massif, OB: Okcheon belt, YM: Yeongnam massif and GB: Gyeongsang Basin.

after the early Paleozoic. As an example this study presents some results of a study on quartzite gravels occurring in the Tetori Group in central Japan (the Inner Zone).

The Tetori Group is a representative Middle Jurassic to Lower Cretaceous deposit in central Japan. It is subdivided into three subgroups, namely, the Kuzuryu, Itoshiro and Akaiwa subgroups with decreasing age. The depositional environments of the Tetori Group changed with time from shallow marine environments (Kuzuryu Subgroup) to brackish environments (Itoshiro Subgroup) and to nonmarine environments (Akaiwa Subgroup) [40]. Quartzite gravels are abundantly included in conglomerates of the basal parts of the Itoshiro and Akaiwa subgroups. They are predominantly subrounded to rounded pebbles and cobbles, and are pink, reddish purple, gray, white and bluish gray in color. Considering no

existence of quartzite strata in Japan the quartzite clasts of the Tetori Group are commonly interpreted to have been derived from Precambrian quartzite in East Asia: the Sinian System of China and the Sangweon System of North Korea (e.g., [41]). However, textural observation suggests that they were not derived from Precambrian quartzites of China and North Korea but from Precambrian and lower Paleozoic quartzites of South Korea.

The Korean Peninsula is composed of variably aged rocks from the Precambrian to the recent. Quartzite beds occur only in Precambrian and Paleozoic sequences. In the southern peninsula twelve quartzite strata occur in the Gyeonggi massif and Okcheon belt (Fig. 2). All quartzites in the Gyeonggi massif are of Precambrian in age and are characterized by high-grade metaquartzites, whereas those in the Okcheon belt are diverse from

orthoquartzite to medium-grade metaquartzite [42]. Comparison of textural characteristics of Tetori Group quartzite gravels suggests that the most probable source rocks for the Tetori quartzite gravels would be the Lower Cambrian Jangsan Formation, distributed in eastern central Korea (NE Okcheon belt), as well as Late Proterozoic quartzose strata, which was newly discovered in the same area recently [43]. The paleocurrent data indicate that the Kamiaso conglomerates were derived from the north [30]. The proposed source area agrees with the paleocurrent data after correction of the clockwise rotation of Southwest Japan.

## VI. Radiolaria-bearing Chert Pebbles in Cretaceous Sediments in Korea

The Lower Cretaceous Gyeongsang Supergroup distributed in the southeastern Korean Peninsula contains three strata having chert-pebble conglomerate beds. The Gyeongsang Supergroup is composed of sediments deposited in nonmarine environments including alluvial fan, fluvial and lacustrine settings. The chert pebbles in conglomerates contain many radiolarian faunas ranging in age from Late Permian to Late Jurassic [44, 45, 46]. In the Korean Peninsula Permian to Jurassic marine chert sequences are not exposed. The directions of paleocurrent for the sediments containing radiolaria-bearing chert pebbles are mainly from NE to SW [44, 47]. It was revealed that the radiolarian faunas are similar to those of Jurassic accretionary complexes of Southwest Japan such as the Mino-Tamba-Ashio and Chichibu terranes [44, 45, 46]. Based on terrane analysis [48], the Mino-Tamba belt was located northeast of the Gyeongsang Basin during Cretaceous time, which indicates that the paleo-drainage was connected between the nonmarine Gyeongsang Basin, Korea and the Mino-Tamba-Ashio terrane in Japan. Also, the occurrence of chert pebbles derived from the Jurassic accretionary complexes suggests that Permian accretionary complexes, which are supposed to have been located between the Gyeongsang Basin and the Jurassic accretionary complexes, were peneplained and thus, they did not act as a possible sediment source. Or the Jurassic accretionary complexes might have covered more widely over the Permian accretionary complexes and the present distribution of the Jurassic accretionary complexes may be resulted from exhumation. The erosion of the Jurassic accretionary complexes is known to have started during Late Jurassic time at least in the Outer Zone and possibly in the Inner Zone and became intensified during the Early Cretaceous [49].

### Acknowledgements

This research was supported by the Korea Science and Engineering Foundation (R01-2000-000-00056-0) and by the Korea Research Foundation (DS0071). Thanks to Prof. Shoji Arai for

invitation to the Kanazawa International Symposium and Dr. Ken-ichiro Hisada for help in the field and research collaboration.

### References

- [1] A. Hayashida, Y. Ito, "Paleoposition of Southwest Japan at 16 Ma: Implication from paleomagnetism of the Miocene Ichishi Group," *Earth and Planetary Science Letters* Vol. 68, pp. 335-342, 1984.
- [2] S. Lallemand, L. Jolivet, "Japan Sea: a pull-apart basin?," *Earth and Planetary Science Letters* Vol. 76, pp. 375-389, 1986.
- [3] Y. Otofujii, T. Itaya, T. Matsuda, "Rapid rotation of Southwest Japan: Paleomagnetism and K-Ar ages of Miocene volcanic rocks of Southwest Japan," *Geophysical Journal International* Vol. 105, pp. 397-405, 1991.
- [4] Y. Otofujii, "Large tectonic movement of the Japan Arc in late Cenozoic times inferred from paleomagnetism: Review and synthesis," *The Island Arc* Vol. 5, pp. 229-249, 1996.
- [5] I. Kaneoka, K. Notsu, Y. Takigami, K. Fujioka, H. Sakai, "Constraints on the evolution of the Japan Sea based on 40Ar-39Ar ages and Sr isotopic ratios for volcanic rocks of the Yamato Seamount chain in the Japan Sea," *Earth and Planetary Science Letters* Vol. 97, pp. 211-225, 1990.
- [6] N. Isezaki, "Possible spreading centers in the Japan Sea," *Marine Geophysical Research*, Vol. 2, pp. 265-277, 1975.
- [7] N. Isezaki, "A magnetic anomaly map of the Japan Sea," *Journal of Geomagnetism and Geoelectricity* Vol. 38, pp. 403-410, 1986.
- [8] N. Seama, N. Isezaki, "Sea-floor magnetization in the eastern part of the Japan Basin and its tectonic implication," *Tectonophysics* Vol. 181, pp. 285-297, 1990.
- [9] Scientific Drilling Party Leg 127, "Exploring the Japan Sea," *Geotimes* Vol. 34, pp. 19-21, 1989.
- [10] Shipboard Scientific Drilling Party Leg 127 and 128, "Evolution of the Japan Sea," *Nature* Vol. 346, pp. 18-20, 1990.
- [11] J. C. Ingle, Jr., "Subsidence of the Japan Sea: Stratigraphic evidence from ODP sites and onshore sections," In: K. Tamaki, K. Suyehiro, J. Allan et al. (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results*, Vol. 127/128, pp. 1197-1218. College Station, Texas, 1992.
- [12] L. Jolivet, K. Tamaki, "Neogene kinematics in the Japan Sea region and volcanic activity of the Northeast Japan arc," In: K. Tamaki, K. Suyehiro, J. Allan et al. (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results*, Vol. 127/128, pp. 1311-1331. College Station, Texas, 1992.
- [13] K. Suzuki, M. Adachi, "Middle Precambrian detrital monazite and zircon from the Hida gneiss on Oki-Dogo Island, Japan: Their origin and implications for the correlation of basement gneiss of Southwest Japan and Korea," *Tectonophysics* Vol. 235, pp. 277-292, 1994.
- [14] Y. Isozaki, "Anatomy and genesis of a subduction-related orogen: A new view of geotectonic subdivision and evolution of the Japanese Islands," *The Island Arc* Vol. 5, pp. 289-320, 1996.
- [15] L. Jolivet, K. Tamaki, M. Fournier, "Japan Sea, opening history and mechanism: A synthesis," *Journal of Geophysical Research* Vol. 99B, pp. 22237-22259, 1994.
- [16] T. Sohma, L. Kunugiza, M. Terabayashi, "Hida metamorphic belt," *Excursion Guidebook of the 96<sup>th</sup> Annual Meeting of Geological Society of Japan* pp. 27-53 (in Japanese with English abstract), 1990.
- [17] Y. Isozaki, S. Maruyama, "Studies on orogeny based on plate tectonics in Japan and new geotectonic subdivision of the Japanese Islands," *Journal of Geography* Vol. 100, pp. 697-761 (in Japanese with English abstract), 1991.
- [18] T. Kobayashi, "The Sakawa orogenic cycle and its bearing on the origin of the Japanese Islands," *Journal of Faculty of Science, Imperial University of Tokyo Section II*, pp. 219-278, 1941.
- [19] S. Mizutani, I. Hattori, "Hida and Mino tectonostratigraphic terranes in central Japan," In: M. Hashimoto, S. Uyeda (Eds.), *Accretion Tectonics in the Circum-Pacific Regions*, Terra Science, Tokyo, pp. 169-178, 1983.
- [20] W. G. Ernst, R. Cao, J. Jiang, "Reconnaissance study of Precambrian

- metamorphic rocks, northeastern Sino-Korean shield, People's Republic of China," *Geological Society of America Bulletin* Vol. 100, pp. 692-701, 1988.
- [21] S. Maruyama, J. G. Liou, T. Seno, "Mesozoic and Cenozoic evolution of Asia," In: Z. Ben Avraham (Ed.), *The Evolution of the Pacific Ocean Margins*, Oxford Monograph in Geology and Geophysics Volume 8, p. 75-99, 1989.
- [22] M. Komatsu, K. Suwa, "The structure and tectonics of pre-Jurassic serpentinite mélangé and overlying Hida nappe in central Japan," Abstracts of the International Symposium on Pre-Jurassic East Asia. IGCP 224, pp. 97-103, 1986.
- [23] M. Komatsu, M. Nagase, K. Naito, T. Kanno, M. Ujibara, T. Toyoshima, "Structure and tectonics of the Hida massif, central Japan," *Memoir of Geological Society of Japan* No. 42, pp. 39-62 (in Japanese with English abstract), 1993.
- [24] T. Sohma, K. Kunugiza, "The formation of the Hida nappe and the tectonics of Mesozoic sediments: the tectonic evolution of the Hida region, central Japan," *Memoir of Geological Society of Japan* No. 42, pp. 1-20, 1993.
- [25] Y. Arakawa, Y. Saito, H. Amakawa, "Crustal development of the Hida belt, Japan: Evidence from Nd-Sr isotopic and chemical characteristics of igneous and metamorphic rocks," *Tectonophysics* Vol. 328, pp. 183-204, 2000.
- [26] H. Igo, "Paleozoic strata in the Hida 'Gaien' belt," In: K. Ichikawa, S. Mizutani, I. Hara, S. Hada, A. Yao (Eds.), *Pre-Cretaceous Terranes of Japan*, Publication of IGCP #224, Osaka, pp. 41-48, 1990.
- [27] M. Kato, "Paleozoic corals," In: K. Ichikawa, S. Mizutani, I. Hara, S. Hada, A. Yao, (Eds.), *Pre-Cretaceous Terranes of Japan*, Publication of IGCP #224, Osaka, pp. 307-312, 1990.
- [28] M. Adachi, "Permian intraformational conglomerate at Kamiasso, Gifu Prefecture, central Japan," *Journal of Geological Society of Japan* Vol. 77, pp. 471-482, 1971.
- [29] N. Tachibana, H. Okada, H. Sano, "Sedimentary facies, mineral composition of sandstones and depositional setting of Jurassic clastic rocks of the Mino terrane, Southwest Japan," *Memoir of Geological Society of Japan* No. 38, pp. 121-133, 1992.
- [30] M. Adachi, "Paleogeographic aspects of the Japanese Paleozoic-Mesozoic geosyncline," *Journal of Earth Science, Nagoya University* Vol. 23/24, pp. 13-55, 1976.
- [31] M. Adachi, S. Mizutani, "Sole markings and paleocurrent system in the Paleozoic group of the Mino terrane, central Japan," *Memoir of Geological Society of Japan* No. 6, pp. 39-48 (in Japanese with English abstract), 1971.
- [32] K. Shibata, M. Adachi, "Rb-Sr whole-rock ages of Precambrian metamorphic rocks in the Kamiasso conglomerate, central Japan," *Earth and Planetary Science Letters* Vol. 21, pp. 277-287, 1974.
- [33] M. Adachi, K. Suzuki, "Were Precambrian gneiss clasts in the Kamiasso conglomerate derived from the eastern Korean Peninsula?," *Bulletin of Nagoya University Furukawa Museum* Vol. 9, pp. 25-45, 1993.
- [34] Y. J. Kim, S. H. Joo, D. R. Jo, "Rb/Sr of leucogranitic rocks in the Sameheog area," *Journal of Korean Institute of Mining Geology*, Vol. 17, pp. 231-236, 1984 (in Korean with English abstract).
- [35] K. Wakita, S. Kojima, K. Tsukada, *Middle Mesozoic accretionary complex of the Mino terrane, and Paleozoic to Mesozoic sedimentary rocks of the Hida marginal belt*, ISRGA Field Workshop (FW-C1), pp. 165-235, 2001.
- [36] M. Huh, K. G. Hwang, I. S. Paik, C. H. Chung, "Dinosaur tracks from the Cretaceous of South Korea: distribution, occurrences and paleobiological significance," *The Island Arc* Vol. 12, in press.
- [37] M. Huh, D. K. Zelenitsky, "Rich dinosaur nesting site from the Cretaceous of Bosung County, Chullanam-do Province, South Korea," *Journal of Vertebrate Paleontology* Vol. 22, pp. 716-718, 2002.
- [38] Y. Azuma, Y. Tomida, "Japanese Dinosaurs," In: P. J. Currie, K. Padian (Eds.), *Encyclopedia of Dinosaurs*, New York, Academic Press, pp. 375-379, 1997.
- [39] T. Tokuoka, "The Simanto Terrain in the Kii Peninsula, Southwest Japan-with special reference to its geologic development viewed from coarser clastic sediments," *Memoir of Faculty of Science, Kyoto University Series in Geology and Mineralogy* Vol. XXXIV, pp. 35-74, 1967.
- [40] S. Maeda, "On the geological history of the Mesozoic Tetori Group in Japan," *Journal of College of Arts and Sciences, Chiba University* Vol. 3, pp. 369-426 (in Japanese with English abstract), 1961.
- [41] M. Fujita, "A new contribution to the stratigraphy of the Tetori Group, adjacent to Lake Kuzuryu, Fukui Prefecture, central Japan," *Memoir of the Fukui Prefectural Dinosaur Museum* No. 1, pp. 41-53, 2002.
- [42] Y. Kim, Y. I. Lee, "Characterization of quartzites in the southern Korean Peninsula," *Gondwana Research* Vol. 6, in press.
- [43] Y. Kim, Y. I. Lee, "Discovery of a new late Proterozoic stratum in South Korea," *Geosciences Journal* Vol. 7, in press.
- [44] K. H. Chang, B. G. Woo, J. H. Lee, S. O. Park, A. Yao, "Cretaceous and early Cenozoic stratigraphy and history of eastern Kyongsang Basin, S. Korea," *Journal of Geological Society of Korea* Vol. 26, pp. 471-487, 1990.
- [45] Y. Kamata, K. Hisada, Y. I. Lee, "Late Jurassic radiolarians from pebbles of Lower Cretaceous conglomerates of the Hayang Group, southeastern Korea," *Geosciences Journal* Vol. 4, pp. 165-174, 2000.
- [46] T. Mitsugi, K. Ishida, B. G. Woo, K. H. Chang, S. O. Park, H. Hirano, "Radiolarian-bearing conglomerate from the Hayang Group, the Kyongsang Supergroup, southeastern Korea," *Journal of Asian Earth Sciences* Vol. 19, pp. 751-763, 2001.
- [47] B. G. Woo, "Cretaceous stratigraphy and evolution of the central part of Kyongsang Basin," PhD thesis, Kyungpook National University, 256p, 1997.
- [48] S. Kojima, "Mesozoic terrane accretion in northeast China, Sikhote-Alin and Japan regions," *Palaeogeography, Palaeoclimatology, Palaeoecology* Vol. 69, pp. 213-232, 1989.
- [49] K. Ishida, T. Kozai, S. O. Park, T. Mitsugi, "Gravel bearing radiolaria as tracers for erosional events: a review of the status of recent research in SW Japan and Korea," *Journal of Asian Earth Sciences*, in press.