

# The Correlation of Plio-Pleistocene Strata in Japan Based on Volcanic Ash Keybeds

メタデータ	言語: eng 出版者: 公開日: 2017-10-05 キーワード (Ja): キーワード (En): 作成者: メールアドレス: 所属:
URL	<a href="http://hdl.handle.net/2297/6371">http://hdl.handle.net/2297/6371</a>

## The Correlation of Plio-Pleistocene Strata in Japan Based on Volcanic Ash Keybeds

Hiroshi OHKUBO

*Graduate School of Natural Sciences and Technology, Kanazawa University, Kanazawa, Ishikawa 920-1192, JAPAN*

**Abstract** - Volcanic ash keybed is a most precise indicator of contemporaneous datum planes of geological age. Recently, several Pliocene to Early Pleistocene wide spread volcanic ashes were reported from central Japan. Volcanic ash keybeds contribute chronologically to the study of paleoclimatic changes and tectonic events. While most of the correlations of the Pliocene-Pleistocene strata by volcanic ash keybeds are consistent with the correlations based on biostratigraphy, several of them are not consistent between the biostratigraphic bases and volcanic ash bases.

### I. Introduction

Volcanic ash beds are most precise contemporaneous datum planes of geological age. Many volcanic ash beds are intercalated in the Pliocene to Pleistocene strata in Japan. Volcanic ash beds are reported to be the useful marker bed for the chronology of the Quaternary climatic or tectonic events in or around Japan. For example, middle to late Pleistocene glacial-interglacial records were correlated between the Boso Peninsula and the Oiso hill [1]. High-resolution sequence stratigraphy of submarine fan system was reported using volcanic ash keybeds at the Kazusa Group [2]. Glacial eustatic sea level changes during the Pliocene to Early Pleistocene were reported to have 41k years periodicities, which were short compared with the Middle Pleistocene to Holocene sea level changes [3]. These facts indicate that chronological study of the Pliocene-Pleistocene sea level changes needs more precise datum planes. In the Middle Pleistocene to Holocene, many wide spread volcanic ashes have been reported by previous works. The Pliocene to early Pleistocene wide spread volcanic ashes were recently started to report by several studies [4,5]. For example, the Hotaka-Kd38 and the Ebisutoge-Fukuda Tephra were reported that were widely distributed in the central Japan [5]. Some of those volcanic ash keybeds were even traced to other distant regions [6].

The Pliocene to Pleistocene strata are widely distributed from the northeastern to the central Japan. The Pliocene-Pleistocene marine or non-marine strata in Japan were correlated with each other based on the biostratigraphy, magnetostratigraphy, tephrostratigraphy, and the radiometric age. Calcareous nannofossil biostratigraphy provides global and the most useful datum plane for stratigraphical age in marine sediments. However, calcareous nannofossil biostratigraphy is not enough precise to evaluate the correlation of the record of periodic glacial eustatic sea-level changes [7], and impossible to date non-marine strata. Volcanic ash keybeds are possible to dating of non-marine

strata with the help of palaeoclimate records. The author introduces examples of the contribution of volcanic ash keybeds to paleoclimatic study, and compare the correlation of the volcanic ash keybeds with biostratigraphy and/or radiometric age.




Fig. 1. The distribution of the Pliocene to the Early Pleistocene strata in the northeast-central Japan.

### II. Correlation of Strata the Pacific Ocean Coastal Areas

Localities of the representative of Pliocene-Pleistocene sequences of the Pacific coast in Japan are the Boso peninsula, the Kakegawa district, the Ise-Noubi Plain and the Osaka Plain. These localities have the strata that are called the Kazusa, the Kakegawa, the Tokai and the Osaka Groups, respectively. Many tephrostratigraphic studies were reported from these Groups based on the petrographical characteristics. The Kazusa Group is comprised of marine facies. In the Kazusa group, high-resolution sequence stratigraphic studies recognize eighteen depositional sequences, and these sequences are correlated to the oxygen isotope stages from deep-sea cores based on

fission-track age of the volcanic ash bed, magnetostratigraphy and calcareous nannofossil biostratigraphy [2]. The Osaka Group is mainly comprised of non-marine facies, and is intercalated twenty marine clay beds and many volcanic ash beds. Twenty marine clay beds are called the Ma-1, Ma0, Ma0.5, Ma1, Ma2... and Ma12 from the bottom to the top. These marine clay beds were traced using the volcanic ash keybeds under the Osaka plain, and correlated to oxygen isotope stages based on volcanic ash keybeds and magnetostratigraphic study [8]. The volcanic ash keybeds correlate the records of sea-level changes between the Kazusa and the Osaka groups. At the Early Pleistocene, the Azuki Volcanic Ash keybed in the Osaka Group was correlated to the Ku6c pyroclastic keybed. The Correlation of the Azuki and Ku6c volcanic ash beds indicates that the correlation of depositional sequence 7 of the Kazusa Group [2] and the Marine clay bed Ma3 of the Osaka Group [8].

### III. Correlation of Strata in the Japan Sea Coastal Areas

Localities of the representatives of the Japan Sea coastal Pliocene-Pleistocene sequence are the Oga Peninsula, the Niigata district and the Ishikawa-Toyama district. These localities have the strata called the Funakawa-Kitaura Formations, the Nishiyama-Haitsume Formations and the Himi Group, respectively.

Tephrostratigraphic studies based on petrographical characteristics were reported from the Niigata and Ishikawa-Toyama districts. In the Niigata district, the representative of the Pliocene-Pleistocene strata are the Nishiyama Formation, the Haitsume Formation and the Uonuma Group. Sequence stratigraphic study was reported from the Uonuma Group, and the Group was correlated to oxygen isotope stages based on tephrostratigraphy and the age of volcanic ash beds at the Uonuma Group [24]. In the Himi Group, sedimentary cycles were reported from the Omma and the Junicho Formations [25, 26]. Sedimentary cycles of the Junicho Formation were dated and they turned out to include the Plio-Pleistocene boundary. Then, they were correlated to the sedimentary cycles of the Uonuma Group based on volcanic ash keybed named the Ebisutoge-Fukuda Tephra [14]. The calcareous nannofossil biostratigraphy was studied for the Ishikawa-Toyama district [27,28]. In several sections of these districts, the correlation in the Himi Group suggested by tephrostratigraphic study is different from the correlation in the Himi Group based on calcareous nannofossil biostratigraphy [13,15]. The Ishikawa-Toyama district is geographically close to the Osaka Plain where non-marine sediments were widely distributed. This district is expected to have volcanic keybeds that can be correlated to volcanic ash beds of the Osaka Plain.




Fig. 2. Correlation of the Pliocene-Pleistocene strata in the Central Japan. Ages of the Kazusa Group are based on biostratigraphy [21], magnetostratigraphy [22] and radiometric age [23]. Ages of other strata are based on tephrostratigraphy and magnetostratigraphy.

## VI. Summary

Volcanic ash keybeds are not used as global datum planes. These keybeds are, however, precise indicators for chronological study of paleoclimatic or tectonic events. Volcanic ash keybeds are especially important marker-beds for non-marine strata. These keybeds are possible to be cross-checked and directly correlate sea-level change records between non-marine and marine sediments. In several sections, the geological age estimated by calcareous nannofossil biostratigraphy was reported to be different from those based on tephrostratigraphy. Volcanic ash keybeds are expected to provide important contemporaneous datum planes in the Pliocene-Pleistocene strata for the study of paleoceanographic change at the Japan Sea and the Pacific Ocean coasts.

## References

- [1] S. Sugihara, F. Arai, H. Machida, "Tephrochronology of the Middle to Late Pleistocene sediments in the northern part of the Boso Peninsula, central Japan," *the Journal of the Geological Society of Japan* Vol. 84, pp. 583-600, 1978 (in Japanese with English abstract).
- [2] M. Ito, "Volcanic ash layers facilitate high-resolution sequence stratigraphy at convergent plate margins: an example from the Plio-Pleistocene forearc basin fill in the Boso Peninsula, Japan", *Sedimentary Geology* Vol. 122, pp. 187-206, 1995.
- [3] N. J. Shackleton, "New data on the evolution of Pliocene climatic variability", In Vrba, E. S., Denton, D. H., Partridge, T. C. and Burckle, L. H., eds, *Palaecoclimate and evolution with emphasis on human origins*, Yale University Press, New Haven, pp. 242-248, 1997.
- [4] H. Kamata, T. Danhara, T. Yamashita, H. Hoshizumi, A. Hayashida and K. Takemura, "Correlation of the Azuki volcanic ash of the Osaka Group and the Ku6C volcanic ash of the Kazusa Group to the Imaichi pyroclastic-flow deposit in central Kyushu, Japan—A co-ignimbrite ash erupted from Shishimuta", *the Journal of the Geological Society of Japan* Vol. 100, pp. 583-600, 1978 (in Japanese with English abstract).
- [5] Y. Nagahashi, Y. Satoguchi and S. Yoshikawa, "Correlation and stratigraphic eruption age of the pyroclastic flow deposits and wide spread volcanic ashes intercalated in the Pliocene-Pleistocene strata, central Japan", *the Journal of the Geological Society of Japan* Vol. 106, pp. 51-69, 2000 (in Japanese with English abstract).
- [6] Y. Satoguchi, Y. Nagahashi, K. Kurokawa, S. Yoshikawa, "Tephrostratigraphy of the Pliocene to Pleistocene formations in central Honshu, Japan", Vol. 53, pp. 275-290, 1999 (in Japanese with English abstract).
- [7] A. Kitamura, K. Kimoto, T. Takayama, "Reconstruction of the thickness of the Tsushima Current in the Quaternary from molluscan fossils", *Palaeoecology, Palaeoecology, Palaeoecology* Vol. 135, pp. 51-69, 1997.
- [8] S. Yoshikawa and M. Mitamura, "Quaternary stratigraphy of the Osaka Plain, central Japan and its correlation with oxygen isotope record from deep sea cores", *the Journal of the Geological Society of Japan* Vol. 105, pp. 332-340, 1999 (in Japanese with English abstract).
- [9] K. Kikkawa, K. Mizuno, Y. Sugiyama, "Long distance correlation of the early to middle Pleistocene tephros from Kyushu to Kanto", *The earth Monthly* Vol. 13, pp. 223-234, 1991 (in Japanese).
- [10] Y. Tomita, K. Kurokawa, "Correlation between the Osaka Pink ash in the Osaka Group and the Ogi ash in the Niigata region", *Earth Science (Chikyu Kagaku)* Vol. 48, pp. 467-476, 1994 (in Japanese with English abstract).
- [11] S. Yoshikawa, F. Yoshida, E. Sugawa, "Volcanic ash layers of the Tokai Group and their correlation" *Earth Science (Chikyu Kagaku)* Vol. 45, pp. 453-467, 1991 (in Japanese with English abstract).
- [12] K. Kurokawa, H. Sakai, H. Hiwatari, T. Aoki, "Correlation of early Pleistocene volcanic ash layers between Niigata region and Boso Peninsula", *Abstracts of the 106th annual meeting of the Geological Society of Japan*, pp. 31, 1999 (in Japanese).
- [13] H. Okubo, "Tephrostratigraphy of late Pliocene to early Pleistocene in the Hokuriku region, and its petrographical characteristics and correlation with volcanic ash layers in other areas of Central Japan" Doctoral course *Thesis of Graduate School of natural science and Technology, Kanazawa University*, pp.123, 2000, (in Japanese with English abstract).
- [14] H. Okubo, K. Arai, "The Hotaka-Kd39 Tephra and Ebisutoge-Fukuda Tephra from the middle part of the Junicho Formation, Toyama prefecture, central Japan, and their stratigraphic position, *The Quaternary Research* Vol. 41, pp.429-442, 2002 (in Japanese with English abstract).
- [15] I. Tamura, H. Yamazaki, The widespread tephra layers of the Omma Formation distributed in Kanazawa area *Abstracts of 2002 Joint Meeting of Japan Earth Planetary Science*, 2002.
- [16] K. Kurokawa, " Tephrostratigraphy of the Nanatani to Unuma Formations of 13Ma to 1Ma in the Niigata region, central Japan", *Journal of the Japanese Association for Petroleum Technology* Vol. 64, pp. 80-93, 1999 (in Japanese with English abstract).
- [17] H. Okubo, "Plio-Pleistocene tephrostratigraphy of the Omma, Zukawa and Junicho Formations and uppermost part of the Takakubo Formation in the Hokuriku Region, Central Japan", *the Journal of the Geological Society of Japan* Vol. 105, pp. 836-851, 1999 (in Japanese with English abstract).
- [18] H. Okubo, T. Sato, M. Watanabe, "Stratigraphic correlation between the Plio-Pleistocene Yabuta and Junicho Formations using volcanic ash beds, and calcareous nannofossil biostratigraphy of lower part of the Junicho Formation in northwestern Toyama Prefecture, Central Japan", *the Journal of the Geological Society of Japan* Vol. 106, pp. 583-596, 2000 (in Japanese with English abstract).
- [19] Y. Tomita, K. Kurokawa, "A widespread volcanic ash layer of about 2.7Ma in central Japan; correlation of the Habutaki I (Osaka Group), the MT2 (Himi group) and the Arg-2 (Nishiyama Formation) ash layers" *the Journal of the Geological Society of Japan* Vol. 105, pp. 63-72, 1999 (in Japanese with English abstract).
- [20] T. Sato, K. Kameo, I. Mita, "Validity of the latest Cenozoic calcareous nannofossil datums and its application to the tephrochronology", *Earth Science (Chikyu Kagaku)* Vol. 53, pp. 265-274, 1999 (in Japanese with English abstract).
- [21] T. Sato, T. Takayama, M. Kato, T. Kudo, K. Kameo, "Calcareous microfossil biostratigraphy of the uppermost Cenozoic formations distributed in the coast of the Japan Sea", *Journal of the Japanese Association for Petroleum Technology* Vol. 53, pp. 13-29, 1988 (in Japanese with English abstract).
- [22] N. Niitsuma, "Magnetic stratigraphy in the Boso Peninsula", *the memoirs of the Geological Society of Japan* Vol. 82, pp. 163-181, 1976 (in Japanese).
- [23] M. Watanabe, T. Danhara, "Fission track ages of volcanic ash

layers of the Kazusa Group in the Boso Peninsula, central Japan”, Vol. 102, pp. 545-556, 1996 (in Japanese with English abstract).

[24] A. Urabe, T. Tateishi and O. Kazaoka, “Depositional cycle of marine beds and relative sea-level changes of the Plio-Pleistocene Uonuma Group, Niigata, Japan”, *the memoirs of the Geological Society of Japan* Vol. 45, pp. 140-153, 1995 (in Japanese with English abstract).

[25] A. Kitamura, Y. Kondo, H. Sakai, M. Horii, “Cyclic changes in lithofacies and molluscan content in the early Pleistocene Omma Formation, Central Japan related to the 41,000-year orbital obliquity”, *Palaeogeography, Palaeo-climatology, Palaeoecology* Vol. 112, pp. 345-361, 1994.

[26] K. Arai, H. Sakai, K. Konishi, “High-resolution rock-magnetic variability in shallow marine sediments: a sensitive paleoclimatic metronome”, *Sedimentary Geology* Vol. 110, pp. 7-23, 1997.

[27] T. Takayama, M. Kato, T. Kudo, T. Sato and K. Kameo, “Calcareous microfossil biostratigraphy of the uppermost Cenozoic Formations distributed in the coast of the Japan Sea”, *Journal of the Japanese Association for Petroleum Technology* Vol. 53, pp. 9-27, 1988 (in Japanese with English abstract).

[28] C. Shida, K. Kameo and T. Takayama, “Lateral and vertical distributions of calcareous nannofossils in the Omma Formation”, *professor H. Matsuo Memorial Volume*, pp. 111-124, 1991 (in Japanese with English abstract).