

1 **Factors associated with tuberculosis cases in Semarang District Indonesia; case-control**  
2 **study performed in the area where Case Detection Rate was extremely low**

3

4 Sri Ratna Rahayu<sup>1</sup>, Hironobu Katsuyama<sup>2</sup>, Masashi Demura<sup>1</sup>, Midori Katsuyama<sup>1</sup>, Yoko Ota<sup>1</sup>,  
5 Hideji Tanii<sup>1</sup>, Tomomi Higashi<sup>1</sup>, Ngakan Putu Djaja Semadi<sup>3</sup> and Kiyofumi Saijoh<sup>1</sup>.

6 <sup>1</sup>Kanazawa University School of Medicine, Kanazawa 9208640 Japan,

7 <sup>2</sup>Department of Public Health, Kawasaki Medical University, Kurashiki 7011092, Japan

8 <sup>3</sup>Health Department Semarang District, MT Haryono 29 Ungaran, Semarang District,  
9 Indonesia.

10

11 Correspondence:

12 Kiyofumi Saijoh, MD, PhD

13 Professor and chairperson

14 Department of Hygiene

15 Kanazawa University School of Medicine

16 13-1 Takaramachi,

17 Kanazawa 9208640, Japan

18 Email: [saijohk@med.kanazawa-u.ac.jp](mailto:saijohk@med.kanazawa-u.ac.jp)

19 **Keywords**

20 Tuberculosis, Case Detection Rate, Risk factors, Indonesia, DOTS.

21 **Abstract**

22 *Objectives* Indonesia is ranked to be the 4th as a contributor to the Tuberculosis (TB) in the  
23 world. Semarang District in Central Java displays extremely low Case Detection Rate (CDR),  
24 possibly contributing to the local prevalence of TB.

25 *Methods* Case-control study was performed to explore the factors cause of such low CDR.  
26 We recruited 129 TB cases and 83 controls that visited the same centers and were not  
27 diagnosed as TB.

28 *Results* The cases had  $7.5 \pm 2.3$  symptoms/person in average, indicating the delay of  
29 diagnosis because the controls only displayed  $1.0 \pm 1.7$ . The multiple logistic regression  
30 analysis comparing the cases/controls extracted following factors as a risk to have TB; farmer,  
31 close contact with TB patients, ignorance of whether Bacillus Calmette-Guérin (BCG) was  
32 accepted or no, smoking, low income, a lot of people living in the same room, irregular hand  
33 wash before meals, not wash hands after blow, soil floor, and no sunlight and no ventilation in  
34 the house.

35 *Conclusions* Neither the cases nor the controls always knew the symptoms and how to avoid  
36 TB infection probably causes the delay of diagnosis. It is difficult to change the current living  
37 condition. Thus, the amendment of the community based education program of TB seems to

38 be required.

39 **Abbreviations**

40 BCG : Bacillus calmette-guérin

41 CDR : Case detection rate

42 CI : Confidence Interval

43 DOTS : Directly observed treatment short-course

44 HIV : Human immunodeficiency virus

45 MA : Monovariate analysis

46 MDR : Multidrug resistant

47 MLR : Multiple logistic regression analysis

48 TB : Tuberculosis

## 49 **1. Introduction**

50 Tuberculosis (TB) is one of the biggest problems throughout the world and a leading cause of  
51 death and major public health problem [1]. Many people with TB remain poor in diagnosis or  
52 are diagnosed only after long delays. The high burden of undiagnosed TB causes much  
53 suffering economic hardship and sustained transmission [2-4].

54 According to Global Tuberculosis Report 2012 [5], Indonesia is ranked to the 4th  
55 (0.4 million-0.5 million) as a contributor to the TB world after India (2.0 million-2.5 million),  
56 China (0.9 million-1.1 million), and South Africa (0.4 million-0.6 million). Since 2000, 22  
57 high burden countries account for 82 % of all estimated incident cases [5].

58 The discovery of TB in Indonesia in 2011 is 807 per 100,000 populations and that in  
59 Central Java is 637 per 100,000 populations [6]. The TB Case Detection Rate (CDR) of  
60 Central Java Province is estimated to be 56.93 % [7], but among its local districts, Semarang  
61 District displays extremely low CDR being 19.21 % [8]. National targets for TB control  
62 strategy Indonesia, the strategic plan of the Ministry of Health from 2010-2014, is to reduce  
63 the TB prevalence to 224 per 100,000 populations. Output targets are (1) to increase the CDR  
64 of new cases of smear-positive pulmonary TB from 73 % to 90 %, (2) to reach the percentage  
65 of successful treatment of new cases of smear positive pulmonary TB to 88 %, (3) to increase

66 “the percentage of province with CDR above 70 %” to reach 50 %, and (4) to increase “the  
67 percentage of the province with treatment success rate being above 85%”, from 80 % to 88 %  
68 [9].

69 To achieve the national targets controlling TB, some strategies seem to be very  
70 important; empowerment in the community; early detection and registration of TB patients;  
71 improvement of the quality Directly Observed Treatment short-course (DOTS) services;  
72 facing the challenges of TB/HIV, Multidrug Resistant (MDR-TB) and childhood TB; to meet  
73 the demands of society poor and other vulnerable groups [9]. Domain knowledge is very  
74 important in the formation of action. In several developing countries, TB patients are  
75 perceived to seek late care or avoid care, due to misunderstanding to popular TB etiologies  
76 like sharing utensils, heavy labor, smoking, bewitchment and hereditary transmission  
77 [10-12]. Thus, it seems very important to know the basic knowledge, attitude, and behavior of  
78 the people in such a low CDR region to complete the TB programs.

79 In the present study, differences in the basic knowledge, attitude, and behavior were  
80 compared among the TB cases registered in 2012 at the health centers in Semarang District  
81 and the controls that visited the same health centers and diagnosed differently.

82

83 **2. Methods**

84 This study was designed to be Case-Control study [13] during 2012, January until 2013,  
85 October. The TB cases (129) were recruited among the cases registered with TB diagnosis at  
86 the health centers in Semarang District, and 50 % were female. The controls (83) were  
87 recruited among those who visited the same health centers and were diagnosed differently,  
88 and 52 % were female. Average age of the cases and controls (Mean  $\pm$  SD) was  $41.2 \pm 15.3$   
89 and  $35.7 \pm 11.7$ , respectively. This research was approved by the ethical committees of  
90 Kanazawa University School of Medicine, Japan and Diponegoro University School of  
91 Medicine, Dr Kariadi Hospital Semarang, Indonesia and Semarang State University,  
92 Indonesia. All participants approved this research by written informed consent.

93 *TB information of the cases*

94 TB cases were sent to the Health Center from clinics and diagnosed to have TB using  
95 physical examination, microscopic examination by Ziehl Neelsen staining, X-ray, etc.  
96 Way of diagnosis, symptoms they had, and way of treatment were examined. All of them  
97 were treated with DOTS properly regardless of having side effects or not.

98 *Characteristics of the cases and controls*

99 Demographic characteristics like gender, age, occupation, origin, BCG experience, close

100 contact with TB patients, smoking habit, having pets, and income was examined. Source of  
101 TB information (multiple answers) was also examined. Thereafter, differences in living  
102 condition and attitudes in daily life were examined.

### 103 *Knowledge and opinion about TB*

104 Symptoms of TB (multiple answers) and the way to avoid TB (multiple answers) that the  
105 participants thought were questioned. Differences in the opinion and attitude against TB  
106 between the cases and controls were examined.

### 107 *Statistical analysis*

108 The difference in the age and numbers of complaint of the cases and controls was estimated  
109 using Student's t-test. The differences in the frequency of answers between the cases and the  
110 controls were estimated by Chi-square ( $\chi^2$ ) test (monovariate analysis, MA). Multiple logistic  
111 regression analysis (MLR) with the cases/the controls as the dependent valuable was utilized  
112 with the independent variables using groups classified by characteristics of the cases and  
113 controls. All analyses were performed with SPSS ver. 19 (SAS Institute Inc., Cary, NY). In  
114 all analyses,  $\rho < 0.05$  was taken to indicate statistical significance.

115

## 116 **3. Results**



117 The 129 cases included 65 male and 64 female, respectively, and the average age was  $41.2 \pm$   
118  $15.3$  (Table 1). Although all the cases were registered at the health centers, around 20 % were  
119 diagnosed at the different medical facilities (Table 2). Sputum smear was the first choice for  
120 diagnosis [1, 14], but 20 cases were diagnosed without any clinical examination. Chest X-ray  
121 was utilized for the diagnosis of more than half of the cases. Long lasting cough with sputum  
122 was the most common symptom that was followed by chest pain, malaise, anorexia, and  
123 weight loss. Around two third displayed hemoptysis, dyspnea, sweat at night, and long lasting  
124 sub fever at night. The cases had many symptoms being  $7.5 \pm 2.3$  complaints/person in  
125 average. These cases were registered, hence all of them underwent to treatment, whether they  
126 knew it was under DOTS or not (Table 3). Nine had no supervisors and at least one fourth had  
127 to pay treatment fee, suggesting that they were not under DOTS. More than 85 % of the cases  
128 quit taking medicine at 6 months, regardless of frequency of medication in first 2 months.  
129 Treatment with 4 drugs was the most common, but that with 2 drugs was also observed in  
130 around 15 % of the cases. Sputum smear was the most common examinations during  
131 treatment followed by chest X-ray.

132 The controls were selected from people visited the same health center and  
133 diagnosed not having TB. We tried to obtain age-gender-matched control. We could achieve

134 gender-matched, but their age was slightly younger than the cases (Table 1). Around 80 % of  
135 the cases were farmers and around 30 % of them graduated from elementary school alone.  
136 More than half of the controls received BCG whereas more than half of the cases did not  
137 know whether they received BCG or not. Nine percent of the cases experienced close contact  
138 with TB patients but none of the controls did. Income of the cases was significantly lower  
139 than that of the controls, and three fourth of them got less than 100 \$/month. The cases were  
140 diagnosed already, hence they got the TB information from medical staff more than the  
141 controls, but they usually did not use other sources (Table 1). When the living conditions  
142 were compared, all conditions were significantly different between the cases and the controls  
143 (Table 4). “Ceramic floor”, “outside kitchen”, “gas for cooking”, “open windows everyday”,  
144 “sunlight into the house”, and “ventilation in every room” were more common in the controls,  
145 whereas “window in each room” and high “humidity in the house” were in the cases.

146           The cases less frequently “washed their hands before eating” but more frequently  
147 “shared the dishes with others” and “drunk from the same glasses/bottles” than the controls  
148 (Table 5). They also less frequently “washed their hands after blowing” than the controls,  
149 whereas no difference was observed in the frequency of whether “they worked when they felt  
150 unwell” between these two groups.

151           In order to clarify what kinds of these physical factors were most affected  
152 difference in the cases and the controls, the MLR was applied (Table 6). The cases/controls  
153 were the dependent variables and the groups divided by above mentioned information were  
154 used as determinants. The way of obtaining TB information was removed from the  
155 determinant because that from the cases was modified as described above. The obtained risks  
156 were “farmers”, ”close contact with TB patients”, “whether or not they did not know they  
157 received BCG”, “smoking”, and “low income; < 100 \$/month”. “High income; > 150 \$/month”  
158 was also extracted as a risk compared with “middle income; 100-150 \$/month”. To “wash  
159 hands before eating”, “wash hands after blow” and “not work when unwell” were protective.  
160 Among living conditions, “live with  $\geq 3$  person in the same room”, “soil floor”, no “sunlight  
161 in the house” and no “ventilation in the house” were extracted as risk.

162           Regardless of many symptoms, the cases did not always display significant  
163 differences with the controls among the opinion regarding what they thought was TB  
164 symptoms (Figure 1). “Long lasting sub fever” alone was significantly higher in the controls.

165           More than 70 % of both the cases and the controls thought that it was important to  
166 “cover mouth/nose when someone sneezed” (Figure 2). More than half of the controls thought  
167 that “avoid sharing dish”, “avoid drinking from the same glass/bottle”, “wash hands after

168 touching items in the public” and “maintain good nutrition” were the way to avoid getting TB  
169 and the rates were significantly higher than the cases. In fact, 76 % of the controls washed  
170 hands before eating and only 35 % of the cases did so (Table 5). Moreover, 76 % (9 + 67) of  
171 the cases at least sometimes “shared the dish” and” drunk from the same glass”, that was  
172 significantly higher than the controls. On the other hand, 87 % (39 + 48) of the controls at  
173 least sometimes “washed hands after blowing”. “Vaccination” was also higher in the controls  
174 than the cases (Figure 2).

175 Opinions related to the seriousness and shame, did not display any significant  
176 difference between the cases and the controls (Table 7). Although many of the cases and  
177 controls thought TB to be “serious”, they did not always think that TB was “serious at  
178 workplaces” and “affected work performance”. Significant difference was not observed in  
179 “be ashamed of having TB” but the cases were tended to want to “hide having TB”.  
180 Significantly more controls thought “TB affected relationship with others” and “wanted to  
181 live isolated”, whereas there was no significant difference in “TB affected family  
182 responsibility” against the controls. Both of the cases and controls usually tried to be good at  
183 TB patients. Around 50 % of the controls believed “TB treatment was very costly” but  
184 around one fourth of the cases thought so. “HIV positive people should concern about TB”

185 was significantly higher in the controls than in the cases. Around one fourth to one third of  
186 the cases and controls believed that TB was hereditary.

#### 187 **4. Discussion**

188 In Indonesia, regular health examination was not mandatory [6]. Therefore, after symptoms  
189 became apparent, the person visited the clinic where sputum smear was not always available.  
190 The cases in the present study displayed  $7.5 \pm 2.3$  complaints/person in average, indicating  
191 the delay of diagnosis [15]. Although early diagnosis and initiation of treatment of infectious  
192 cases is the best measure to reduce transmission [3, 16, 17], in some countries, 20 % of  
193 patients were not diagnosed for over 6 months from the onset of symptoms [18]. Even after  
194 the symptoms became obvious, for diagnosis, it took at least 2 more days because positive TB  
195 was defined as more than 2 positive sputum smears in the smear performed three times within  
196 2 days [14]. Household contacts continued meantime, when patients were with potentially  
197 infectious forms proceeding to high prevalence of TB [19, 20]. On the other hand, culture was  
198 not common, whereas patients with smear-negative, culture-positive TB were reportedly  
199 responsible for TB transmission [21, 22]. Immediate introduction of culture examination is  
200 required because, in addition to high sensitivity, it allows determining whether the patient is  
201 sensitive to anti-TB drugs and useful for finding extra pulmonary TB [14]

202 The cases did not always know whether they were under DOTS treatment or not, but all the  
203 cases could luckily quit taking medicine regardless of the obvious delay of diagnosis. Around  
204 80 % of the cases were farmer, and around 30 % graduated elementary school alone, hence  
205 their income was lower than the controls. Low income and low education are reportedly  
206 associated with TB infection [4, 23-25]. They also had lost the chance of BCG injection. This  
207 occupation was also extracted as a risk by the MLR. However, the MLR extracted high  
208 income as a risk as well. Such a result is not always in accordance with several studies [4,  
209 23-25], whereas it is conceivable that people with high income, regardless of their occupation,  
210 had more chances to live and/or work at the places with a lot of people where a risk of TB  
211 transmission was supposed to be high. The number was small but only cases had a chance to  
212 close contact with TB patients. TB contact was absolutely the risk of TB transmission [18, 26].  
213 No significant difference in the rate of “smoking” by the MA, but the MLR extracted  
214 “smoking” as a risk to be the case. This is in good accordance with the previous report [23]. It  
215 is natural that the cases “source of TB information” was medical staff, but the cases were not  
216 always eager to collect information from other sources comparing with the controls. All the  
217 items related living condition was significantly different between the cases and the controls  
218 by the MA. Among them, the MRL extracted “small number of the person in the room”,

219 “ceramic floor”, “sunlight in the house”, and “ventilation in the house” as protective.  
220 Importance of good ventilation was emphasized elsewhere [3, 23]. Excluding “work when  
221 unwell”, their attitudes displayed significant differences between the cases and the controls by  
222 the MA. “Share the dish” and “drink from the same glasses/bottles” were not extracted by the  
223 MLR. Instead, “work when unwell” was extracted as well as “wash hands before eating” and  
224 “wash hands after blow”. These findings may be a reflection that TB is airborne. In general,  
225 the cases were not aware of danger in their attitudes, which was in good accordance with  
226 previous reports [10-12].

227 Both of the cases and the controls did not recognized “dyspnea” and “chest pain” as TB  
228 symptoms. Significant differences existed, but “long lasting sub fever” was also not  
229 considered as TB symptoms. “dyspnea”, “chest pain” and “long lasting sub fever” were less  
230 frequent than “cough with sputum”, “malaise” and so on but number of the cases complaint  
231 them. Thus, it seems necessary to let the people know the TB symptoms [10-12].

232 TB itself was recognized to be dangerous both by the cases and the controls, but they did not  
233 recognize its dangerousness at work places. Many of the cases were farmers; hence it seems  
234 less possible to spread TB than workers. However, TB positive workers can work and be able  
235 to transmit TB to their colleagues. Comparing with the rate of both the cases and the controls

236 who thought “having TB was a shame”, that of “wanted to hide having TB” was less. It seems  
237 natural that more controls who did not receive TB treatment believed that TB was “affected  
238 relationship with others” and wanted to “live isolated in case of TB” than the cases. DOTS  
239 performed under the governmental hospitals and health centers were free [9], but some cases  
240 visiting private hospitals/clinics had to pay the treatment fee. Higher rate of “HIV positive  
241 people should be concerned about TB” in the control was reflection that they were more eager  
242 to collect information than the cases. HIV infection reportedly affected TB infection [27].  
243 However, the number who believed “TB was hereditary” was not different between two  
244 groups.

245 Some aspects underlying the low coverage CDR are problems of socioeconomic,  
246 education/knowledge and stigma [4]. Economic conditions will affect the public in getting not  
247 only good environmental home conditions but also the excellent level of education. The level  
248 of education in this study was relatively low because many people only finished elementary  
249 school where sufficient TB education was impossible. Poor education will cause shortage of  
250 knowledge about TB, leading the public into embarrassment and sometimes attitude to hide  
251 their disease if they exposed to TB. Such conditions may cause the delay for some people to  
252 go to the health service [18]. As a result, TB was spread among the farmers even when their



253 contact was not always intense like workers. It is very difficult to change occupation, income  
254 and housing condition, immediately. Thus, community based TB education is very important.  
255 It may be useful to educate and expose not only public but also private practitioners to the  
256 community based TB program [9, 28].

257 Some cases were not dependent on the free DOTS program. To inform the existence of this  
258 program is also a good education. Utilization of this program not only reduces multi-drug  
259 resistant TB, but also helps reducing out-of-pocket expenses to patients [27]. Number of  
260 syndromes of the cases absolutely indicated the delay of diagnosis.

261 The classic symptoms of TB are fever, cough and weight loss, but they are non-specific and  
262 can be mimicked by other conditions, including malignancy and other pulmonary infections.  
263 That is, in an early stage, such syndromes are not always specific to TB. However,  
264 importance of these classic lung related syndromes should be aware that they are possible  
265 signs of initiation of TB expansion [29].

266

## 267 **Acknowledgements**

268 We would like to thank to Semarang State University, Directorate General of Higher  
269 Education (DGHE or DIKTI) and Health Department Semarang District, for all support.

270 **Conflict of Interests**

271 The authors declare that there is no conflict of interests regarding the publication of this  
272 paper.

273 **References**


- 274 1. WHO. Systematic Screening for active TB. WHO: World Health Organization, 20  
275 Avenue Appia, 1211 Geneva 27 Switzerland, 2013.
- 276 2. Bauer, M., A. Leavens, and K. Schwartzman, A systematic review and meta-analysis  
277 of the impact of tuberculosis on health-related quality of life. *Qual Life Res.* 2013; 22:  
278 2213-35.
- 279 3. Saxena, S., V. Karkhanis, and J.M. Joshi, Tuberculosis prevention: an enigma worth  
280 unravelling. *Indian J Tuberc.* 2012; 59: 65-7.
- 281 4. Wu, J. and K. Dalal, Tuberculosis in Asia and the pacific: the role of socioeconomic  
282 status and health system development. *Int J Prev Med.* 2012; 3: 8-16.
- 283 5. WHO. Global Tuberculosis Report 2012. WHO, 20 Avenue Appia, 1211–Geneva–27,  
284 Switzerland, 2012:11.
- 285 6. Ministry of Health , R.I. Current Report of Development of Tuberculosis in Indonesia  
286 January-December 2012. Ministry of Health Republic Indonesia: Jakarta, Indonesia,  
287 2012.
- 288 7. Ministry of Health , R.o.I. Profile of Health Data of Indonesia in 2011. Ministry of  
289 Health of The Republic of Indonesia: Jakarta, Indonesia. 2012,
- 290 8. Health Department , S.D. Reported TB Cases in Semarang District, 2011. Semarang  
291 District, Central Java, Indonesia.2011.
- 292 9. Ministry of Health, R.O.I, Directorate General of Disease Control and Environmental  
293 Health National. Strategy for Controlling TB in Indonesia 2010-2014. Republic Of  
294 Indonesia Health Ministry Directorate General of Disease Control and Environmental  
295 Health , Jakarta, 2011.
- 296 10. Notoatmodjo, S. Health promotion and behavioral sciences. Jakarta: Rineka Cipta.  
297 2007.
- 298 11. Tasnim, S., A. Rahman, and F.M. Hoque, Patient's Knowledge and Attitude towards  
299 Tuberculosis in an Urban Setting. *Pulm Med.* 2012; 2012: 352850.
- 300 12. Buregyeya, E., et al., Tuberculosis knowledge, attitudes and health-seeking behaviour  
301 in rural Uganda. *Int J Tuberc Lung Dis.* 2011; 15: 938-42.
- 302 13. Raj S. Bhopal, A.B.a.J.U.P.o.P.H., et al., Concepts of Epidemiology, an itegrated  
303 introduction to the ideas, theories, principles and methods of epidemiology. 2002,  
304 United States,New York: Oxford University.
- 305 14. Health Minister of The Republic of Indonesia, N.M.S.V. Tuberculosis Control

- 306 Guidelines. Health Minister of The Republic of Indonesia Jakarta, Indonesia, 2009.
- 307 15. Ahmad, R.A., et al., Diagnostic delay amongst tuberculosis patients in Jogjakarta  
308 Province, Indonesia is related to the quality of services in DOTS facilities. *Trop Med*  
309 *Int Health*. 2011; 16: 412-23.
- 310 16. Dye, C., Tuberculosis 2000-2010: control, but not elimination. *Int J Tuberc Lung Dis*.  
311 2000; 4: S146-52.
- 312 17. Reid, M.J. and N.S. Shah, Approaches to tuberculosis screening and diagnosis in  
313 people with HIV in resource-limited settings. *Lancet Infect Dis*. 2009; 9: 173-84.
- 314 18. Yang, Y.R., et al., Evaluation of the tuberculosis programme in Ningxia Hui  
315 Autonomous region, the People's Republic of China: a retrospective case study. *BMC*  
316 *Public Health*. 2012; 12: 1110.
- 317 19. Batra, S., et al., Childhood tuberculosis in household contacts of newly diagnosed TB  
318 patients. *PLoS One*. 2012; 7: e40880.
- 319 20. Jensen, P.A., Centers for Disease Control and Prevention (U.S.), and National Center  
320 for HIV STD and TB Prevention (U.S.), Guidelines for preventing the transmission of  
321 *Mycobacterium tuberculosis* in health-care settings, 2005. MMWR recommendations  
322 and reports. 2005, Atlanta, GA: U.S. Dept. of Health and Human Services, Public  
323 Health Service, Centers for Disease Control and Prevention. 141 p.
- 324 21. Kanaya, A.M., D.V. Glidden, and H.F. Chambers, Identifying pulmonary tuberculosis  
325 in patients with negative sputum smear results. *Chest*. 2001; 120: 349-55.
- 326 22. Sarmiento, O.L., et al., Assessment by meta-analysis of PCR for diagnosis of  
327 smear-negative pulmonary tuberculosis. *J Clin Microbiol*. 2003; 41: 3233-40.
- 328 23. Oxlade, O. and M. Murray, Tuberculosis and poverty: why are the poor at greater risk  
329 in India? *PLoS One*. 2012; 7: e47533.
- 330 24. Shen, X., et al., Tuberculosis in an urban area in China: differences between urban  
331 migrants and local residents. *PLoS One*. 2012; 7: e51133.
- 332 25. Clark, M., P. Riben, and E. Nowgesic, The association of housing density, isolation  
333 and tuberculosis in Canadian First Nations communities. *Int J Epidemiol*. 2002; 31:  
334 940-5.
- 335 26. Sacchi, F.P.C., et al., Sugar cane manufacturing is associated with tuberculosis in an  
336 indigenous population in Brazil. *Transactions of the Royal Society of Tropical*  
337 *Medicine and Hygiene*. 2013; 107: 152-157.
- 338 27. Ismail, I. and A. Bulgiba, Determinants of unsuccessful tuberculosis treatment  
339 outcomes in Malaysian HIV-infected patients. *Prev Med*. 2013; 57: S27-30.


- 340 28. Artawan Eka Putra, I.W., et al., Factors associated to referral of tuberculosis suspects  
341 by private practitioners to community health centres in Bali Province, Indonesia. BMC  
342 Health Serv Res. 2013; 13: 445.
- 343 29. Davies, P.D.O., P.F. Barnes, and S.B. Gordon, Clinical tuberculosis. 4th ed. 2008,  
344 London: Hodder Arnold
- 345
- 346



**Figure 1. TB symptoms**



**Figure 2. The way to avoid getting TB**





## Figure legends

Fig.1

Differences in the knowledge about TB symptoms between the cases and controls (multiple answer). \*Significant differences ( $p < 0.05$ ,  $\chi^2$  test and Student's t-test).

Fig. 2

Differences in the way that the cases and controls thought it good to avoid getting TB

\*Significant differences ( $p < 0.05$ ,  $\chi^2$  test).

**Table 1. Demographic characteristics of the cases and control**

	Cases		Controls			Cases		Controls	
	N	%	N	%		N	%	N	%
Total	129		83		Smoking				
Gender					No	41	32	34	41
Male	65	50	40	48	No, but family smokes	45	35	28	34
Female	64	50	43	52	Yes	44	34	21	25
Occupation*					Pet				
Farmer	101	78	31	37	Yes	67	52	34	41
Others	28	22	52	63	No	62	48	49	59
Education*					Income*				
Elementary school	38	29	8	10	<100	96	74	38	46
Jr high school	59	46	32	39	100-150	19	15	31	37
High school or higher	32	25	43	52	>150	14	11	14	17
Origin					Source of TB information (multiple answer)				
Rural	106	82	70	84	Broadcast <sup>#</sup>	74	57	64	77
Urban	23	18	13	16	Billboards <sup>#</sup>	10	8	20	24
Have you ever had BCG?*					Newspaper <sup>#</sup>	18	14	23	28
Yes	33	26	46	55	Medical <sup>#</sup>	89	69	37	45
No	30	23	21	25	Acquaintance <sup>#</sup>	9	7	24	29
Don't know	66	51	16	19	Teacher <sup>#</sup>	6	5	25	30
Close contact with TB patients*									
Yes	18	9	0	0					
No	111	86	83	100					

Average age of the case and control (Mean ± SD) was 41.2 ± 15.3 and 35.7 ± 11.7, respectively, and the difference was significant ( $p < 0.05$ , Students' t-test).

\*Significant difference in the rate of answers between the case and the control ( $p < 0.05$ ,  $\chi^2$  test).

<sup>#</sup>For multiple answer questions, the difference in the rate of each source was examined using  $\chi^2$  test. The case and control displayed significant differences in all sources ( $p < 0.05$ ).

**Table 2. The way of diagnosis and symptoms**

---

N%
Location of diagnosis
Health center
Private clinic
Others
Way of diagnosis
No clinical examination
Smear alone
Smear, X-ray
Symptoms (multiple answer)
Cough with sputum
Cough >2w
Haemoptysis
Dyspnea
Chest pain
Malaise
Anorexia
Weight loss
Sweat at night
Subfever >1 Mo

---

Mantoux test was utilized only in 2 cases. The mean  $\pm$  SD of complaints/person was  $7.5 \pm 2.3$ , whereas that of the controls only have  $1.0 \pm 1.7$  ( $p < 0.05$ , Students' t-test.)

**Table3. Information related to treatments of the cases**

	N	%
Was treatment made under DOTS?		
Yes	80	62
No	14	11
Don't know	35	27
Who was a supervisor ? <sup>a</sup>		
Family	86	67
Medical Service	67	52
None	9	7
Was treatment free of charge?		
Yes	80	62
No	33	26
Don't know	16	12
How long was the duration of treatment?		
<6 Mo	7	5
6Mo	112	87
>6Mo	10	8
How frequent do you take medication within first 2 Mo?		
Everyday	80	62
1x/week	35	27
Longer	14	11
How many kinds of drugs did you take?		
2	19	15
3	34	26
4	70	54
5	6	5
Were you suffered from side effects?		
+	70	54
Which examinations were utilized during treatment ? <sup>b</sup>		
Sputum smear	104	81
Chest X-ray	79	61
Sputum culture	11	8.5
Mantoux test	11	8.5
Distance to the health center		
≤5 min	85	66
≤30 min	21	16
Longer	26	18

<sup>a</sup>35 cases were supervised both by family and medical staff.

<sup>b</sup>76 cases were subjected more than 2 examinations.

**Table 4. Comparison in living condition of the participants**

	Cases		Controls	
	N	%	N	%
Number of person in the same room*				
1	15	12	10	12
2	84	65	72	87
>3	30	23	1	1
House floor*				
Soil	35	27	2	2
Plaster	49	38	18	22
Ceramics	45	35	63	76
Location of kitchen*				
Outside	51	40	60	72
Inside	78	60	23	28
Fuel for cooking*				
Cordwood	52	40	8	10
Gas	77	60	75	90
Window in each room*				
Yes	52	40	8	10
No	77	60	75	90
Open windows every day*				
Yes	43	33	60	72
No	86	67	23	28
Sunlight into the house*				
Yes	66	51	76	92
No	63	49	7	8
Ventilation in every room*				
Yes	50	39	75	90
No	79	61	8	10
Humidity in the house*				
Humid	84	65	17	20
Not humid	45	35	66	80

\*Significant difference between the cases and the controls ( $p < 0.05$ ,  $\chi^2$  test).

**Table 5. The differences in attitudes that the participants usually do**

	Cases		Control	
	N	%	N	%
Do you wash your hands before eating?*				
Yes	45	35	63	76
Sometimes	72	56	18	22
No	12	9	2	2
Do you eat from the same dish with others?*				
Yes	11	9	8	10
Sometimes	86	67	28	33
No	32	25	47	57
Do you drink from the same glasses/bottles with others?*				
Yes	7	5	9	11
Sometimes	92	71	35	42
No	30	23	39	47
Do you wash your hands after blowing?*				
Yes	25	19	40	48
Sometimes	69	53	32	39
No	35	27	11	13
Do you work when you are unwell?				
Yes	23	18	25	30
Sometimes	78	60	40	48
No	28	22	18	22

\*Significant difference between the cases and the controls ( $p < 0.05$ ,  $\chi^2$  test).

**Table 6. Multiple logistic regression analysis using case/control as the dependent valuable and living status as determinants**

	Comparison	Reference	Odds	P	(95 % CI)	
Occupation						
	Workers	Farmers	0.05	<0.01	0.00	0.20
	Others	Farmers	0.05	0.01	0.00	0.54
Close contact with TB patients						
	No	Yes	0.00	0.02	0.00	0.49
BCG						
	Yes	Don't know	0.04	<0.01	0.00	0.32
Smoking						
	No	Yes	0.14	0.04	0.01	0.92
Income (\$/Mo)						
	100-150	<100	0.05	<0.01	0.00	0.33
	100-150	>150	0.08	0.04	0.01	0.89
Person/room						
	1	≥3	0.00	<0.01	0.00	0.35
	2	≥3	0.00	<0.01	0.00	0.27
Wash hands before eating						
	Yes	Sometimes	0.06	<0.01	0.01	0.32
Wash hands after blow						
	Sometimes	No	0.06	0.01	0.01	0.46
Work when unwell						
	No	Yes	0.09	0.02	0.00	0.66
Floor						
	Ceramics	Soil	0.06	0.04	0.00	0.90
Sunlight in the house						
	Yes	No	0.06	0.02	0.00	0.67
Ventilation in the house						
	Yes	No	0.02	<0.01	0.00	0.24

Source of TB information (multiple answer) was excluded because this information was modified by consultation of the cases to the health center. Among determinants, age class, gender, education, pet, share the dishes, drink from the same bottle/glass, location of kitchen, fuel for cooking, open the window everyday, and humidity in the house were not selected.

**Table 7. Differences in the opinion against TB between the cases and controls**

	Cases		Controls	
	N	%	N	%
Do you think that TB is serious?				
Yes	91	71	67	81
No	2	2	3	4
Don't know	36	28	15	18
Do you think that TB is serious at work places?				
Yes	46	36	34	41
No	14	11	10	12
Don't know	69	53	39	47
Does TB affect your work performance?				
Yes	51	40	43	52
Not always	60	47	29	35
No	18	14	11	13
Do you be ashamed of having TB?				
Yes	49	38	22	27
No	53	41	35	42
Don't know	27	21	26	31
Do you want to hide having TB?*				
Yes	14	11	10	12
Not always	74	57	30	36
No	41	32	43	52
Does TB affect relationship with others?*				
Yes	20	16	36	44
Not always	72	56	35	42
No	37	29	12	14
Does TB affect family responsibilities?				
Yes	58	45	41	49
Not always	53	41	35	42
No	18	14	7	8
Will you want to live isolated due to having TB?*				
Yes	5	4	12	14
Not always	43	33	25	30
No	81	63	46	55
How do you feel about the person with TB?				
Desire to help	94	73	59	71
Want to stay away	28	22	20	24
No particular feeling	7	5	4	5
Is TB treatment very costly?*				



Yes	32	25	42	51
Not always	44	34	21	25
No	53	41	20	24
Do you think that HIV positive people should concern about TB?*				
Yes	23	18	39	47
Not always	79	61	30	36
No	27	21	14	17
Do you think that TB is hereditary				
Yes	30	23	24	29
No	79	61	47	57
Don't know	20	16	12	14

---

\*Significant difference between the cases and the controls ( $p < 0.05$ ,  $\chi^2$  test).