

# Ostracodes reveal the sea-bed origin of tsunami deposits

メタデータ	言語: eng 出版者: 公開日: 2021-04-15 キーワード (Ja): キーワード (En): 作成者: メールアドレス: 所属:
URL	<a href="https://doi.org/10.24517/00061630">https://doi.org/10.24517/00061630</a>

This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 International License.



## Ostracodes reveal the sea-bed origin of tsunami deposits

Gengo Tanaka,<sup>1</sup> Hajime Naruse,<sup>2</sup> Shota Yamashita,<sup>3</sup> and Kazuno Arai<sup>3</sup>

Received 14 February 2012; accepted 15 February 2012; published 15 March 2012.

[1] In Rikuzentakata City, Ostracode assemblages in sediment deposited by the Tohoku-Oki earthquake and tsunami of 11 March 2011 revealed that the sediment was derived from the seafloor from at least 9 m water depth, and was transported inland more than 1 km. The tsunami wave height at this location was higher than 10 m. Four hundred fifty seven modern ostracode assemblages were used in the modern analogue technique to estimate the depth source of the tsunami deposited assemblages. The application of this method to paleo-tsunami deposits may provide insight into past tsunami wave height and potentially earthquake slip and magnitude. **Citation:** Tanaka, G., H. Naruse, S. Yamashita, and K. Arai (2012), Ostracodes reveal the sea-bed origin of tsunami deposits, *Geophys. Res. Lett.*, 39, L05406, doi:10.1029/2012GL051320.

### 1. Introduction

[2] Holocene tsunami events have been recognized from sedimentological, micropaleontological, geological, and historical evidence [e.g., Nelson *et al.*, 2008; Sawai *et al.*, 2009]. The terrestrial or marine origin of sediments in the geologic record must be ascertained in order to distinguish between deposits from tsunami events and deposits from other processes. Micropaleontological evidence can be used to determine sediment source [e.g., Jenkins, 1993]. For example, the inundation of terrestrial environments by marine waters during Holocene tsunami events has been recognized by examining marine phytoplankton assemblages [Sawai *et al.*, 2004; Cisternas *et al.*, 2005].

[3] Ostracodes are small crustaceans (0.3–30 mm long) with calcified valves that are adapted to practically every aquatic environment [e.g., Horne *et al.*, 2002]. Thus, fossilized valves are an important paleoenvironmental indicator, particularly with regard to Holocene oceanographic, climatologic, and geologic events [e.g., Boomer *et al.*, 2005; Hussain *et al.*, 2010; Ruiz *et al.*, 2010]. Marine podocopid ostracodes are exclusively benthic crustaceans that are abundant in marine sediments [e.g., Horne *et al.*, 2002]. Furthermore, most species show regional endemism, therefore they can serve as an important indicator of local bottom-water environment. Their valves can behave like sediment grains in the water column [Ikeya and Cronin, 1993].

[4] The modern analog technique (MAT) statistical method is used to infer paleoenvironmental conditions by comparing fossil ostracode assemblages in terrestrial sediments with similar assemblages in modern environments

[Ikeya and Cronin, 1993]. In this paper we estimate the water depth of tsunami excavated ostracode assemblages from the Japan 2011 earthquake and tsunami at Rikuzentakata city.

### 2. Materials and Methods

[5] MAT is a method for estimating paleoenvironmental conditions by comparing modern and fossil assemblages [Overpeck *et al.*, 1985]. The squared chord distance (SCD), which is a measure of dissimilarity between a fossil assemblage and a modern assemblage, is calculated as follows:

$$SCD = \sum_{i=1}^n (Fp_i^{1/2} - Mp_i^{1/2})^2$$

where  $Fp_i$  is the  $i$ th number of the specified fossil species, and  $Mp_i$  is the  $i$ th number of the specified modern species.

[6] Ikeya and Cronin [1993] used MAT to estimate the paleoenvironments of Pliocene and Pleistocene formations in Japan by comparing fossil ostracode assemblages with 273 recent samples from the seas around Japan. To examine the source origin of the Japan 2011 Rikuzentakata City tsunami deposits, we used a total of 476 ostracode samples, including 367 recent species from Japan and its adjacent seas, from the published sources [Zhou, 1993; Tsukawaki *et al.*, 1998; Yamane, 1998; Ozawa *et al.*, 1999; Tsukawaki *et al.*, 2000; Yasuhara and Irizuki, 2001; Nakao and Tsukagoshi, 2002; Irizuki *et al.*, 2006; Tanaka, 2008] and the modern analog data matrix from the U.S. Geological Survey [Ikeya and Cronin, 1993]. The SCD values are computed using the Microsoft 2007 with VBA software. The SCD values can range from 0 to 2.0. A value of 0 indicates that the fossil assemblage is completely concordant with the modern population with which it is being compared.

[7] We compared modern ostracode assemblages from 476 sampled stations around Japan (small black plots in Figure 1a) with ostracode assemblages recovered from sediments deposited by the Tohoku-Oki earthquake generated tsunami of 11 March 2011. Our study area is about 4 km southeast of Rikuzentakata City (samples, collected on 25 April; Figure 1b and Table 1). A total of 50 sediment samples were collected in the study area, and 21 samples were used for this study.

### 3. Results and Discussion


[8] Although we found ostracode assemblages from 5 samples, only one sample (sample c) consisted of more than 100 individuals. So, we only deal with sample c in the following discussion.

[9] The assemblage of sample c is characterized by inner bay species, such as *Bicornucythere bisanensis* (Figure 1c),

<sup>1</sup>Gunma Museum of Natural History, Tomioka, Japan.

<sup>2</sup>Department of Geology and Mineralogy, Kyoto University, Kyoto, Japan.

<sup>3</sup>Department of Science, Chiba University, Chiba, Japan.



**Figure 1.** (a) Map of Japan showing the study area location and the epicenter of the Tohoku-Oki earthquake of 11 March 2011 and the black dots represent the 476 modern ostracode samples used in the MAT analysis. (b) Detailed map of the study area. Circled letters a–e show the sampling sites where ostracode specimens were found (see Table 1 for a complete list although only sample c is analyzed herein). Scanning electron microscope images of characteristic species recovered from sample c: (c) *Bicornucythere bisanensis* (preserved soft parts shown in yellow); (d) *Nipponocythere bicarinata*; (e) *Spinileberis quadriaculeata*; (f) *Cytheromorpha acupunctata*.

*Nipponocythere bicarinata* (Figure 1d), *Spinileberis quadriaculeata* (Figure 1e), and *Cytheromorpha acupunctata* (Figure 1f). This assemblage also contains *Aurila corniculata* and *Xestoleberis hanaii* which also occur on sediment substrates that support plant growth (seaweed, seagrass) [Kamiya, 1988]. The two species indicate derivation from within the photic (shallow) zone (Table 1). *Pontocythere subjaponica* and *Semicytherura miurensis* has been found at the shallow, sandy bottom environments

[Kamiya, 1988]. Living *Neonesidea oligodentata* has been found at calcareous algae of the intertidal zone [Smith et al., 2005]. Some of the ostracode valves (18%) of sample c were well preserved (Figures 1d–1f) and translucent. Soft parts were preserved in one *B. bisanensis* specimen (Figure 1c) which implied immediate derivation from a biocoenosis. However, many of the ostracode valves (82%) were opaque and fragmented, indicating that the ostracode assemblage in sample c was derived from another part [Zhou, 1995] on the seafloor. Brouwers [1988] discussed autochthonous or allochthonous ostracode assemblages from recent sediments off Alaska, USA. She mentioned that the assemblage is identified as allochthonous when the ratio of the number of adult valves and that of juvenile valves in a sample is 1:3 to 1:5. Table 1 shows that the ratio of adult and juvenile of sample c is 1:3. Thus, the ostracode assemblages from our sample c is identified as autochthonous based on Brouwers' [1988] opinion. This is supported by the observation that 27% of total individuals from sample c were articulated carapace. To summarize, the ostracode assemblage in tsunami deposit appears autochthonous from the viewpoint of the ratio of adult and juvenile individuals in a sample. This characteristic ostracode assemblage and the preservation of soft parts are useful when we identify the tsunami deposit from the (deep) older sediments. Thus, it is appropriate to use MAT to compare the sample c assemblage with modern ostracode assemblages from around Japan.

[10] By applying MAT, we determined that the ostracode assemblage in sample c was most similar to the ostracode assemblage of sample OK 28 from Osaka Bay, which was collected in 9 m of water depth (Figure 1a and Table S1 in the auxiliary material).<sup>1</sup> Table S1 shows that the valves of SCD of five samples were calculated <0.8, with depths ranging from 9 m to 30 m. Among them, the most closely matched sample was derived from a water depth at 9 m. The frequency distribution of water depths of 476 samples are shown in Figure S1. The number of samples used the MAT which were collected from <10 m water depths ( $n > 20$ ) is larger than that from 10 m to 20 m water depths. This result indicates that the database of the MAT used in this study is not artificially affected, for example by the difficulty of collecting sediment less than 10 m from a boat. Thus, sample c was derived from at least a depth of 9 m.

[11] Therefore, we infer that the sediments of sample c were derived from the seafloor at least 9 m water depth and deposited inland more than 1 km from the sea coast by the tsunami of 11 March 2011 (Figure 1b). These results suggest that the tsunami waves generated by the 2011 Tohoku-Oki earthquake eroded the seafloor under at least 9 m of water.

[12] Thus, the wave height of the tsunami and the sediment water depth source were similar. This suggests a relationship between tsunami wave height and wave excavation depth which is then translated onshore and preserved in the ostracode assemblage of tsunami deposits, although tsunami wave heights are generally influenced by many factors, such as width, depth and shape of a baymouth, dip, width, length and shape of the aggradation lowland [Mizutani, 2011].

<sup>1</sup>Auxiliary materials are available in the HTML. doi:10.1029/2012GL051320.

**Table 1.** Ostracode Assemblages (Raw Numbers) Discovered From 5 Tsunami Deposit Samples in Rikuzentakata City

Species	a					b					c					d					e				
	ac	av	jc	jv	to	ac	av	jc	jv	to	ac	av	jc	jv	to	ac	av	jc	jv	to	ac	av	jc	jv	to
<i>Ambtonia obai</i>											1	0	2	0	3	0	0	0	1	1					
<i>Angulicytherura miii</i>	0	1	0	1	1																				
<i>Aurila corniculata</i>											1	4	1	8	14	2	0	3	6	11					
<i>Bicornucythere bisanensis</i>											0	6	2	28	36	0	5	0	12	17					
<i>Bythoceratina hanaii</i>											0	0	0	2	2										
<i>Bythoceratina cf. hanaii</i>											0	0	0	2	2										
<i>Callistocythere japonica</i>											0	0	0	1	1										
<i>Callistocythere undulatifacialis</i>											0	0	0	2	2										
<i>Coquimba ishizakii</i>											0	1	0	2	3										
<i>Cornucoquimba tosaensis</i>											0	1	0	2	3						1	0	0	0	1
<i>Cytherois nakanoumiensis</i>											1	0	0	1	2										
<i>Cytheromorpha acupunctata</i>											4	0	1	1	6	2	0	1	0	3					
<i>Hemicytherura kajiyamai</i>											3	0	0	0	3	1	0	0	0	1					
<i>Howeina leptocytheroidea</i>											3	0	0	0	3	0	1	0	0	1					
<i>Kobayashiina donghaiensis</i>											0	0	0	2	2										
<i>Loxoconcha epeterseni</i>											0	1	0	0	1						0	0	0	1	1
<i>Loxoconcha japonica</i>											0	1	0	1	2										
<i>Loxoconcha ozawai</i>											0	2	0	0	2										
<i>Loxoconcha uranouchiensis</i>											0	0	0	3	3	0	0	0	1	1					
<i>Neonesidea oligodentata</i>											0	0	1	8	9	0	0	1	1	2					
<i>Nipponocythere bicarinata</i>											0	3	3	2	8	0	1	0	0	1					
<i>Parakrithella pseudadonta</i>											1	0	0	0	1										
<i>Pistocythereis bradyformis</i>											1	0	0	1	2										
<i>Pontocythere subjaponica</i>						0	1	0	0	1	3	0	3	0	6	0	1	0	1	2	0	1	0	3	4
<i>Schizocythere kishinouyei</i>																0	0	0	2	2					
<i>Semicytherura miurensis</i>											5	0	0	0	5										
<i>Spinileberis quadriculeata</i>											4	4	2	6	16	9	1	0	6	16					
<i>Xestoleberis hanaii</i>											0	2	0	17	19	0	1	0	2	3	0	0	0	1	1
<i>Xestoleberis sagamiensis</i>											0	0	0	1	1										
<i>Xestoleberis setouchiensis</i>																1	0	1	0	2					
Sub-total (Adult carapace)	0				0						27					15					1				
Sub-total (Adult valve)		1					1					25					10					1			
Sub-total (Juvenile carapace)			0					0					15					6					0		
Sub-total (Juvenile cvalve)				0					0					90					32					5	
Number of individual(s)					1					1					157					63					7
Carapace/total			0					0					0.27					0.33					0.14		
Adult/total			100					100					0.33					0.40					0.29		

[13] **Acknowledgments.** I (G.T.) dedicate this paper to the late N. Ikeya. We thank Robin J. Smith and Gary Parker for reviewing and discussing the early draft. Two editors from ELSS, Inc. edited the early manuscript. Our paper was extremely improved by David Horne, an anonymous reviewer, and the editor of GRL. This research was funded by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology, Government of Japan (21740370 to G. T.).

## References

- Boomer, I., U. V. Grafenstein, F. Guichard, and S. Bieda (2005), Modern and Holocene sublittoral ostracod assemblages (Crustacea) from the Caspian Sea: A unique brackish, deep-water environment, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 225, 173–186, doi:10.1016/j.palaeo.2004.10.023.
- Brouwers, E. M. (1988), Sediment transport detected from the analysis of ostracod population structure: An example from the Alaskan continental shelf, in *Ostracoda in the Earth Science*, edited by P. De Deckker, J.-P. Colin, and J.-P. Peypouquet, pp. 231–244, Elsevier, Amsterdam.
- Cisternas, M., et al. (2005), Predecessors of the giant 1960 Chile earthquake, *Nature*, 437, 404–407, doi:10.1038/nature03943.
- Horne, D. J., A. Cohen, and K. Martens (2002), Taxonomy, morphology and biology of Quaternary and living Ostracoda, in *The Ostracoda: Applications in Quaternary Research*, edited by J. A. Holmes and A. R. Chivas, pp. 5–36, AGU, Washington, D. C., doi:10.1029/131GM02.
- Hussain, S. M., S. P. Mohan, and M. P. Jonathan (2010), Ostracoda as an aid in identifying 2004 tsunami sediments: A report from SE coast of India, *Nat. Hazards*, 55, 513–522, doi:10.1007/s11069-010-9543-4.
- Ikeya, N., and T. M. Cronin (1993), Quantitative analysis of Ostracoda and water masses around Japan: Application to Pliocene and Pleistocene paleoceanography, *Micropaleontology*, 39, 263–281, doi:10.2307/1485900.
- Irizuki, T., H. Takata, and K. Yamada (2006), Recent Ostracoda from Urauchi Bay, Kamikoshiki-jima Island, Kagoshima Prefecture, southwestern Japan, *Laguna*, 13, 13–28.
- Jenkins, J. M. (1993), *Applied Micropaleontology*, Kluwer Acad., Dordrecht, Netherlands.
- Kamiya, T. (1988), Morphological and ethological adaptations of Ostracoda to microhabitats in *Zostera* beds, in *Evolutionary Biology of Ostracoda: Its Fundamentals and Applications*, edited by T. Hanai, N. Ikeya, and K. Ishizaki, pp. 303–318, Kodansha, Tokyo, doi:10.1016/S0920-5446(08)70191-2.
- Mizutani, H. (2011), M9 earthquake, tsunami, nuclear accident, *Newton*, 31, 18–117.
- Nakao, Y., and A. Tsukagoshi (2002), Brackish-water Ostracoda (Crustacea) from the Obitsu River Estuary, central Japan, *Species Diversity*, 7, 67–115.
- Nelson, A. R., Y. Sawai, A. Jennings, L. Bradley, B. Sherrod, J. Sabean, and B. P. Horton (2008), Great-earthquake paleogeodesy and tsunamis of the past 2000 years at Alsea Bay, central Oregon coast, USA, *Quat. Sci. Rev.*, 27, 747–768, doi:10.1016/j.quascirev.2008.01.001.
- Overpeck, J. T., T. Webb III, and I. C. Prentice (1985), Quantitative interpretation of fossil pollen spectra: Dissimilarity coefficients and the method of modern analogs, *Quat. Res.*, 23, 87–108, doi:10.1016/0033-5894(85)90074-2.
- Ozawa, H., K. Ikehara, and H. Katayama (1999), Recent ostracode fauna in the northeastern part of the Japan Sea, off northwestern Hokkaido, in *Comprehensive Study on Environmental Changes in the Western Hokkaido Coastal Area and Study on Evolution of Marine Active Faults*, edited by K. Ikehara and Y. Okamura, pp. 103–117, Geol. Surv. of Jpn., Tsukuba.

- Ruiz, F., M. Abad, L. M. Cáceres, J. R. Vidal, M. I. Carretero, M. Pozo, and M. L. González-Regalado (2010), Ostracods as tsunami tracers in Holocene sequences, *Quat. Res.*, *73*, 130–135, doi:10.1016/j.yqres.2009.08.005.
- Sawai, Y., K. Satake, T. Kamataki, H. Nasu, M. Shishikura, B. F. Atwater, B. P. Horton, H. M. Kelsey, T. Nagumo, and M. Yamaguchi (2004), Transient uplift after a 17th-century earthquake along the Kuril subduction zone, *Science*, *306*, 1918–1920, doi:10.1126/science.1104895.
- Sawai, Y., et al. (2009), A periodic recurrence of geologically recorded tsunamis during the past 5500 years in eastern Hokkaido, Japan, *J. Geophys. Res.*, *114*, B01319, doi:10.1029/2007JB005503.
- Smith, R. J., T. Kamiya, D. J. Horne, and A. Tsukagoshi (2005), Evaluation of a new character for the phylogenetic analysis of Ostracoda: The podocopan maxillular branchial plate, *Zool. Anz.*, *243*, 139–153, doi:10.1016/j.jcz.2004.07.005.
- Tanaka, G. (2008), Recent benthonic ostracod assemblages as indicators of the Tsushima warm current in the southwestern Sea of Japan, *Hydrobiologia*, *598*, 271–284, doi:10.1007/s10750-007-9162-6.
- Tsukawaki, S., T. Kamiya, H. Ozawa, and M. Kato (1998), Preliminary results on the sediment samplings during the R. V. Tansei-maru Cruise KT96-17 Leg 2 in the southwestern part of the Japan Sea—Sediments, benthic foraminifers and ostracodes, *Bull. Jpn. Sea Res. Inst.*, *29*, 67–89.
- Tsukawaki, S., H. Ozawa, H. Domitsu, T. Kamiya, M. Kato, and M. Oda (2000), Preliminary results from the R. V. Tansei-maru Cruise KT 98-17 in the southwestern marginal part of the Japan Sea—Sediments, benthic and planktonic foraminifers, and ostracodes, *Bull. Jpn. Sea Res. Inst.*, *31*, 89–119.
- Yamane, K. (1998), Recent ostracode assemblages from Hiuchi-nada Bay, Seto Inland Sea of Japan, *Bull. Ehime Pref. Sci. Mus.*, *3*, 81–91.
- Yasuhara, M., and T. Irizuki (2001), Recent Ostracoda from the north-eastern part of Osaka Bay, southwestern Japan, *J. Geosci. Osaka City Univ.*, *44*, 57–95.
- Zhou, B. (1993), Ostracode fauna in the Pacific off southwest Japan, PhD thesis, Dep. Geol. and Mineral., Kyoto Univ., Kyoto, Japan.
- Zhou, B. (1995), Recent ostracode fauna in the Pacific off southwest Japan, *Mem. Fac. Sci. Kyoto Univ. Ser. Geol. Mineral.*, *47*, 21–98.
- 
- K. Arai and S. Yamashita, Department of Science, Chiba University, 1-33, Yayoi-cho, Inage-ku, Chiba 263-8522, Japan.
- H. Naruse, Department of Geology and Mineralogy, Kyoto University, Kitashirakawa-oiwake-cho, Sakyo-ku, Kyoto 606-8502, Japan.
- G. Tanaka, Gunma Museum of Natural History, 1674-1 Kamikuroiwa, Tomioka, Gunma 370-2345, Japan. (tanaka@gmnh.pref.gunma.jp)